pain relief was required earlier in these nine children than in the children who had alternative intraoperative analgesia (n=9) (3.3 h v 7.6 h).

A conflict therefore arises over whether we should persist with tonsillar block and improve our performance even though we know that some children will be in pain while we are improving or whether we should abandon it for the method that gives better analgesia in our hands. We would now have difficulty in gaining the approval of an ethical committee for a trial of tonsillar block, so such a trial would not be included in an evidence based database even though it would answer a useful question. We also thought that parents would not consent to a trial investigating a method that we had shown to be inferior. This means that the question of pain relief may not be amenable to investigation with the criteria that have been imposed for evidence based medicine. Many more questions of interest would also be unanswerable.

Side effects of drugs come squarely into this category. Parke *et al*'s report of deaths after administration of propofol in children did not meet the criteria for inclusion in an evidence based database.² In our hospital an early death occurred after administration of propofol. We were told that there were so many confounding factors in our case that no inference was possible: Parke *et al*'s report certainly changed our practice. We conclude that some questions cannot be decided by the rigours of evidence based medicine and that, for others, answers are provided only after an unacceptable delay.

We challenge Frank Davidoff and colleagues to apply their criteria for evidence based medicine to the change of practice that they are advocating: is there any good evidence that evidence based practice works? The issue of the *BMJ* containing Davidoff and colleagues' editorial also contains a letter that investigates use of the Cochrane database.³ A more relevant question, however, is whether the outcome data for doctors who use the Cochrane database are better or worse than the data for those who do not.

Although evidence based medicine is a good way of deciding some issues, it is not the only way forward. If we use it as the only way forward we should recognise that we will not be able to answer some questions for which we will need to know the answers.

OLIVER DEARLOVE
Consultant anaesthetist

A SHARPLES
Consultant anaesthetist

K O'BRIEN
Senior registrar

C DUNKLEY
Registrar

Anaesthetic Department,
Royal Manchester Children's Hospital,
Pendlebury,
Manchester M27 1HA

1 Davidoff F, Haynes B, Sackett D, Smith R. Evidence based medicine. *BMJ* 1995;310:1085-6. (29 April.)

2 Parke TJ, Stevens JE, Rice ASC, Greenaway CL, Bray RJ, Smith PJ, *et al*. Metabolic acidosis and fatal myocardial failure after propofol infusion in children: five case reports. *BMJ* 1992;305: 613-6.

3 Hyde C. Who uses the Cochrane pregnancy and childbirth database? *BMJ* 1995;310:1140-1.

No guidance is provided for situations for which evidence is lacking

EDITOR,—The paradigm of evidence based medicine provides clear guidance for making decisions about the delivery of health care in populations.^{1,2} Therefore, it must be qualified so that it can be used for managing specific patients.

Evidence based medicine finds answers only to those questions open to its techniques; randomised controlled trials are its capstone. Other types of evidence, including natural observations and pathophysiological principles, are ranked lower and heavily discounted. Nevertheless, for many

rare disorders and an increasing number of subgroups of patients we will never have higher levels of evidence. Evidence based medicine does not provide guidance when one is trying to make sense of situations for which better types of evidence are lacking. Even when we have better evidence it is difficult to apply it to a particular patient. When one treatment is shown to be better than another on a population basis this does not mean that it is the best treatment for the patient. Only when we cannot reliably predict which patient will benefit from each treatment option should we take the general conclusion for the population and apply it to the individual patient.³ Establishing predictive assays and baseline variables to stratify for risk and benefit may help to match treatment options to subgroups of patients. The evidence for this, however, is likely to be of a lower level, partly because of increasingly small sample sizes.

Furthermore, we present treatment options to a patient to elicit informed consent. Some patients may prefer to receive a treatment that has been classified as less efficacious by evidence based medicine. Some doctors may be willing to give the treatment, and some taxpayers may be willing to pay for it, to empower the patient by offering a meaningful choice. Also, patients have to compare different procedures of techniques and trade off the chances of benefits and of toxicities, the severities of outcomes, and their temporal relationships. Each patient will do this slightly differently and may prefer a treatment option that evidence based medicine and its derivatives (practice guidelines⁴ and economic analyses) suggest should be abandoned.

We enjoy using evidence based medicine in our practice and research efforts but believe that it fails to address how individual, public, and social choices are made. It is useful in the design and appraisal of studies, which can provide us with quality data about multiple treatment options. While high quality evidence based medicine definitely informs us, however, it cannot make decisions for our patients.

GLENN W JONES
Consultant oncologist
STEPHEN M SAGAR
Consultant oncologist

Hamilton Regional Cancer Centre,
Hamilton,
Ontario L8V 5C2,
Canada

1 Rosenberg W, Donald A. Evidence based medicine: an approach to clinical problem solving. *BMJ* 1995;310:1122-6. (29 April.)

2 Evidence-Based Medicine Working Group. Evidence-based medicine: a new approach to teaching the practice of medicine. *JAMA* 1992;268:2420-5.

3 Charlton BG. Mega-trials: methodological issues and clinical implications. *J R Coll Physicians Lond* 1995;29:96-100.

4 Browman GP, Levine MN, Mohide EA, Hayward RSA, Pritchard KI, Gafni A, *et al*. The practice guidelines development cycle: a conceptual tool for practice guidelines development and implementation. *J Clin Oncol* 1995;13:502-12.

Quality cannot always be quantified

EDITOR,—William Rosenberg and Anna Donald examine some of the disadvantages of evidence based medicine but, for each "threat," argue that an overall "opportunity" to improve existing practice ensues (with the implication that the debate is more polarised than perhaps it is).¹ One important disadvantage is not addressed, though it is encapsulated in another context earlier in the article where the effectiveness of evidence based medicine is considered: "the approach is difficult to evaluate . . . since many [outcomes] are difficult to quantify."

Surely this is a problem with evidence based medicine. Most outcomes of health care are multifactorial, and it is mainly those outcomes that lend themselves to direct quantitative assessment that are used in randomised controlled trials—measuring the measurable. To extrapolate from

the example given in the article, it is important to consider all the effects on a patient of initiating a lifelong course of warfarin; these include the inconvenience of daily treatment and regular monitoring and review; disruption of existing and future drug treatment because of drug interactions; and potential psychological morbidity resulting from imposition of the sick role by treatment of an asymptomatic condition. These outcomes cannot be measured, but the risk of haemorrhage and reduction in the risk of stroke can be, so these are the figures on which decisions are to be based, argue the authors.

Where gaps in evidence are likely to exist we are exhorted to surmount the problem by attempting to answer only questions "for which there is likely to be good evidence" and awaiting the results of future research. There are no suggestions on how our practice relating to unmeasurable aspects is to be guided.

An analogy might be where hospital managers are tempted to highlight waiting time for patients at outpatient clinics as an indicator of the clinics' performance. As doctors and patients we believe that the content of the eventual consultation is more important than the waiting time, but the measurement is easy and relatively precise. Neither of the implicit assumptions—that patients' satisfaction equates with their waiting time and that quality of health care given equates with patients' satisfaction—can be made.

The principle of critical appraisal is laudable in all branches of higher education and professional practice and may have been deficient in medical education and practice. The wholesale adoption of evidence based medicine, given the current limitations of comprehensive outcome measures, is premature. A gradual and integrated introduction may be more appropriate.

BLAIR H SMITH
Lecturer in general practice

Department of General Practice,
University of Aberdeen,
Foresterhill Health Centre,
Aberdeen AB9 2AY

1 Rosenberg W, Donald A. Evidence based medicine: an approach to clinical problem solving. *BMJ* 1995;310:1122-6. (29 April.)

Journal of Evaluation in Clinical Practice will start publication in September

EDITOR,—The "Editor's choice" in the issue of 29 April welcomes the publication of the new journal, *Evidence Based Medicine*, which is described more fully in Frank Davidoff and colleagues' editorial.¹ In addition, Aneez Esmail reviews a new textbook entitled *Evidence-based General Practice*.² These are important contributions to the inexorable development of evidence based health care.

I wish to draw attention to a new clinical journal that is similarly dedicated to evidence based medicine but within the broader context of multidisciplinary clinical care and analysis. The *Journal of Evaluation in Clinical Practice*, whose first edition will be published in September, will focus on systematic reviews of research on clinical effectiveness; the implementation of evidence based care in routine clinical practice; systematic medical and clinical audit; methodologies for quantifying clinical benefit from audit and after the implementation of evidence based care; systematic reviews of the current status of audit and its development in medicine, nursing, midwifery, and the range of professions allied to medicine; and the functional integration of clinical audit, research and development, and quality assurance in health care organisations.

A textbook focusing on such central issues in modern medicine, *Effective Clinical Practice*, will be published in November³ and is concerned with clinical effectiveness, clinical appropriateness, the efficiency of the delivery of clinical services, the