produce a tarry stool.² If there are, as it would be reasonable to expect, a fairly large number of small gastrointestinal bleeds then those who were alerted about what to look for could expect to detect a few more. Aspirin prophylaxis is only recommended to be undertaken on medical advice. Presumably most doctors giving such advice would warn the patient of possible gastrointestinal bleeding and give advice on reporting it, if it occurs. The control groups would not be expected to have had similar advice.

John Weil and colleagues could throw light on whether this confounding factor could be material to their results, by showing the breakdown of melaena and haematemesis in each of the three groups they studied.

GORDON R FRYERS
Honorary president

European Aspirin Foundation,
London NW14 6DR

1 Weil J, Colin-Jones D, Langman M, Lawson D, Logan R, Murphy M, *et al.* Prophylactic aspirin and risk of peptic ulcer bleeding. *BMJ* 1995;310:827-30. (1 April.)

2 Daniel WA, Egan S. The quantity of blood required to produce a tarry stool. *JAMA* 1939;113:2232.

Author's reply

EDITOR,—We cannot check on the proportions of overt haematemesis to melaena without going back to original records. We think it unlikely that Dr Fryer's explanation will account for the trends observed. If doctors had warned patients to watch for the ill effects of aspirin on the upper gut then we would not have expected to see the steadily increasing risk with aspirin dose. Also, very low dose aspirin was believed to be safe and therefore precautions are unnecessary, so warnings by doctors about the possible hazards of low dose aspirin will have been unlikely.

MICHAEL LANGMAN
Professor of medicine

Queen Elizabeth Hospital,
University of Birmingham,
Birmingham B15 2TH

Treatment of childhood asthma

EDITOR,—The study of Fiona Bryce and colleagues is elegantly designed to test the effects of an audit facilitator on treatment and diagnosis of childhood asthma by randomising within practices.¹

It is a pity, however, given the large number of time related significant changes in treatment found for both intervention and control groups, that the study did not have a control group of children from practices where no intervention was made. This would better confirm the conclusion that intervention increased the likelihood of diagnosis of childhood asthma and planned review but did not make major changes in treatment. From the evidence presented, time related changes for both groups (decrease in use of oral bronchodilators, theophyllines, antibiotics, and cough linctus; increase in inhaled and oral steroids) suggest that practice prescribing was also influenced for children in the control group. In support of the conclusion that practices were changing habits at this time, it may be of interest that over much the same period, in the neighbouring Grampian and Highland regions, a similar decrease in use of oral bronchodilators and increase in use of inhaled steroids was found for adults.²

None the less, we all need to remember that studies which introduce management interventions for (randomly) selected patients within a practice are not as "clean" as clinical trials of procedures or drug treatment. The same doctors and nurses whose behaviour is being changed in relation to the intervention group are also managing the control group. Without a completely

independent control group it is difficult to interpret findings if few differences are found between intervention and control.

LIESLOS MAN
Senior research fellow

Chest Clinic,
City Hospital,
Aberdeen AB9 8AU

SUE ROSS
Lecturer in health care research

Department of General Practice,
University of Glasgow,
Glasgow

1 Bryce FP, Neville RG, Crombie IK, Clark RA, McKenzie P. Controlled trial of an audit facilitator in diagnosis and treatment of childhood asthma in general practice. *BMJ* 1995;310:838-42. (1 April.)

2 Ross S. Asthma medication use in Northern Scotland 1989-91. *Eur Respir J* 1993;6:341S.

Mineralocorticoid effects of high dose hydrocortisone

EDITOR,—In their recent Lesson of the Week B H Ramsahoye and colleagues highlight the mineralocorticoid effect of high dose hydrocortisone as the principal factor that resulted in the fall in serum potassium concentrations (2.9 to 2.0 mmol/l) in their patient.¹ The episodes of ventricular fibrillation which followed were attributed to this fall in serum potassium concentration. Whereas intravenous hydrocortisone certainly contributed to the reduction in serum concentration of potassium, we suggest that other factors responsible for this change were a combination of inadequate supplementation, inappropriate cessation of potassium supplementation (at a serum concentration of 2.9 mmol/l), and, perhaps most importantly, significant continued urinary losses of potassium in the face of magnesium depletion.

The authors suggest that a reduction in serum potassium concentration of 1 mmol/l represents depletion in total body potassium of 100-200 mmol. Other workers would put the figure at twice this, with serum potassium concentrations of 3.0 mmol/l and 2.0 mmol/l representing total body potassium deficits of 350 and 700 mmol respectively.^{2,4} Thus, potassium supplementation of 40-80 mmol per day, in the face of such a large deficit and continuing losses, was inadequate. This case illustrates clearly the need for substantial intravenous potassium supplementation in patients with potassium depletion.

NAVEED A SATTAR
Registrar in pathological biochemistry
ALLAN GAW
Clinical lecturer

Department of Pathological Biochemistry,
University of Glasgow,
Royal Infirmary,
Glasgow G4 0SF

1 Ramsahoye BH, Davies SV, El-Gayani N, Sandeman D, Scanlon MF. The mineralocorticoid effects of high dose hydrocortisone. *BMJ* 1995;310:656-7. (11 March.)

2 Sterns RH, Cox M, Feig PU, Singer I. Internal potassium balance and the control of the plasma potassium concentration. *Medicine* 1981;60:339.

3 Stanaszek WF, Romankiewicz JA. Current approaches to management of potassium deficiency. *Drug Intelligence and Clinical Pharmacy* 1985;19:176-84.

4 Marino PL. *The ICU book*. Philadelphia: Lea and Febiger, 1991.

Interventions in OXCHECK study waste resources

EDITOR,—The OXCHECK study showed significant reductions in total cholesterol concentration (3.1%) and blood pressure (2.5/1.5 mm Hg) with health checks over three years.¹ The reductions in cholesterol concentration are in line with responses shown in previous controlled trials.² The authors

seem in some doubt about whether these effects are large enough to justify the cost of the health checks. This may be clearer when the projected benefits are considered in absolute rather than relative terms.

When the Framingham parametric model is used to estimate risk³ with the mean blood pressure and mean total cholesterol concentration in the OXCHECK study, assuming a mean age of 50 and with weighting for the proportion of smokers, the risk of coronary death in the control group is about 0.075% a year in non-diabetic men without electrocardiographic evidence of left ventricular hypertrophy. In the intervention group the estimated rate of coronary death is lowered to 0.066% a year. This is a reduction in risk of 12% in relative terms, but in absolute terms 11 000 men would need to have health checks for one coronary death to be prevented each year. For women the reduction in relative risk seems even more impressive, at 25%, but over 80 000 women would need health checks for one coronary death to be prevented each year. If the study population had only a single initial health check lasting one hour and no follow up visits, seven practice nurses would be employed full time for one year to prevent one coronary death in men, and an additional 41 would need to be employed to prevent one death on women.

It is important to advise opportunistic measures such as stopping smoking and increasing exercise, but whether the time and resources needed to implement these measures systematically, as tested in the OXCHECK study, can be justified is doubtful. It seems more appropriate to concentrate resources on lowering cholesterol with 3-hydroxy-3-methylglutaryl coenzyme A reductase inhibitors after myocardial infarction⁴ and in patients with other forms of coronary heart disease.⁵

I U HAQ
Research registrar
W W YEO
Lecturer
P R JACKSON
Senior lecturer
L E RAMSAY
Professor

Section of Clinical Pharmacology and Therapeutics,
University Department of Medicine and Pharmacology,
Royal Hallamshire Hospital,
Sheffield S10 2JF

1 Imperial Cancer Research Fund OXCHECK Study Group. Effectiveness of health checks conducted by nurses in primary care: final results of the OXCHECK study. *BMJ* 1995;310:1099-104. (29 April.)

2 Ramsay LE, Yeo WW, Jackson PR. Dietary reduction of serum cholesterol concentration: time to think again. *BMJ* 1991;303:953-7.

3 Anderson KM, Odell PM, Wilson PWF, Kannel WB. Cardiovascular disease risk profiles. *Am Heart J* 1990;121:293-4.

4 Scandinavian Simvastatin Survival Study Group. Randomised trial of cholesterol lowering in 4444 patients with coronary heart disease: the Scandinavian simvastatin survival study (4S). *Lancet* 1994;344:1383-9.

5 Oliver M, Poole-Wilson P, Shepherd J, Tikkanen MJ. Lower patients' cholesterol now. *BMJ* 1995;310:1280-1. (20 May.)

Assessing coronary risk

EDITOR,—Steven A Grover and colleagues compared estimates by doctors of gains in life expectancy as a result of a reduction in risk factors with those estimated from their computer model of prevention of coronary heart disease.¹ The model they used is based on logistic regression analyses, using data from the Framingham heart study, the Canada health survey, and Canadian life tables.² They calculated potential gains in life expectancy with intervention by applying to such a statistical model the levels of the risk factors before and after the intervention.

Three major problems are inherent in this approach. Firstly, the benefits of a reduction in risk factors may not be immediately apparent, since there is a substantial lag period before full benefits are achieved.³ Secondly, there may be irreversible damage to blood vessels or irreversible side effects of drugs. Finally, there is some

Intervention	For 40 year old man (coronary heart disease prevention model) ¹	For 45 year old man (meta-analysis application) ²
Stopping smoking	4.9	3.84 (3.68 to 4.04)
Reducing blood pressure	0.9	0.83 (-0.07 to 1.79)
Lowering cholesterol	1.2	-0.01 (-0.78 to 0.66)

suspicion, particularly with cholesterol lowering interventions, that adverse effects may be seen on non-cardiovascular mortality.⁴

I have used a different method to calculate potential gains in life expectancy with a reduction in risk factors, using observations from meta-analyses of intervention rather than "modelling out" the effect of the risk factor.⁵ The table shows the benefits calculated by Grover and colleagues using the coronary heart disease prevention model for a 40 year old man and compares these with figures I calculated using the data from meta-analyses of intervention for a 45 year old man. While the estimated benefits of stopping smoking and reducing blood pressure are of similar magnitude, the effect of cholesterol lowering treatment is much greater when the model of prevention of coronary heart disease is used, largely because of the three factors I have outlined.

Prevention of coronary heart disease needs a concerted attack on risk factors, with smoking way out in front as the most important. Possibly the newer drugs to treat hypercholesterolaemia will be shown to reduce cardiovascular mortality in relatively low risk subjects. In the meantime, it is important both for patients and for doctors not to overestimate likely benefits from treating 40 year old men free of coronary heart disease and with a serum cholesterol concentration of 6.2 mmol/l.

JOHN S YUDKIN
Professor of medicine

Academic Division of Medicine,
Whittington Hospital,
London N19 3UA

- 1 Grover SA, Lowenstein I, Esrey KL, Steinert Y, Joseph L, Abrahamowicz M. Do doctors accurately assess coronary risk in their patients? Preliminary results of the coronary health assessment study. *BMJ* 1995;310:975-8. (15 April.)
- 2 Grover SA, Abrahamowicz M, Joseph L, Brewer C, Coupal L, Suissa S. The benefits of treating hyperlipidemia to prevent coronary heart disease. *JAMA* 1992;267:816-22.
- 3 Law MR, Wald NJ, Thompson SG. By how much and how quickly does reduction in serum cholesterol concentration lower risk of ischaemic heart disease? *BMJ* 1994;308:367-72.
- 4 Davey Smith G, Pekkanen J. Should there be a moratorium on the use of cholesterol lowering drugs? *BMJ* 1992;304:431-4.
- 5 Yudkin JS. How can we best prolong life? The benefits of coronary risk factor reduction in diabetic and non-diabetic subjects. *BMJ* 1993;306:1313-8.

Authors' reply

EDITOR,—We agree with John S Yudkin that there are many problems inherent in a statistical modelling approach to estimate the benefits of a reduction in risk factors. We disagree with him, however, about whether there are only three problems and whether those he identifies are major.

Yudkin suggests that the first major problem is that the benefits of a reduction in risk factors may be apparent only after a substantial lag period. We agree that there may be a lag before the benefits of treatment are seen in primary prevention trials, and we have built a variable lag period into our model.¹ What is not clear is whether this is due to a biological delay associated with the benefits of treatment or simply a statistical artefact resulting from few fatal events among either treatment or control groups in the first year or two of a randomised clinical trial, which makes it difficult to appreciate the cumulative difference over the short term. Consistent with the second possibility is the recently published Scandinavian simvastatin survival study, in which Kaplan-Meier curves for major coronary events among treatment and

control groups diverged in less than one year while all cause mortality diverged at approximately 1.5 years.² Moreover, even if there is a two year lag for the benefits of treatment to manifest themselves, the net effect on long term life expectancy for a 40 year old man may be relatively small. For instance, when we use our model the difference in life expectancy is 1.23 years without a two year lag and 1.16 years with a lag.

We agree with Yudkin that the risk of coronary disease may not be totally reversible, particularly in the context of a reduction in blood pressure.³ None the less, we note that our estimates of the benefits of reducing blood pressure are nearly identical with his results based on a meta-analysis of outcomes of clinical trials.

Finally, we agree with the important concern regarding the potential adverse effects that may be associated with cholesterol lowering treatments. We have always emphasised that our model simulations have not been adjusted for a potential increased risk in non-coronary death associated with lipid lowering treatment.⁴ To assume, however, that the results of a meta-analysis can identify this relation without any a priori hypothesis or clear cut biological explanation is somewhat simplistic. For instance, there is no explanation of why the increase in traumatic death seen in some primary prevention trials is generally not seen in secondary prevention trials. The Scandinavian simvastatin survival, a large clinical trial, did not show any evidence of increased non-coronary death associated with lipid modification.

Finally, the focus of our paper was the accuracy of doctors' predictions of the reduction in coronary risk. Comparing their estimates with our computer model, we concluded that doctors may systematically overestimate the benefits of modification of coronary risk factors. If Yudkin is correct in stating that our computer model overestimates the benefits of modification of risk factors for some groups of people, then these conclusions are even more valid.

STEVEN A GROVER
Director
ILKA LOWENSTEIN
Research fellow

KATJA ESTREY
Research fellow
LAWRENCE JOSEPH
Biostatistician
MICHAL ABRAHAMOWICZ
Biostatistician

Division of Clinical Epidemiology,
Centre for the Analysis of Cost-Effective Care,
Montreal General Hospital,
Montreal,
Quebec,
Canada H3G 1A4

YVONNE STEINERT
Psychologist

Department of Family Medicine,
Sir Mortimer B Davis Jewish General Hospital,
Montreal,
Quebec,
Canada H3T 1E2

- 1 Grover SA, Gray-Donald K, Joseph L, Abrahamowicz M, Coupal L. Life expectancy following dietary modification or smoking cessation: estimating the benefits of a prudent lifestyle. *Arch Intern Med* 1994;154:1697-704.
- 2 Scandinavian Simvastatin Survival Study Group. Randomised trial of cholesterol lowering in 4444 patients with coronary heart disease: the Scandinavian simvastatin survival study (4S). *Lancet* 1994;344:1383-9.
- 3 Collins R, Peto R, MacMahon S, Herbert P, Fiebach NH, Eberlein KA, et al. Blood pressure, stroke, and coronary heart disease. Part 2. Short-term reductions in blood pressure: overview of randomised drug trials in their epidemiological context. *Lancet* 1990;335:827-38.
- 4 Grover SA, Abrahamowicz M, Joseph L, Brewer C, Coupal L, Suissa S. The benefits of treating hyperlipidemia to prevent coronary heart disease: estimating changes in life expectancy and morbidity. *JAMA* 1992;267:816-22.

Hospital security

EDITOR,—We wish to draw attention to an important incident that occurred in Princess Alexandra Hospital Trust, Harlow, partly because of the need for increased hospital security. The cardiac team was called to the paediatric ward when an asthmatic child had a respiratory arrest. The anaesthetics registrar ran towards the ward but found that the corridor was blocked by a door operated by a number code, which had been installed earlier that week. Not knowing the code and being unable to attract attention, the registrar took a circuitous route through another security door, for which he knew the code. The ward door, which was usually operated by a buzzer linked to a video camera, was being held open by a member of staff, who did not know that there were any other doors between the main hospital and the paediatric ward. Overall there was an estimated four minute delay before the child was attended by the cardiac team.

While recognising the importance of security for staff and patients, we urge those responsible for installing new systems to consider the implications for emergencies. Our hospital resolved this problem by using the same code on all doors. The code number is printed on stickers, which are attached to bleeps. Other possible solutions include a central override facility, an override button on each door that sounds an alarm indicating that security has been breached, and a system that uses swipe cards. The hospital resuscitation officer should be involved in the planning and running of such systems to prevent similar incidents occurring.

JEFFREY PHILLIPS
Registrar
MICHELLE SOSKIN
Senior registrar

Department of Anaesthesia,
Princess Alexandra Hospital Trust,
Harlow,
Essex CM20 1XQ

Handling scientific fraud

Pearce's editors were not to blame

EDITOR,—In his editorial on the Pearce affair Stephen Lock states that the review of the clinical trial was clearly inadequate.¹ He is amazed at the credulity that the *British Journal of Obstetrics and Gynaecology* showed in publishing the trial, on the basis of his observation that over three years Pearce purported to have collected 191 women with a syndrome so uncommon that a major referral centre (unnamed) saw only one or two new cases a month. I imagine that even editors of the *BMJ* sometimes have problems with superficial and uncritical referees' reports.

After receiving a clinical referee's report recommending publication of the paper Professor Chamberlain delegated editorial responsibility to me. At an editorial meeting in November 1993, attended by Professor Chamberlain and all four editors, I gave a detailed account of the paper, noting the relatively large number of cases (although the time span of their recruitment was not stated in the manuscript) and suggesting that the source must be a tertiary referral centre. Pearce confirmed that it was, and his assertion was accepted. His acknowledged skill in ultrasound examination could be expected to attract large numbers of women with recurrent miscarriages to his pregnancy loss clinic. The paper was then sent to a statistical referee, and among 11 enumerated points in his report was one asking the authors whether "all [his emphasis] women with RSM [recurrent spontaneous miscarriage] associated with PCOS [polycystic ovaries] in a particular time interval were invited to take part in the trial." The revised manuscript responded to the question and gave the time interval.