- 2 Sever P, Beevers G, Bulpitt C, Lever A, Ramsay L, Reid J, *et al.* Management guidelines in essential hypertension: report of the second working party of the British Hypertension Society. *BMJ* 1993;306:983-7.
- 3 Grimshaw JM, Russell IT. Achieving health gain through clinical guidelines. I. Developing scientifically valid guidelines. *Quality in Health Care* 1993;2: 243-8.
- 4 Grimshaw JM, Russell IT. Achieving health gain through clinical guidelines. II. Ensuring guidelines change medical practice. *Quality in Health Care* 1994;3:45-52.
- 5 Grimshaw JM, Russell IT. Effect of clinical guidelines on medical practice: a systematic review of rigorous evaluations. *Lancet* 1993;342:1317-22.
- 6 Grimshaw J, Freemantle N, Grimshaw J, Wallace S, Russell I, Hurwitz B, *et al.* Developing and implementing clinical practice guidelines. *Quality in Health Care* 1995;4:55-64.
- 7 NHS Management Executive. *Improving clinical effectiveness*. Leeds: Department of Health, 1993. (EL(93)115.)
- 8 Lomas J, Anderson GM, Domnick-Pierre K, Vayda E, Enkin MW, Hannah WJ. Do practice guidelines guide practice? The effect of a consensus statement on the practice of physicians. *N Engl J Med* 1989;321:1306-11.
- 9 Jones RH, Lydeard S, Dunleavy J. Problems with implementing guidelines: a randomised controlled trial of consensus management of dyspepsia. *Quality in Health Care* 1993;2:217-21.
- 10 Neville RG, Clark RC, Hoskins G, Smith B. National asthma attack audit 1991-2. *BMJ* 1993;306:559-62.
- 11 Davis DA, Thomson MA, Oxman AD, Haynes RB. Evidence for the effectiveness of CME. A review of 50 randomized controlled trials. *JAMA* 1992;268:1111-7.
- 12 Mugford M, Banfield P, O'Hanlon M. Effects of feedback of information on clinical practice: a review. *BMJ* 1991;303:398-402.
- 13 Lomas J, Enkin M, Anderson GM, Hannah WJ, Vayda E, Singer J. Opinion leaders versus audit and feedback to implement practice guidelines. *JAMA* 1991;265:2202-7.
- 14 Brook RH, Park RE, Chassin MR, Solomon DH, Keesey J, Koscoff J. Predicting the appropriate use of carotid endarterectomy, upper gastrointestinal endoscopy, and carotid angiography. *N Engl J Med* 1990;323: 1173-7.
- 15 Hicks NR. Some observations on attempts to measure appropriateness of care. *BMJ* 1994;309:730-3.
- 16 Ashton CM, Kuykendall DH, Johnson ML, Wun CC, Wray NP, Carr MJ, *et al.* A method of developing and weighting explicit process of care criteria for quality assessment. *Med Care* 1994;32:755-70.
- 17 Kessner DM, Kalk CE, Singer AA. Assessing health care quality—the case for tracers. *N Engl J Med* 1973;288:189-94.
- 18 Irvine D. Managing for quality in general practice. London: King's Fund Centre, 1990. (Medical audit series 2.)
- 19 Oxman AD. Checklists for review articles. *BMJ* 1994;309:648-51.
- 20 Donabedian A. The quality of medical care. Methods for assessing and monitoring the quality of care for research and for quality assurance. *Science* 1978;200:856-63.
- 21 Baker R, Khunti K, Lakhani M. *Monitoring diabetes*. Leicester: Eli Lilly National Clinical Audit Centre. University of Leicester, 1993.
- 22 Haines A, Jones R. Implementing findings of research. *BMJ* 1994;308:1488-92.
- 23 Baker R. Audit and standards in new general practice. *BMJ* 1991;303:32-4.
- 24 North of England Study of Standards and Performance in General Practice. Medical audit in general practice: effects on doctors' clinical behaviour and the health of patients with common childhood conditions. *BMJ* 1992;304: 1480-8.

(Accepted 1 May 1995)

Commentary: The perils of checklist medicine

Iona Heath

This paper has a logic and clarity that does not seem to belong in the messy world where I and my patients strive, often against daunting socioeconomic odds, to be "co-producers of health."¹ The honourable intentions of the authors are evident, but the notion of defining and using criteria with which purchasers can monitor compliance with agreed guidelines implies levels of coercion and control which are potentially destructive of the fragile good that is the doctor-patient relationship.² Guidelines are constructed from evidence from research derived from studies of populations and are predicated on the notion of a composite patient which may have little immediate relevance to the troubled person who presents in the consulting room.³ In hospitals, diseases stay and patients come and go; in general practice, patients stay and diseases come and go. Guidelines depersonalise individual patients and turn them into diseases even in general practice.

The authors give a single example of what a review criterion will look like in reality. As far as I understand it, the evidence from research is that excessive intake of alcohol increases the patient's risk of raised blood pressure. This becomes translated into a review criterion with the suggested wording "the records show that annually the patient has been advised to limit weekly alcohol intake to less than 21 units if male and less than 14 units if female." I have objections at two levels. Firstly, there is evidence of the role of excessive alcohol in the aetiology of hypertension, but I am not aware of evidence to support the contention that advising patients annually to limit their alcohol consumption improves the control of raised blood pressure, let alone that a note in the records to this effect has any impact. Surely, this is to confuse process with outcome. Secondly, patients may resent the imposition of the doctor's agenda for the consultation at the expense of their own. The doctor will feel compelled to give this advice repeatedly and increasingly mechanistically whether the patient is a longstanding alcoholic or a renowned teetotaler. A year is a short time in general practice, and patients who never drink will not take it kindly when their doctor reminds them yet again of the perils of excess. The patient will be concerned about both the memory and the sensitivity

of the doctor. At the other end of the range, an alcoholic patient often faces social and economic problems which render the risks of hypertension unimportant. Will he or she be helped by being repeatedly informed of the recommended limits?

Patients, perhaps even more than doctors, make many perverse decisions. They persist in smoking and drinking to excess despite being aware of the risks to their future health; they refuse to comply with medication and technology, such as hearing aids, of proved efficacy. This seems to suggest that health, narrowly defined as the absence of disease, is not the pre-eminent aspiration of human endeavour.⁴ Doctors, managers, and politicians need to recognise that self determination, dignity, and autonomy are important components of a wider definition of health and that personal and socioeconomic pressures can make longevity less desirable.

Guidelines and review criteria can contribute to the pursuit of quality in clinical care, but if politicians and purchasers seek to implement them without acknowledging the complexities of real clinical settings they may do more harm than good and consume huge resources in the process.⁵

- 1 Hart JT. *Feasible socialism: the National Health Service, past, present and future*. London: Socialist Health Association, 1994.
- 2 Willis J. *The paradox of progress*. Oxford: Radcliffe Medical Press, 1995.
- 3 Royal College of General Practitioners. *The development and implementation of clinical guidelines*. Exeter: Royal College of General Practitioners, 1995. (Report from general practice 26.)
- 4 Toon P. *What is good general practice?* Exeter: Royal College of General Practitioners, 1994. (Occasional paper 65.)
- 5 McKee M, Clarke A. Guidelines, enthusiasms, uncertainty, and the limits to purchasing. *BMJ* 1995;310:101-4.

Correction

Ethical debate: Financial ties as part of informed consent to postmarketing research

The contributions to this debate by R E Ferner and Malcolm VandenBurg and Ian Dews (24 June, pp 1660-3) contained an error in the references. The second author of the guidelines on safety assessment of marketed medicine (references 2 and 1 respectively) should be the Royal College of General Practitioners not the Royal College of Physicians.

Caversham Group
Practice, Kentish Town
Health Centre, London
NW5 2AJ
Iona Heath, general
practitioner