Niu Huang Wan was found to contain 47 mg mercury and Zhusha Aushen 659 mg mercury (in the form of cinnabar). Arsenic and lead were also detected. When taken together as prescribed this would have resulted in a daily intake of 706 mg of cinnabar. We are concerned that the use of this amount of mercuric sulphide in long term treatment is potentially hazardous and could lead to mercury intoxication. Kew *et al* reported symptoms of mercury poisoning in a patient after daily exposure to 180–252 mg cinnabar for four weeks.²

Cinnabar has been used for a long time in traditional medicines. The toxic effects of inorganic mercury are well recognised, but because of its insolubility it has been assumed that this compound would not be significantly absorbed from the gastrointestinal tract. However, a study by Yeon *et al* of the oral absorption of cinnabar in mice found a significant increase in mercury concentrations in the liver and kidney. Concomitant use of cinnabar and drugs containing bromides, sulphates, sulphides, nitrates, and iodine may enhance its toxicity by increasing the gastrointestinal absorption.³

We wish to bring the availability and use of these preparations to the attention of medical professionals and to emphasise the importance of obtaining full information on all the medications used by their patients. We are carrying out a traditional remedies surveillance project and would be interested in receiving reports of adverse reactions to herbal remedies and food supplements.

This study is supported by the Ministry of Agriculture, Fisheries and Food.

DEBBIE SHAW Research scientist	IVAN HOUSE Trace elements analyst
STOYKO KOLEV Toxicologist	VIRGINIA MURRAY Consultant toxicologist

Traditional Remedies Surveillance,
Medical Toxicology Unit,
Guy's and St Thomas's Hospital Trust,
London SE14 5ER

- 1 De Smet PAGM. Should herbal medicine-like products be licensed as medicines? *BMJ* 1995;310:1023. (22 April.)
- 2 Kew J, Morris C, Alhie A, Fysh R, Jones S, Brooks D. Arsenic and mercury intoxication due to Indian ethnic remedies. *BMJ* 1993;306:507-8.
- 3 Pharmacopoeia of the People's Republic of China. English ed. Guangzhou: Guangdong Science and Technology Press, 1992: 21-2.
- 4 Yeon TS, Lee AS, Lee HS. Absorption of mercuric sulphide following oral administration in mice. *Toxicology* 1986;41: 107-11.
- 5 Kang-Yum E, Oransky SH. Chinese patent medicine as a potential source of poisoning. *Vet Hum Toxicol* 1992;34:235-8.

recommended as the drug of choice for sedation on the basis of pharmacokinetic and pharmacodynamic properties⁴ and lends itself to a careful titration technique. Combinations of opioids and benzodiazepines are best avoided owing to the additive and synergistic effects on the cardiovascular and respiratory systems. It must be remembered that the antagonist drug flumazenil has a shorter half life than the benzodiazepines in common clinical use.

Local anaesthetic sprays have been implicated in the development of pneumonia after gastroscopy.⁵ Their use may be unnecessary in most diagnostic endoscopic procedures. Formal training of junior endoscopists in sedation techniques together with adequate supervision should ensure the rational use of the available drugs.

When single operator sedationists practise endoscopy the British Society of Gastroenterology recommends at a minimum that a nurse trained in endoscopy techniques and dedicated to patient care is present throughout. Supplemental oxygen and monitoring should ideally be continued throughout the period of recovery, and premature discharge to maximise turnover must be resisted. The employment of qualified staff who are able to monitor patients, interpret changes in the monitored variables, and start cardiopulmonary resuscitation when necessary may have implications for general practice endoscopy services, where it may prove difficult to ensure both safety and cost effectiveness.

We may be able to influence these worrying statistics by centralising services, ensuring proficiency of endoscopists, providing skilled help, and adopting the excellent guidelines already available.

IAN R APPADURAI Lecturer in anaesthetics	RAYMOND J DELICTA Research fellow in surgery
	P DECLAN CAREY Senior lecturer in surgery

University of Wales College of Medicine,
Cardiff CF4 4XN

- 1 Charlton JE. Monitoring and supplemental oxygen during endoscopy. *BMJ* 1995;310:886-7. (8 April.)
- 2 Buck N, Devlin HB, Lunn JN. *The report of a confidential enquiry into perioperative deaths*. London: Nuffield Provincial Hospitals Trust and King's Fund, 1987.
- 3 Bell GD, McCloy RF, Charlton JE, Campbell D, Dent NA, Gear MWL, *et al*. Recommendations for standards of sedation and patient monitoring during gastrointestinal endoscopy. *Gut* 1991;32:823-7.
- 4 Royal College of Surgeons of England Commission on the Provision of Surgical Services. *Report of the working party on guidelines for sedation by non-anaesthetists*. London: RCS, 1993.
- 5 Quine MA, Bell GD, McCloy RF, Charlton JE, Devlin HB, Hopkins A. Prospective audit of upper gastrointestinal endoscopy in two regions of England: safety, staffing, and sedation methods. *Gut* 1995;36:462-7.

valuable, and prior knowledge is needed to prevent misinterpretation. Many limitations of pulse oximetry have been described, including the fact that it is a "lag" monitor; that it measures oxygenation, not ventilation; and that the function of the oximeter is adversely affected by a variety of factors.⁴ Anaesthetists are required, as part of their training, to have detailed knowledge of all the monitoring devices they use on patients perioperatively, including pulse oximeters. A recent survey of junior doctors in specialties other than anaesthesia showed poor overall knowledge of the principles of pulse oximetry, its limitations, and its application to clinical practice.⁵ A failure to interpret oxygen saturation values correctly could be as dangerous as not having them available.

If monitoring techniques such as pulse oximetry are introduced widely this should be accompanied by an increase in training in the use of such devices for staff. Widening the remit of the national confidential inquiry into perioperative deaths to include endoscopy would certainly be a start in auditing the safety of current endoscopic practice, but perhaps the safest method of all would be to add endoscopy under sedation to the list of procedures that are monitored by anaesthetists.

MARK STONEHAM
Senior registrar

Department of Anaesthesia,
Stoke Mandeville Hospital,
Aylesbury HP21 8AL

- 1 Charlton JE. Monitoring and supplemental oxygen during endoscopy. *BMJ* 1995;310:886-7. (8 April.)
- 2 Verhoef F, Sykes MK. Delayed detection of hypoxic events by pulse oximeters: computer simulations. *Anaesthesia* 1990;45: 103-9.
- 3 Davidson JAH, Hosie HE. Limitations of pulse oximetry: respiratory insufficiency—a failure of detection. *BMJ* 1993; 307:372-3.
- 4 Ralston AC, Webb RK, Runciman WB. Potential errors in pulse oximetry. III. Effects of interferences, dyes, dysaemoglobins and other pigments. *Anaesthesia* 1991;46:291-5.
- 5 Stoneham MD, Saville GM, Wilson IH. Knowledge about pulse oximetry among medical and nursing staff. *Lancet* 1994;334: 1339-42.

Gastric lavage poses risks to patient

EDITOR.—J E Charlton comments on the safety of diagnostic and therapeutic endoscopy.¹ Gastric lavage is a procedure akin to upper gastrointestinal endoscopy that is invariably performed on an already sedated patient. It is likewise associated with cardiovascular and respiratory compromise² and should perhaps be undertaken with a similar degree of care and monitoring.

JAMES B ROBINS

Senior registrar in obstetrics and gynaecology
Leeds General Infirmary,
Leeds LS2 9NS

- 1 Charlton JE. Monitoring and supplemental oxygen during endoscopy. *BMJ* 1995;310:886-7. (8 April.)
- 2 Thompson AM, Robins JB, Prescott LF. Changes in cardio-respiratory function during gastric lavage for drug overdose. *Hum Toxicol* 1987;6:215-8.

Paediatric endoscopy should be carried out under general anaesthesia

EDITOR.—J E Charlton highlights the hazards of gastrointestinal endoscopy carried out under sedation.¹ These dangers are not confined to adults.

In a prospective study of 32 children undergoing gastrointestinal endoscopy under sedation, desaturation (oxygen saturation of 90% or less) occurred in 37.5%; this desaturation was unpredictable and failed to correlate with the patient's weight, age, or tolerance of the procedure.² In another prospective analysis, of 57 small children, 10% of the examinations had to be interrupted by removal of the endoscope because the child developed stridor and a low oxygen saturation.³

Monitoring during endoscopy

Attention to sedation techniques may reduce mortality

EDITOR.—J E Charlton draws attention to the estimated overall death rate of 1 in 2000 from upper gastrointestinal endoscopy, which is usually performed under sedation or local anaesthesia, or both.¹ When compared with the overall mortality solely attributable to anaesthesia, which is 1 in 185 000,² this mortality seems unacceptably high.

Recommendations for standards of sedation and for monitoring patients during gastrointestinal endoscopy were published by the British Society of Gastroenterology in 1991,³ and guidelines for sedation by non-anaesthetists were published by a working party of the Royal College of Surgeons in 1993.⁴ Despite this, practice across Britain seems to vary widely.

We were surprised to find that in some regions the preferred benzodiazepine for sedation during endoscopy is diazepam,⁵ which has an active principal metabolite (desmethyldiazepam) with a half life of 48–96 hours. Midazolam has been

Pulse oximetry has limitations

EDITOR.—Most anaesthetists are already aware of the dual problems of oxygenation and monitoring during endoscopy on sedated patients.¹ The solution, however, may not be as simple as introducing supplemental oxygen and using pulse oximetry for all patients. There are crucial differences between a sedated patient undergoing endoscopy and an anaesthetised patient undergoing surgery.

Firstly, for patients undergoing endoscopy there is usually only one operator, who performs the endoscopy as well as giving sedation and cannot therefore give the patient's oxygenation and monitoring full attention during the procedure. The revised recommendations for standards of monitoring during anaesthesia and recovery issued by the Association of Anaesthetists of Great Britain and Ireland state that the anaesthetist should be present throughout the general anaesthesia and that this consideration applies to any technique or sedation carrying a risk of unconsciousness.

Secondly, and most importantly, pulse oximetry is not an infallible monitoring device, though it is

Despite these findings and the well documented data on adults, pulse oximetry and cardiac monitoring have not been universally used in children undergoing endoscopy under sedation.⁴ Clearly, if paediatric endoscopy is performed in this way similar recommendations concerning careful monitoring and supplemental oxygen must apply.

There is, however, a safer and possibly better alternative. After witnessing several major hypoxic events and a respiratory arrest in centres where paediatric gastrointestinal endoscopy is performed under sedation, we have made it our policy to undertake these procedures under short general anaesthesia. This permits both detailed cardio-respiratory monitoring and constant supervision of the patient's airway by the anaesthetist. The procedure is well tolerated (and much preferred by those who have previously undergone endoscopy or jejunal biopsy under sedation) and can usually be carried out on a day case basis. General anaesthesia allows the endoscopist to concentrate on the examination without distraction, and therapeutic interventions such as dilatation of an oesophageal stricture or injection of oesophageal varices can be undertaken during the same session.

In a personal audit of over 300 diagnostic or therapeutic flexible gastrointestinal endoscopies performed in children aged 1 month to 17 years under general anaesthesia since January 1994 there were no major adverse events and all patients recovered fully. The patients included children with severe cerebral palsy, congenital heart disease, and other complex medical problems, who are at greater risk of cardiorespiratory disturbances.

MARK D STRINGER Consultant paediatric surgeon
United Leeds Teaching Hospitals Trust,
Leeds LS2 9NS

PAT J M MCHUGH Consultant anaesthetist

- 1 Charlton JE. Monitoring and supplemental oxygen during endoscopy. *BMJ* 1995;310:886-7. (8 April.)
- 2 Casteel HB, Fiedorek SC, Kiel EA. Arterial blood oxygen desaturation in infants and children during upper gastrointestinal endoscopy. *Gastrointest Endosc* 1990;36:489-93.
- 3 Bendig DW. Pulse oximetry and upper intestinal endoscopy in infants and children. *J Pediatr Gastroenterol Nutr* 1991;12:39-43.
- 4 Ament ME, Berquist WE, Vargas J, Perisic V. Fiberoptic upper intestinal endoscopy in infants and children. *Pediatr Clin North Am* 1988;35:141-55.

Italian data support upper gastrointestinal endoscopy without sedation

EDITOR,—J E Charlton claims that closer monitoring, as well as routine oxygen, during upper gastrointestinal endoscopy would reduce morbidity related to this procedure.¹ Charlton wonders whether it is safe to dismiss cardiographic abnormalities and low oxygen desaturation, since in a recent survey nearly all the complications were respiratory or cardiovascular.² In the same survey 85% of upper gastrointestinal endoscopies were performed under sedation, a figure similar to that detected in a survey in 1990, in which sedation was shown to be the main cause of adverse outcomes.³

We believe that performing upper gastrointestinal endoscopy without sedation would greatly improve its safety and cost effectiveness without any appreciable reduction of efficiency. In our institution, since May 1993, we have performed upper gastrointestinal endoscopies routinely without sedation, with local anaesthesia by lignocaine spray of the oropharynx. We have reviewed the incidental endoscopies performed in the past three months (n=353) to assess the true need for sedation (table); we separated emergency or "operative" from diagnostic upper gastrointestinal endoscopies, because in emergency cases sedation is often performed without prior assessment of the patient's preference. Four per cent of the 285 diagnostic examinations were done under sedation, partly because of patients' explicit request. Among the remaining 273 diagnostic

Number of upper gastrointestinal endoscopies in which sedation was given

	Diagnostic	Operative or emergency	Total
No Sedation given	285	68	353
Sedation given	12	33	45
No sedation given (interrupted)	273 (2)	35 (0)	308 (2)

endoscopies, which were performed without sedation, only two were interrupted or judged technically unsatisfactory, or both, because of the patient's intolerance. These data support our opinion that sedation is not generally needed for upper gastrointestinal endoscopy and that the procedure is well tolerated. We therefore conclude that the easiest, safest, cheapest way to reduce morbidity and mortality associated with upper gastrointestinal endoscopies is to perform them without sedation.

We use sedation at the patient's request, if the patient is overconcerned about the procedure, and in "operative" or emergency settings. This allows a reduction in the number of sedations (and hence in adverse outcomes related to sedation) of over 80%. Pulse oximetry and administration of oxygen could be reserved for patients undergoing sedation—who should be a minority—and their role eventually assessed in a formal randomised controlled trial.

VITTORIO PERI Research fellow in gastroenterology
GIOVANNI GATTO Senior registrar

MARIANO AMUSO Senior registrar
MARIO TRAINA Senior registrar

Endoscopy Unit,
Department of Internal Medicine,
Ospedal "V Cervello,"
Palermo, Italy

- 1 Charlton JE. Monitoring and supplemental oxygen during endoscopy. *BMJ* 1995;310:886-7. (8 April.)
- 2 Quine MA, Bell GD, McCloy RF, Charlton JE, Devlin HB, Hopkins A. Prospective audit of upper gastrointestinal endoscopy in two regions of England: safety, staffing, and sedation methods. *Gut* 1995;36:462-7.
- 3 Daneshmend TK, Bell GD, Logan RFA. Sedation for upper gastrointestinal endoscopy: results of a nation-wide survey. *Gut* 1991;32:12-5.

Hypoxia during endoscopy also occurs in unsedated patients

EDITOR,—J E Charlton emphasises the risks of upper alimentary endoscopy, particularly in high risk patients undergoing the procedure with sedation,¹ but fails to emphasise that the risk can also occur in unsedated patients. We have previously reported an oxygen saturation of $\leq 92\%$ in 41% (24/59) of unsedated patients breathing room air.² The proportion of unsedated patients with an oxygen saturation of $\leq 92\%$ fell significantly to 11% (6/56) when supplemental oxygen was given. Similar findings were found in a group of patients undergoing endoscopy with sedation. In line with Charlton's comments, however, there were no significant differences in maximum pulse rate in any of these groups, indicating that alleviating the hypoxia may not reduce the cardiac risk. Despite this we agree that it would not be safe to dismiss the benefits of supplemental oxygen during endoscopy and believe that it should be part of routine practice.

M W R REED Senior lecturer in surgery
C S REILLY Professor of anaesthetics

Department of Surgical and Anaesthetic Sciences,
Royal Hallamshire Hospital,
Sheffield S10 2JF

- 1 Charlton JE. Monitoring and supplemental oxygen during endoscopy. *BMJ* 1995;310:886-7. (8 April.)
- 2 Reed MWR, O'Leary D, Duncan JL, Majeed AW, Wright B, Reilly CS. Effects of sedation and supplemental oxygen during upper alimentary tract endoscopy. *Scand J Gastroenterol* 1993; 28:319-22.

Open access echocardiography

EDITOR,—We have six years of experience of open access echocardiography for general practitioners, so we read the paper by C M Francis and colleagues¹ and the accompanying editorial by J R Hampton and A R Barlow² with interest.

We have performed a retrospective study of echocardiograms ordered by general practitioners at a health teaching centre of family medicine between January 1985 and April 1991. The centre has six general practitioners and covers a population of 22 000. There are also six trainees in family medicine and a paediatrician. The echocardiography was done by an experienced cardiologist working at a large hospital, which covers a population of 350 000.

There were no predetermined criteria for ordering echocardiograms. During the study period 127 echocardiograms were requested, but data were missing in 14, leaving 113 for analysis.

Eighty echocardiograms had been ordered because of persistent symptoms, 25 because of dyspnoea, 21 because of palpitations, 14 because of chest pain, and five because of syncope. Ninety one had been ordered for diagnosis and 22 to assist management in patients with existing diseases. Sixty eight echocardiograms showed abnormal results, including 39 valve abnormalities and nine conditions that did not affect the valves. Pathological heart murmurs were less common in patients aged 20 and under. Symptoms and pathological findings on echocardiography predicted referral to a cardiologist.

Our results strengthen our impression that echocardiography is useful to exclude or confirm disease in primary care.

We think it necessary to outline some referral criteria in coordination with the cardiology department, which also prepares the trainees in family medicine in the use and interpretation of echocardiograms.

Our experience was wholly satisfactory. Unfortunately, the echocardiography department is now overworked and the cardiologist must decide whether to perform echocardiography.

EDUARDO CALVO CORBELLA Family practitioner

JUAN JOSE RODRIGUEZ ALONSO Vocational trainee in family practice
CELIA QUINTANA LUZON Family practitioner

University Health Centre
Pozuelo de Alarcón,
Teaching Unit of Family Practice,
Universidad Autónoma,
Madrid,
Spain

MAGDALENA GONZÁLEZ Cardiologist

Cardiology Service,
Puerto de Hierro Hospital,
Madrid

- 1 Francis CM, Caruana L, Karney P, Love M, Sutherland GR, Starkey IR, et al. Open access echocardiography in management of heart failure in the community. *BMJ* 1995;310:634-6. (11 March.)
- 2 Hampton JR, Barlow AR. Open access. *BMJ* 1995;310:611-2.

Patients' perceptions of need for primary health care services

Patients could consider general good rather than their own

EDITOR,—Jane L Hopton and Maria Dlugolecka examine the feasibility of using patients' perceptions of need for primary health care services to develop priorities.¹ Their study illustrates three important deficiencies in engaging the general public to develop priorities for health care services. Firstly, we wonder how representative the respondents were of the general population. As 64% of the