did it?" and "How do we stop it?" may be of little relevance after an earthquake, they are uppermost in the mind of anyone subject to continual bombardment and ethnic cleansing. Failure to engage with these issues can increase a sense of victimhood and disempowerment; and attempts to remain politically neutral in the face of genocide may well be construed as tacit collaboration with the aggressor and make effective therapeutic work impossible.

Human identity rests on the network of social relations that we build around ourselves: the tie of family, work, and community, and the emotional bond we make with our physical environment. One might argue that the main psychic injury of war is the disruption of those ties, the destruction of identity through the destruction of our social world. Thus any thorough evaluation of psychological treatments for traumatic stress should compare their effectiveness with the impact on mental health of non-psychological interventions such as community redevelopment projects. Only then can scarce resources be allocated in the most appropriate way.

LYNNE JONES
Senior registrar

Child, Adolescent, and Family Psychiatry,
Institute of Family Psychiatry,
Ipswich,
Suffolk IP1 3TF

- 1 Raphael B, Meldrum L, McFarlane A. Does debriefing after psychological trauma work? *BMJ* 1995;310:1479-80. (10 June.)
- 2 Bracken P, Giller J, Summerfield D. Psychological responses to war and atrocity: the limitations of current concepts. *Soc Sci Med* 1995;40:1073-82.
- 3 Jones L. Letter from Sarajevo. *BMJ* 1995;310:1052-4. (22 April.)

Response to treatment varies

EDITOR,—Beverley Raphael and colleagues' editorial raises many questions about psychological debriefing and post-traumatic stress reactions.¹ A military debriefing is an analysis of the events occurring on a mission and the lessons learnt. The primary aim of psychological debriefing is to provide the person with as much information as possible from all the sources available to enable the cognitive appraisal and emotional processing of a traumatic experience. This may be therapeutic or may not be. Traumatic incidents irrevocably change people as they challenge fundamental beliefs and values—spiritual, philosophical, moral, and existential. In any experience many post-traumatic stress reactions are chimeric, and even the *Diagnostic and Statistical Manual of Mental Disorders*, fourth edition, reflects the importance of the meaning of an event for the person.²

Any post-traumatic stress reaction is the product of three almost infinitely variable factors: the person's personality, coping mechanisms, and life experiences or abuses; aspects and meaning of the traumatic event itself; and the psychosocial environment into which the person returns. My, and others',³ experience is that people with no, or few, scars from the vicissitudes of life and with good psychosocial support can survive almost anything. This model also explains how psychological debriefing may be followed by idiosyncratic responses or an apparent worsening in the person's condition; explains why no one intervention will "work" either before or after the development of a post-traumatic stress reaction; and suggests why it will be difficult to find controls for any randomised trial.

After traumas there is a natural need and desire to help, especially as social, cultural, spiritual, and humanistic dimensions exist alongside the medical and psychological ones. But there are darker aspects to providing help to the victims of traumas. These lie in voyeurism—vicariously listening to or experiencing salacious, sadistic, violent, even

pornographic details of another person's life. It can prove an abusive experience for all parties. For this reason, those who provide psychological debriefing or treatment for post-traumatic stress reactions must have good psychotherapeutic skills and supervision.

Whatever the outcome of trials, I suspect that psychological debriefing is here to stay. Perhaps it has always been with us: "But we two snugly indoors here may drink and eat and revel in an interchange of sorrows—sorrows that are memories, I mean; for when a man has endured deeply and strayed far from home he can cull solace from the rehearsal of old griefs. And so I will meet your questioning."⁴

IAN P PALMER
Lecturer in military psychiatry

Royal Army Medical College,
London SW1P 4RJ

- 1 Raphael B, Meldrum L, McFarlane A. Does debriefing after psychological trauma work? *BMJ* 1995;310:1479-80. (10 June.)
- 2 Davidson J, Foa E, eds. *Post-traumatic stress disorder, DSM IV and beyond*. Washington: American Psychiatric Press, 1993.
- 3 Bisson J, Deahl M. Psychological debriefing and prevention of post-traumatic stress. *Br J Psychiatry* 1994;165:717-20.
- 4 Solomon S, Gerrity E, Muff A. Efficacy of treatments for post-traumatic stress disorder. *JAMA* 1992;268:633-7.
- 5 Homer. *The odyssey*. Book XV. Oxford: Oxford University Press, 1991. (Translator Lawrence TE.)

Voluntary euthanasia commands majority support

EDITOR,—I fail to see how Anne Rodway can assert that voluntary euthanasia in Britain enjoys only minority support among the general public¹ when results of scientifically conducted polls emphatically tell a different story. A poll commissioned by the Voluntary Euthanasia Society in April 1993—the latest in a series each showing an increase on the last—showed 79% in support of euthanasia,² and countless other surveys, albeit less scientific, after newspaper and magazine articles and television and radio programmes show similar results. If Rodway still doubts the veracity of these surveys, I would draw her attention to a poll in 1987 commissioned by an antieuthanasia group, the World Federation of Doctors Who Respect Human Life, which found 72% in favour of euthanasia.³

Finally, proponents of voluntary euthanasia do not see it as "a substitute for caring and supporting"; they see it as a widening of choice for dying patients and a final act of care.

JOHN OLIVER
General secretary

Voluntary Euthanasia Society,
London W8 5PG

- 1 Rodway A. Euthanasia debate. *BMJ* 1995;310:1466. (3 June.)
- 2 National Opinion Poll. *Euthanasia*. London: NOP, 1993.
- 3 Market and Opinion Research International. *Euthanasia*. London: MORI, 1987.

Their lordships on medical research

EDITOR,—I am surprised that Richard Smith finds the report of the select committee on science and technology, based on the work of the sub-committee that I chaired, "too backward looking."¹ In the report the committee commends the government's initiative on research and development, the work of Professor Michael Peckham and his regional directors, the increasing development of health services research, and new trends in knowledge based and evidence based medical practice. The committee also recommends explicitly that when the recommendations of the Culyer taskforce are implemented a separate stream of funds should be identified at regional

level to support research in general practice and community care as well as research by nurses and members of the other health care professions.

It is true that the committee is concerned about the infrastructure funding in the NHS for biomedical research and research driven by curiosity. The committee does not, however, reject the principle of a research assessment exercise within the NHS as in the universities; it suggests that, to avoid massive duplication of effort, the two processes should be combined and that a core of facilities funding should be preserved to maintain the ability of hospitals and other university based research departments (including academic departments of general practice) to continue with that biomedical and predominantly clinical research on which the future wellbeing of the nation depends. This is in no sense promoting an "ivory tower" approach, but it is a truism that today's discovery in basic medical science brings tomorrow's practical development in patient care. The report also quotes conclusive evidence to indicate that the major specialist research centres, some of which are indeed being starved of tertiary referrals, not only find that their research brings new knowledge about the management of disease but also improves clinical outcomes.

I am glad that Smith commends the committee's wish to see an urgent inquiry into the present and future of clinical academic medicine. He is right in saying that career problems in this field are not new, but the many new factors referred to in the report have greatly increased the sense of crisis that now abounds in the clinical academic community, and I trust that in this regard, as well as in many others, the committee's warnings will be heeded.

JOHN WALTON
Chairman, subcommittee I
(medical research and the NHS reforms)

Science and Technology Committee,
House of Lords,
London SW1A 0PW

- 1 Smith R. Their lordships on medical research. *BMJ* 1995;310:1552. (17 June.)

Effects of homoeopathy

Trial did not evaluate "true" homoeopathy

EDITOR,—Per Lökken and colleagues have made a laudable attempt to evaluate homoeopathy scientifically.¹ For orthodox methodologists the design of their study will seem rigorous, and therefore most conventional physicians and dentists will conclude that homoeopathy is not effective for the given indication. Homoeopaths, on the other hand, might have doubts about this conclusion, arguing that the protocol did not allow for the necessary freedom of homoeopathic prescription: the doses were fixed (at D30), and so was the treatment schedule (four doses during the first three hours, etc), yet homoeopathy requires these to be flexible and fully individualised. Thus homoeopaths might think that the results would have been different if true homoeopathy had been practised.

It might also be argued that arnica, which was the most commonly used remedy in this study, is ineffective but homoeopathy may still work. Others might think that it was a bad idea for some of the subjects to be students of natural medicine, who might respond typically. Finally, one might speculate that the prescribing homoeopaths (who and how many were they? Did they reach individual or consensus decisions about which drug to use?) were ineffectively using a potentially effective form of treatment.

These arguments highlight the immense difficulties that are encountered when evaluating homoeopathy scientifically. Given that there are hundreds of remedies to be applied in hundreds of

ways, one would, strictly speaking, need hundreds of thousands of randomised trials to arrive at conclusive answers about the efficacy of this form of treatment. Clearly, this is not practical. The Centre for Complementary Health Studies is therefore planning a trial designed to answer the much broader question, "does homoeopathy work (regardless of remedy, dose, etc)?" I would be delighted to hear from potential participants in this multicentre study.

E ERNST
Professor of complementary medicine

Centre for Complementary Health Studies,
University of Exeter,
Exeter EX2 4NT

1 Lökken P, Straumshøj PA, Tveiten D, Skjelbred P, Borchgrevink CF. Effect of homoeopathy on pain and other events after acute trauma: placebo controlled trial with bilateral oral surgery. *BMJ* 1994;310:1439-42. (3 June.)

Trial puts negative gloss on essentially positive results

EDITOR,—It is well known that reports of trials tend to exaggerate differences between treatments.¹ By using multiple end points, repeated measures, subgroup analyses, and statistical "data dredging" authors are often able to put a positive gloss on essentially negative results. Per Lökken and colleagues' trial of homoeopathy after dental surgery is a reversal of this trend.² Four predefined clinical outcome measures were used: pain, swelling, bleeding, and trismus. There was no difference in the first three of these measures between the groups. Homoeopathy was statistically superior to placebo for the fourth measure. Yet the authors conclude that there was "no significant effect of homoeopathy on pain and other events." One of the key messages states, "no positive evidence could be found for homoeopathic effects."

The results of the trial clearly do not validate the claim that homoeopathic remedies are of appreciable clinical value after dental surgery. But ignoring significant results is surely as much of an error as creating them by massaging data.

ANDREW VICKERS
Director, information services

Research Council for Complementary Medicine,
London WC1N 3JF

1 Pocock SJ, Hughes MD, Lee RJ. Statistical problems in the reporting of clinical trials. *N Engl J Med* 1987;317:426-32.

2 Lökken P, Straumshøj PA, Tveiten D, Skjelbred P, Borchgrevink CF. Effect of homoeopathy on pain and other events after acute trauma: placebo controlled trial with bilateral oral surgery. *BMJ* 1994;310:1439-42. (3 June.)

Thyroid cancer in the United States since accident at Chernobyl

EDITOR,—V A Stsiazhko and colleagues' letter on childhood thyroid cancer near Chernobyl raises serious issues on a broader scale.¹ If fallout from Chernobyl travelled hundreds of kilometres it seems reasonable that similar findings may occur in more distant populations.

Fallout from Chernobyl began arriving in Connecticut in the United States, 7250 km west of the reactor, about 15 days after the explosion which occurred on 26 April 1986. The concentration of iodine-131 (one of the few radionuclides tracked) in Connecticut's milk increased in the weeks after the accident (the state average in 1983-90 was about 0.07 Bq/l).²

The rate of thyroid cancer among children aged under 15 in Connecticut rose sharply (from 1.6 to 3.1 per million) after 1989 (figure), when increases in children in the area around Chernobyl began.^{3,4} This change is similar to that seen in the (upwind)

Number of cases of thyroid cancer per million children aged under 15 and age adjusted rate per million population in Connecticut 1935-92 (source: Connecticut tumour registry) and concentration of iodine-131 in milk in Connecticut²

Ukraine.¹ It is also intriguing to juxtapose rates of thyroid cancer for all age groups with those in children. The age adjusted rate in Connecticut between 1985-9 and 1990-2 rose from 34.6 to 42.9 per million, after 10 years of no change.

There is an apparent five year lag between exposure to radioactive fallout from Chernobyl and rising rates of thyroid cancer. In contrast, lag times have been reported to be up to 40 years for iatrogenic thyroid cancer due to irradiation.³ Continued tracking of rates may show a progressive rise in thyroid cancer attributable to fallout of a variety of (iodine and other) high energy nucleotides from Chernobyl. It seems prudent to examine further the long term effects, in particular thyroid cancer, on populations distant from Chernobyl.

WILLIAM REID
Oncologist-haematologist

Biological Therapy Institute,
357 Riverside Drive,
Franklin, TN, USA

Radiation and Public Health Project,
786 Carroll Street,
Brooklyn, NY, USA

JOSEPH MANGANO
Consultant

- 1 Stsiazhko VA, Tsyb AF, Tronko ND, Souchevitch G, Baverstock KF. Childhood thyroid cancer since accident at Chernobyl. *BMJ* 1995;310:801. (25 March.)
- 2 US Environmental Protection Agency. *Environmental radiation data*. Montgomery, AL: EPA, 1986: 94-107. (Report 46.)
- 3 Kazakov VS, Demidchik EP, Astakhova LN. Thyroid cancer after Chernobyl. *Nature* 1992;359:21.
- 4 Likhartev IA, Sobolev BG, Kairo IA, Tronko ND, Bogdanava TI, Oleinic VA, et al. Thyroid cancer in the Ukraine. *Nature* 1995;375:365.
- 5 Raventos A, Winship T. The latent intervals for thyroid cancer following irradiation. *Radiology* 1964;83:501-8.

Spinal cord injury in Rugby Union players

EDITOR,—Timothy Noakes and Ismail Jakoet are right in saying that there are no reliable figures for the number of spinal cord injuries sustained by rugby players.¹ The Duke of Cornwall Spinal Injuries Centre states that 16% of all spinal injuries result from sporting activities, with 2.5% being due to rugby.²

For the past 15 months I have been working as a volunteer for the Trevor Jones Tetraplegic Trust, set up to help people completely paralysed as a

result of sports injuries. The trust has the names of 140 people disabled in this way, and I have analysed details on 98 of these from the registration forms filled in by the patients or, occasionally, a relative. Fifty two of the injuries occurred while the person was playing Rugby Union football: four in the 1970s, 28 in the '80s, and 12 in the '90s. The figures are nowhere near complete: the '90s are only half run, and many of those injured are still in injuries centres.

But I find the figures worrying. There is little evidence of a reduction in the number of such lesions in Britain since the last change in Rugby Union's rules in 1985, as 22 of them were sustained after that date. These are devastating accidents for those who would otherwise have had a promising future. Of the 52 patients, 31 were 25 years old or younger at the time of injury. Only 28 list any present occupation, and 10 of these are students, who may or may not get a job. Insurance compensation is derisory compared with that received by victims of road traffic accidents, and if anyone, with enormous effort, makes a success of a job then any additional income may be clawed back by the social services. "I can't see how I shall ever get off income support," said one man, who had progressed from computer operator to information technology manager for his firm.

It is not only the epidemiology that needs study. Little information is available on the late outcome of spinal injuries: whether those who get adequate compensation fare better, what happens to the unknown number who lose touch with their unit, and how many die at an early age and of what causes. Tetraplegic people live under constant threat of pressure sores, urinary infections, and rejection by society. Rugby, I am told, is a great game; it is certainly a profitable one. But it exacts a dreadful toll from some. My 14 year old, lightly built grandson is now at a school where rugby is played, and I have asked his parents to encourage him to choose another sport. Am I overreacting?

ANNE SAVAGE
Medical adviser

Trevor Jones Tetraplegic Trust,
London W1X 1RA

1 Noakes T, Jakoet I. Spinal cord injury in rugby union players. *BMJ* 1995;310:1345-6. (27 May.)

2 Grundy D, Swain A. *ABC of spinal cord injury*. 2nd ed. London: BMJ Publishing, 1993:1.

Training in family planning encompasses several disciplines

EDITOR,—P D Woolley seems not only to have misunderstood the editorial by Yvonne Stedman and Max Elstein¹ but to be unaware of the developments in family planning and sexual health issues.²

Basic training in family planning has been structured for many years, long before training in most other specialties, and includes knowledge of infections. Two years ago the Faculty of Family Planning and Reproductive Health Care of the Royal College of Obstetricians and Gynaecologists was founded; it has developed a membership examination (MFFP), which will start this autumn and includes genitourinary medicine in its syllabus. In contrast, doctors can and often do become consultants in genitourinary medicine without experience in family planning or gynaecology.

Family planning is rapidly becoming a consultant led specialty, often by people with broad experience and multiple qualifications, including in genitourinary medicine. As in most services, other clinicians in various grades will always be needed to help with family planning clinics, and no one thinks that family planning should encompass all genitourinary medicine.

Surely, however, family planning services should be accessible and acceptable to all who need them