Extracontractual referrals have recently exposed a growing interest among general practitioners, psychologists, and community psychiatric nurses for specialist counselling services.

In our experience accurate data on extracontractual referrals can be collected rapidly, are relatively simple to analyse, and give procedure specific details and the possibility of case by case validation. A crude statistic on extracontractual referrals can be interpreted in a variety of ways. It is most useful when regarded as one piece of a jigsaw of evidence concerning services and interpreted in a context that includes clinical advice. Extracontractual referrals offer purchasers an easy way to tap into the local general practitioners' and hospital consultants' "bush telegraph" concerning the range and quality of clinical services and the views of patients as interpreted by their doctors. Extracontractual referrals should therefore be recognised as valuable nuggets of audit information. Rather than being a foe to the accountant, each extracontractual referral can be a friend to the purchaser of better quality health services.

EDWIN PUGH
Director of public health
DEBORAH WILLIAMS
Purchasing support manager

County Durham Health Commission,
Durham DH1 5XZ

1 Ghodse B. Extracontractual referrals: safety valve or administrative paperchase? *BMJ* 1995;310:1573-6. (17 June.)

Measles data are reliable

EDITOR,—Nigel Higson claims that the only source of data on measles cases which has remained consistent before and after the measles-rubella campaign is statutory notification and that numbers of notified cases have shown little decline.¹ As he admits, numbers of notifications are less reliable than cases confirmed by saliva testing, but this service was not widely available before the campaign. The saliva assay, however, was extensively piloted in 18 districts of the United Kingdom between October 1991 and October 1994.^{2,3} During this pilot 36-38% of patients with measles tested were IgM positive,^{2,3} suggesting recent infection, compared to only 4% of patients tested since the campaign. Since November 1994, saliva samples have been received in over 50% of notified cases and the proportion that is IgM positive has fallen with time (table).

In addition, measles cases confirmed by serology, isolation, or antigen detection have been reported to the Communicable Disease Surveillance Centre by laboratories in England and Wales for over 20 years. Unlike notifications, the number of laboratory reports has fallen dramatically, from 224 cases reported with sample dates between 1 January and 31 May in 1994 compared with only 52 in the same period of 1995. Many of the 1995 cases occurred during late 1994 and early 1995, when the full effect of the campaign had not been felt—only four cases have been reported with sample dates during May 1995.

Both data sources clearly indicate that measles transmission has been substantially reduced by the vaccination campaign, in particular in 5-16 year

Saliva tests performed in notified measles cases from England and Wales

Year/week of notification	Notifications (provisional data)	No (%) of samples received	No (%) of samples positive for IgM
94/44 to 94/51	2585	1356 (52)	117 (8.6)
94/51 to 95/07	1672	1002 (60)	27 (2.7)
95/08 to 95/11	1568	910 (58)	10 (1.0)
95/16 to 95/23	1188	658 (55)	5 (0.8)
Total	7013	3926 (56)	159 (4.0)

olds—the age group targeted.⁴ Small numbers of measles cases are still being confirmed in preschool children and unvaccinated adults, and at least four recent infections seem to have been contracted abroad. A comprehensive surveillance programme for measles is now established and will allow us to monitor the longer term effects of the measles-rubella campaign and to define future vaccination policy in England and Wales.

MARY RAMSAY
Consultant epidemiologist
Immunisation Division,
PHLS Communicable Disease Surveillance Centre,
London NW9 5EQ

DAVID BROWN
Consultant virologist
Enteric and Respiratory Virus Laboratory,
PHLS Virus Reference Division,
London NW9 5EQ

1 Higson N. Evaluating the measles rubella campaign. *BMJ* 1995;311:62.
2 Brown DW, Ramsay ME, Richards AF, Miller E. Salivary diagnosis of measles: a study of notified cases in the United Kingdom, 1991-1993. *BMJ* 1994;308:1015-7.
3 Ramsay M, Gay N, Miller E, Rush M, White J, Morgan-Capner P, Brown D. The epidemiology of measles in England and Wales: rationale for the 1994 national vaccination campaign. *Communicable Disease Report* 1994;4:R141-6.
4 Measles transmission interrupted in school children. *Communicable Disease Report* 1995;5:35.

Treatment of children with asthma

Entrust parents with "crisis pack" of steroids

EDITOR,—In their excellent and comprehensive account of treating asthmatic children John Rees and John Price rightly emphasise the importance of "partnership in management" and "self management plans," including "an agreed action plan of what to do...if the child has an acute asthma attack."¹ However, in the subsequent part of the article dealing with acute severe asthma, the principle of partnership with parents seems to extend only as far as giving four hourly relief medication (dose not specified) and calling a doctor if there is no improvement. The adjacent paragraph refers to the need for a large dose of a β stimulant bronchodilator and considering starting oral prednisolone, treatment that most readers will interpret as restricted to a doctor rather than a parent.

Surely any parent who is fit to be entrusted with the overall care of a child is also fit to be entrusted with these forms of treatment, which can be started early in an attack and can often obviate the need for any medical involvement. There are often difficulties and delays in seeing a doctor, even in Britain, and the increasing number of families taking holidays abroad and in remote places is a strong argument for us trying to increase families' self reliance and reduce dependence on doctors.

For the past 10 years I have given a "crisis pack" of steroids to all the parents of asthmatic children on my list. If sufficient time is taken to supplement good written information, most families develop the confidence to use short courses of oral prednisolone very judiciously. Obviously, the concept of self reliance could be taken too far, and parents need to be told to seek medical help if the crisis pack is not working. In these circumstances, I prefer to give parents free access to our paediatric ward, as calling the general practitioner is unnecessary and only leads to delay in admission.

NIGEL SPEIGHT
Consultant paediatrician
Dryburn Hospital,
Durham DH1 5TW

1 Rees J, Price J. Asthma in children: treatment. *BMJ* 1995;310:1522-7. (10 June.)

Author's reply

EDITOR,—We thank Nigel Speight for highlighting an important issue relating to the management at home of acute asthma in children. In this section of our article we say that consideration should be given to starting oral prednisolone. It was certainly not our intention to imply that parents were excluded from making this decision. We completely agree that many families, given adequate guidance and explanation, will know when to give their child a course of oral steroid. On the other hand some families would be apprehensive about taking this responsibility, and we must continue to provide them with the support they need.

Short courses of oral prednisolone given at home increase rates of recovery and may prevent hospital admissions.¹ We agree that initiation by parents is appropriate and quicker than waiting for a doctor.

Occasional short courses of prednisolone are safe. There is transient suppression of hypothalamic-pituitary-adrenal axis function but this returns to normal within 10 days.² However, function of the hypothalamic-pituitary-adrenal axis remains subnormal in children receiving more than four short courses of prednisolone per year.³ Frequent courses of prednisolone indicate poor asthma control, so the total number given to the child, whether started by the family or the doctor, should be recorded. Parents should be advised to tell their doctor if their child is having asthma attacks sufficiently severe to require repeated courses of oral corticosteroids. This is an indication for a general review of the asthma, including reassessment of the home environment and the need for and adherence with regular inhaled anti-inflammatory medication.

JOHN PRICE
Professor
Department of Thoracic Medicine,
King's College School of Medicine and Dentistry,
London SE5 9PJ

JOHN REES
Consultant physician
Guy's Hospital,
London SE1 9RT

1 Despinande A, McKenzie SA. Short course of steroids in home treatment of children with acute asthma. *BMJ* 1986;293:169-71.
2 Zora JA, Zimmerman D, Carey TL, O'Connell EJ, Yunginger JW. Hypothalamic-pituitary-adrenal axis suppression after short term, high dose, glucocorticoid therapy in children with asthma. *J Allergy Clin Immunol* 1986;77:9-13.
3 Dolan LM, Kesarwala HH, Holroyde JC, Fischer TJ. Short term, high dose, systemic steroids in children with asthma: the effect on the hypothalamic-pituitary-adrenal axis. *J Allergy Clin Immunol* 1987;80:81-7.

Managing the complications of childhood arthritis

EDITOR,—I agree with Aideen Landers that the recommendations for ocular slit lamp screening are confusing.¹ The most important points to emphasise are that children with juvenile chronic arthritis are at greater risk of developing asymptomatic chronic anterior uveitis than the normal population and that irreversible complications may be prevented if the uveitis is detected early by slit lamp examination and treated effectively. Landers cited the guidelines for slit lamp screening issued by a joint working party of the Royal College of Ophthalmologists and the British Paediatric Association,² but unfortunately these guidelines do not seem to take into account two key questions.

Firstly, how quickly can irreversible visual impairment follow the onset of chronic anterior uveitis? This should determine the frequency of screening for at risk children. Irreversible complications are likely to occur if the chronic anterior uveitis is untreated for longer than six months (this is based on anecdotal evidence only). Annual screening would not seem sensible for any child with arthritis.