- 4 Simmons A, Barrington S, O'Doherty MJ, Coakley AJ. Dual energy x ray absorptiometry normal reference range use within the UK and the effect of different normal ranges on the assessment of bone density. *Br J Radiol* 1995;68:903-9.
- 5 Simmons A, Barrington S, O'Doherty MJ, Coakley AJ. A survey of dual-energy x-ray absorptiometry normal reference ranges used within the United Kingdom and their effect on patient classification. *Nucl Med Commun* (in press).

Different guidelines will yield different estimates of prevalence

EDITOR,—J E Compston and colleagues suggest that in clinical practice patients should be selected for bone densitometry on the basis of clinical or historical risk factors.¹ Having recently assessed the guidelines on the detection of osteoporosis that operated during the first year of operation of the bone densitometry service in the Wirral, we would point out that care is needed in selecting criteria for bone densitometry as different criteria lead to different rates of detection of disease.

We have the only densitometer in Wirral, which has a catchment population of 365 000. Our guidelines stated that the following categories of patients would be eligible for scanning: women who had taken hormone replacement therapy for 18 months, to determine whether the treatment should be continued; women who were unable to take hormone replacement therapy; women who had sustained a trivial fracture (including vertebral fractures); women taking corticosteroids; and private referrals.

Altogether 435 scans were obtained in the first year, of which 307 fell within the guidelines. The table shows the number of scans obtained for various indications and the incidence of osteoporosis detected (on the basis of the World Health Organisation's guidelines.² The highest proportions of women with osteoporosis were found in the groups with osteopenia evident in x ray films (58%) and with trivial fractures (50%) ($P=0.002$, χ^2 test). When patients who were already taking antiosteoporotic treatment (those who had had 18 months of hormone replacement therapy or were taking etidronate) were excluded we found no difference in the proportion with osteoporosis between the group referred according to our guidelines and the other referrals ($P=0.74$ for the lumbar spine, $P=0.44$ for the femoral neck, χ^2 test). When, however, we reanalysed our data using the guidelines suggested by Compston and colleagues (trivial fractures, use of corticosteroids, premature menopause, and osteopenia evident in x ray films) we found a significant difference in the proportion of patients with osteoporosis between the group referred according to their guidelines (75/178 (42%) had osteoporosis of the lumbar spine and 99/178 (56%) of the femoral neck) and the others (77/257 (30%) had osteoporosis of the lumbar spine and 109/257 (43%) of the femoral neck; $P=0.02$ for the lumbar spine, $P=0.02$ for the femoral neck, χ^2 test).

In conclusion, 71% of our referrals adhered to our guidelines, which suggests that the guidelines are reasonably well followed. A balance has to be achieved, however, between setting guidelines to detect disease and setting guidelines that take

into account patients' preferences (for example, scanning of women who were already taking hormone replacement therapy made up 15% of our workload) but have a lower rate of detection.

S S YEAP
Senior registrar

Department of Rheumatology,
Leicester Royal Infirmary NHS Trust,
Leicester LE1 5WW

N A KEATING
Research assistant

Department of Public Health and Epidemiology,
Queen's Medical Centre,
Nottingham NG7 2UH

T D KENNEDY
Consultant physician and rheumatologist

Department of Rheumatology,
Arrowe Park Hospital,
Wirral L49 5PE

1 Compston JE, Cooper C, Kanis JA. Bone densitometry in clinical practice. *BMJ* 1995;310:1507-10. (10 June.)

2 World Health Organisation. Assessment of fracture risk and its application to screening for postmenopausal osteoporosis. *WHO Tech Rep Series* 1994;No 843.

Authors' reply

EDITOR,—We agree with Andrew Simmons and colleagues about the need to standardise reference values for bone densitometry; differences between manufacturers' reference data have been reported previously,¹ as have differences in reference data from different centres in Britain.² The number and selection of subjects on whom these data are based and geographic differences in bone density will affect the absolute values measured, the pattern of changes related to age, and the variance. These factors in turn contribute to variations in the classification of patients, whether this is based on T scores, Z scores, or absolute values. The use of Z scores to diagnose osteoporosis is illogical since the prevalence of osteoporosis, thus defined, would not increase with age, while the use of an absolute value as a diagnostic criterion is problematic because differences between measurement systems in calibration, detection of the edge of the bone, and methodological approach produce different absolute values for a given bone density.³ Until there is better standardisation of reference data, inconsistencies in classification will persist, regardless of which diagnostic criteria are used; however, T scores provide the most appropriate means of defining osteoporosis, and problems associated with reference data, while requiring attention, should not be allowed to detract from the clinical value of bone densitometry as currently practised.

In response to S S Yeap and colleagues' letter, it is self evident that different guidelines for bone densitometry will result in varying rates of detection of osteoporosis. Assessment of guidelines should not, however, be based solely on the percentage of patients found to have osteoporosis since high rates of detection may reflect inappropriate use of bone densitometry in patients in whom a diagnosis has already been made; conversely, the exclusion of osteoporosis in patients with risk factors is a valuable component

of bone densitometry services since it provides reassurance and avoids unnecessary treatment. Finally, guidelines that take into account patients' preferences are unlikely to be viewed favourably by health authorities or commissions that are approached to provide funding for bone densitometry.

JULIET COMPSTON
Honorary consultant physician

Department of Medicine,
University of Cambridge Clinical School,
Addenbrooke's Hospital,
Cambridge CB2 2QQ

CYRUS COOPER
Reader in rheumatology

MRC Epidemiology Unit,
Southampton General Hospital,
Southampton SO16 6YD

JOHN KANIS
Professor of human metabolism
and clinical biochemistry

University of Sheffield Medical School,
Sheffield S10 2RX

1 Laskey MA, Crisp AJ, Cole TJ, Compston JE. Comparison of the effect of different reference data on Lunar DPX and Hologic QDR-1000 dual-energy X-ray absorptiometers. *Br J Radiol* 1992;65:1124-9.

2 Murrills AJ, Taylor PA, Cotton AM, Petley GW, Cooper C, Wilkin TJ. Bone mineral density in Southampton women. In: Ring EFJ, Elvins DM, Bhalla AK, eds. *Current research in osteoporosis and bone mineral measurement III*. London: British Institute of Radiology, 1994:61.

3 Laskey MA, Flaxman ME, Barber RW, Trafford S, Hayball MP, Crisp AJ, et al. Comparative performance in vitro and in vivo of Lunar DPX and Hologic QDR-1000 dual energy X-ray absorptiometers. *Br J Radiol* 1991;64:1023-9.

Comparing hospital and chiropractic treatment for back pain

Trial did not compare like with like

EDITOR,—In the move towards evidence based medicine, randomised trials should be welcomed. The study by T W Meade and colleagues, however, is flawed in both its methodology and its analysis.^{1,2}

Essentially, the study does not compare like with like. There is no information on outcomes of treatment for different grades of physiotherapists, for the physiotherapists' differing levels of training in manipulation, or for Cyriax versus Maitland manipulation when compared with the outcomes of chiropractic. Each treatment group had a different number of treatments and probably received treatment for different lengths of time. When the number of treatments was similar (at six weeks) there was no significant difference in the mean changes in the Oswestry scores between the two therapies. While significance was reached at six months, when the number of extra chiropractic treatments "was not yet extensive," there was no significant difference at one year.

We suggest that the key messages from this study should advocate more extensive investment in postgraduate training in physiotherapy and in physiotherapists to achieve treatment that is more equitable with that in the private sector. Furthermore, randomised controlled trials in physiotherapy, chiropractic, and other complementary therapies are needed so that NHS purchasers can choose the most beneficial and cost effective treatments for their patients.

ALISON WAKEFIELD
Senior physiotherapist

Princess Margaret Rose
Orthopaedic Hospital,
Edinburgh EH10 7ED

MARTIN BULL
Registrar in public health medicine

Lothian Health,
Edinburgh EH8 9RS

1 Meade TW, Dyer S, Browne W, Frank AO. Randomised comparison of chiropractic and hospital outpatient management for low back pain: results from extended follow up. *BMJ* 1995;311:349-51. (8 August.)

Incidence of osteoporosis detected for each indication for bone densitometry

	No of women	No (%) with osteoporosis of:	
		Lumbar spine	Femoral neck
Taken hormone replacement therapy for 18 months	67	18 (27)	21 (31)
Unable to take hormone replacement therapy	61	18 (30)	27 (44)
Trivial fracture (including vertebral fracture)	91	45 (50)	55 (60)
Use of steroids	50	17 (34)	27 (54)
Private referral	38	8 (21)	13 (34)
Premature menopause	18	2 (11)	5 (28)
Taking etidronate	16	7 (44)	13 (81)
Osteopenia evident in x ray film	19	11 (58)	12 (63)
Family history of osteoporosis	27	8 (30)	13 (48)
Other	48	18 (38)	22 (46)
Total	435	152 (35)	208 (48)

Results were clinically trivial

EDITOR,—T W Meade and colleagues have provided further data from their study comparing chiropractic with hospital physiotherapy.¹ Unfortunately, their report is far from convincing. The "headline" advantage of chiropractic over hospital management at three years (29%) sounds impressive but refers to an improvement of three points on the 100 point Oswestry scale, or one and a half responses on the questionnaire. This difference may be statistically significant but is clinically trivial.

In their original article the authors noted that chiropractic was 50% more expensive,² and in the latest report they note that the chiropractic group had more treatments in the long term. The measured improvement does not seem to support such an expenditure.

The design of the study was criticised after the first paper was published.³ The patients initially presenting to a chiropractor were self selected on the basis that they believed that chiropractic would be effective (as they were expecting to pay for treatment). It is interesting that only the patients referred by chiropractors showed a significant advantage for chiropractic. Those initially recruited in hospital practice showed no significant difference. The study compares private practice with NHS treatment, with all the implications for environmental and personal factors that this brings. The follow up rate (70% and 77%) is inadequate and would preclude publication in some journals.⁴ Analysis by intention to treat does not obviate this deficiency. Important variables such as psychological disturbance are not addressed, although these are noted to have a greater impact on results than does treatment.⁵

It is disappointing that in the five years between the reports these and the other criticisms that were raised were not more fully addressed. Despite the acclamation of the first report in the popular media, I do not think that this study has advanced our understanding of the treatment of back pain in any useful way.

C G GREENOUGH
Consultant orthopaedic surgeon

Department of Orthopaedic
Surgery,
Middlesbrough General Hospital,
Middlesbrough TS5 5AZ

1 Meade TW, Dyer S, Browne W, Frank AO. Randomised comparison of chiropractic and hospital outpatient management for low back pain: results from extended follow up. *BMJ* 1995;311:349-51. (5 August.)

2 Meade TW, Dyer S, Browne W, Townsend J, Frank AO. Low back pain of mechanical origin: randomised prospective comparison of chiropractic and hospital treatment. *BMJ* 1990;300:1431-7.

3 Low back pain: comparison of chiropractic and hospital outpatient treatment [letters]. *BMJ* 1990;300:1647-50, 1723.

4 Nachemson AL, Larocca H. *Spine* 1987;12:427-9.

5 Greenough CG. Recovery from low back pain: 1-5 year follow-up of 287 injury-related cases. *Acta Orth Scand* 1993;64(suppl 254):1-34.

Authors' reply

EDITOR,—Comparing like with like is a requirement for the randomly allocated groups of patients but not for the treatments, which can be as similar or different as circumstances dictate. We compared the general approaches of chiropractic and hospital treatment for two main reasons. One reason was the absence of evidence of the efficacy of chiropractic despite its growing popularity in the 1980s. The other reason was that those whom we consulted at the outset could not agree on which specific treatments and groups of patients should be included, and only a small number of physiotherapists are fully trained in manipulation.

We welcome Alison Wakefield and Martin Bull's endorsement of our recommendation for further trials. The training of physiotherapists to which they refer is necessary for these as well as for the increasing provision of specialist treatment by spinal therapists.

All the patients initially attending chiropractors who expressed a preference for chiropractic treatment were excluded from the trial, so C G Greenough's comment on this point is misplaced. Our first report described the overall cost effectiveness of chiropractic despite the somewhat higher cost of the treatment itself and also discussed the influence of psychological factors. The authors of both letters about our paper have been selective in their quotation of our results. They have overlooked the considerable improvement in pain due to chiropractic not only early on, when over 90% of patients were followed up and additional treatment had not been given, but at all the other follow up intervals. They also do not refer to the patients' considerably greater satisfaction with chiropractic than hospital treatment regardless of whether the patients initially attended chiropractors or hospitals. Our main conclusion that chiropractic has a part to play in the management of low back pain remains unaltered.

T W MEADE

Director

SANDRA DYER

Statistician

WENDY BROWNE

Research nurse

MRC Epidemiology and Medical Care Unit,
Wolfson Institute of Preventive Medicine,
Medical College of St Bartholomew's Hospital,
London EC1M 6BQ

A O FRANK

Consultant in rheumatology and rehabilitation

Northwick Park Hospital,
Harrow,
Middlesex HA1 3UJ

Practical pulse oximetry

Saturation can be normal with dangerously high pCO₂

EDITOR,—We agree with C D Hanning and J M Alexander-Williams that pulse oximetry has a place in the management of hypoxic patients with respiratory failure and that hypercarbia is less damaging physiologically than hypoxaemia.¹ It is important to point out, however, that arterial carbon dioxide tension can become dangerously high while oxygen saturation remains normal, and this is a limitation of pulse oximetry. From the alveolar gas equations²:

$$\text{alveolar } P_{O_2} = \text{barometric pressure} \times (\text{inspired } O_2 - O_2 \text{ uptake/alveolar ventilation})$$

$$\text{alveolar } P_{CO_2} = \text{barometric pressure} \times (\text{mean inspired } CO_2 + CO_2 \text{ output/alveolar ventilation})$$

Therefore during hypoventilation the fall in alveolar oxygen tension (P_{O_2}) can be overcome by an increase in inspired oxygen while the alveolar carbon dioxide tension (P_{CO_2}) continues to rise.

For example, if a patient hypoventilates with an alveolar ventilation of 1.5 litres/min in room air the alveolar oxygen tension would be 4 kPa (assuming an oxygen consumption of 200 ml/min). If, however, the same patient was breathing 30% oxygen the alveolar oxygen tension would be 12.5 kPa, which is almost within the normal range. At this level of hypoventilation the arterial carbon dioxide tension, on the other hand, would rise to 13 kPa (two to three times normal) whatever the inspired oxygen concentration.

In patients who are hypoventilating while breathing supplemental oxygen, such as post-operative patients receiving intravenous or epidural opiates, the arterial carbon dioxide tension can climb dramatically in the face of a normal oxygen saturation. In this situation hyper-

carbia is life threatening, and as the oxygen saturation may be normal the pulse oximeter fails to warn of impending respiratory failure. Indeed, a normal saturation can lead to a false sense of security, and monitoring of respiratory rate and depth will be as valuable as pulse oximetry, if not more so, to detect impending carbon dioxide narcosis in these patients.

P M ROBBINS
Clinical fellow in anaesthesia

Harefield Hospital,
Harefield,
Middlesex UB9 6JH

M D BRUNNER
Consultant anaesthetist

Northwick Park Hospital,
Harrow,
Middlesex HA1 3UJ

1 Hanning CD, Alexander-Williams JM. Pulse oximetry: a practical review. *BMJ* 1995;311:367-70. (5 August.)

2 Nunn JF. *Applied respiratory physiology*. 4th ed. London: Butterworths, 1993.

Overnight oximetry is easy and useful

EDITOR,—Two aspects of C D Hanning and J M Alexander-Williams' review on pulse oximetry warrant further comment.¹ Firstly, the simplicity of operation of pulse oximeters may not be obvious to general readers. Not only can a pulse oximeter be used by nursing staff on the ward but patients themselves can easily be instructed in its use. This becomes important in assessing respiration during sleep. Patients can perform satisfactory overnight studies in their own home, with the advantages of being able to sleep in a familiar environment and not needing medical staff to supervise them.

Secondly, the article suggests that oximetry should be used only when sleep apnoea is suspected or the patient has important lung disease. It may also be useful, however, to assess the effect of sleep on cardiovascular disease. Clinicians are often faced with the problems of nocturnal angina or nocturnal arrhythmias, and oximetry can often give useful information and aid decisions on treatment.² We suggest that overnight oximetry should be included routinely in the investigations of these disorders.

PAUL C DEEGAN
Senior registrar

Department of Respiratory Medicine,
University Hospital,
Queen's Medical Centre,
Nottingham NG7 2UH

RICHARD LISTON
Clinical research fellow

Department of Geriatric Medicine,
Hope Hospital,
Salford M6 8HD

1 Hanning CD, Alexander-Williams JM. Pulse oximetry: a practical review. *BMJ* 1995;311:367-70. (5 August.)

2 Liston R, Deegan PC, McCreery C, McNicholas WT. Role of respiratory sleep disorders in the pathogenesis of nocturnal angina and arrhythmias. *Postgrad Med J* 1994;70:275-80.

Advance directives increase personal autonomy

EDITOR,—Ian Robertson's arguments against advance directives, or living wills, strike me as biased and specious.¹ The fact that most people who fail in attempts at suicide are later glad that they survived is irrelevant to advance directives. Most suicide attempts occur during mental states that later improve. Advance directives apply only in circumstances in which there can be no improvement.

Robertson also claims that terminally ill people do not commit suicide. This claim can be disputed. Most terminally ill people do not have the knowledge or the means to kill themselves and fear the consequences of a bungled attempt.

Robertson's argument that if there is brain damage the person occupying the body is not