or congestion, a state frequently associated with menorrhagia, and one over which medicine has little control. In such a condition two or three cold sitz-baths daily, with an occasional packing in the wet sheet, or the whole body well rubbed over with a sheet dripping in cold water, will be found most advantageous; the patient will soon improve in strength, and enjoy the remedy, which at first appeared most terrifying. Such treatment, I must say, *en passant*, would do away with much that is evil in the management of non-malignant uterine disease; it would put a stop to the too prevalent practice of speculum using, and be to delicate women a far more agreeable remedy than the weekly exposure and cauterizing, which I believe in numerous cases does great mischief.

Hysteria, in all its forms, may be combatted by the daily cold douch and sitz-baths, so as to remove the morbid susceptibility of the uterus, which is so frequently present in these cases.

In the convulsive variety of hysteria the cold dash is a specific, and indeed has been, as is well known, the remedy most relied on by the best practitioners.

In all cases of impaired health, from excess of any kind, I do not hesitate to say, that the free daily use of the water-cure processes is the most powerful agent we know of. I might enlarge as to the theory of its curative power, but many of my professional brethren around me would be better able to do justice to this part of my subject. I need not remind you of the immense importance of a healthy condition of the capillary circulation, nor of the enormous amount of excretory surface possessed by the skin; nor of the sympathy which exists between the skin and the mucous membranes; nor of the vast extent of the cuticular superficies, all which facts should teach us the value of any remedy which can be brought so exclusively to bear upon it. The uncertainty of effect in our diaphoretic medicines is too well known to require a comment, and in the proposed application of water we have a means at our disposal which brings the skin into the finest possible condition for obtaining perspiration.

A few observations on the value of water treatment in many skin diseases and I will sit down. I have employed it in psoriasis, and in acne punctata with the best results, and a case now under treatment, of severe psoriasis, combined with pityriasis, is receiving very marked benefit from the daily use of the wet sheet, and once or twice a week a sweating bath, followed by ablation in water at 60° Fahrenheit, a point worthy of attention, as water under 60°, in these cases, appears to stimulate the capillaries too powerfully, and reaction is consequently too great.

In all cases of cutaneous disease the necessity of cleanliness cannot be too much insisted on, and many patients will wash, if directed to do it, *secundum artem*, who would not be otherwise persuaded to perform their daily ablutions; indeed I need not say that the mere act of removing the cutaneous exudations is of itself sufficient curative treatment.

I have laid these few observations before the meeting

in the hope that the judicious use of water as a remedy will no longer be neglected by the profession, but that on the contrary, they will calmly look at the remedy, and, where feasible, employ it; for by so doing, they will stop the mouths of ill-conditioned persons and illiterate quacks, who now say that we will not employ a remedy which cures the patient too quickly, too simply, and without the mystery of the craft. We are, as a body, too noble-minded, too disinterested, to act thus, and I am sure if my medical brethren only believe water treatment to be good in chronic disorders, they will adopt it freely, fairly, and without reserve, not fearing on the one hand the accusation of those, who, in their dread of innovation, denounce all new modes of treatment as quackery; nor on the other, regarding the scorn of those who laugh at the failure of physic, and the seeming triumph of the empiric; for, in the words of our learned associate, Dr. Davis—"I believe we shall best consult the interest of legitimate medicine by admitting frankly whatever of truth the particular system under discussion may appear to us to contain, and by pointing out dispassionately where we consider truth to terminate and fiction to commence. By so doing, we shall let it appear that our rejection of the system has not arisen from prejudice, or from want of attention to its claims, but because it presents along with a less or greater substratum of truth, a huge superstructure of assumption and of falsehood."

ON PUERPERAL FEVER.

By EDWARD T. HUGHES, M.D., MOLD.

Read before the North Wales Branch.

DR. HUGHES said he did not propose to enter into any account of the pathology and treatment of the disease, as it would be impossible to do justice to so important a subject without trespassing too long on the time assigned to the meeting, but he took this opportunity of recording his testimony in proof of its occasional origin from the contagion of erysipelas. After briefly noticing the earlier epidemics of puerperal fever, and more particularly such as were coexistent with erysipelas, he referred, among numerous authorities, to the writings of Drs. Ramsbotham, Rigby, Copland, and Holmes, as also to the valuable communications of Mr. Storrs, of Doncaster, and others, in the Journal of the Association, for numerous and undoubted evidence on the same subject. He then reported the following as the result of his own experience:—

He attended daily, from the 30th of September to the 17th of October, 1843, two severe cases of sloughing erysipelas in the same family; both patients occupied the same apartment, and his visit never took less than twenty minutes, as in the one case the arm and fore-arm were extensively involved, and in the other the axilla and integuments of one side of the chest. On the 10th of October he attended a midwifery patient, living at least two miles and a half distant from his cases of

erysipelas, who, after a lingering first labour of twins, succeeded by considerable flooding, appeared doing well. On the 11th she had a rigor, succeeded by very profuse perspiration, rapid pulse, and tympanitic abdomen, without pain on pressure. At first these symptoms seemed to yield to treatment, but diarrhoea supervened, and she died early on the morning of the 15th. On this day he was called to a patient three miles distant from the preceding one; it was her fourth child. She had an easy labour, and was left comfortable in every respect. On the morning of the 16th she complained of a fixed pain in the lower part of the abdomen, without any degree of tension, but increased on pressure, with quick feeble pulse, anxious countenance. &c. No treatment was of the slightest avail; she gradually declined, and died at noon of the 18th.

Dr. Hughes could not now fail to recognize the source of this fatality, and also the distressing fact that he had been the vehicle of the contagion, as no other possible communication could have taken place between the affected parties, and no similar cases had occurred to any other medical man in the neighbourhood. On the 8th of October he had attended a midwifery patient, who did well, and thus on the occurrence of his first fatal case he was scarcely prepared to suspect its origin.

That erysipelas does not at all times generate this contagious poison is certain; for, independent of other authorities, Dr. Hughes, in a practice of nineteen years, has attended many more severe cases than those now reported, with impunity, and has seen erysipelas in the lying-in room, where the circumstances of the family rendered separation impossible, without any ill effects.

Regarding the power of generating puerperal fever as an occasional effect of erysipelas, under what circumstances should we most dread its baneful influence?

From the earliest periods of medicine, variable and unseasonable weather has been considered prejudicial to the recovery of lying-in women, but later experience has shown the impossibility of attaching certain importance to any such atmospheric conditions. We have it reported, in respect to the epidemic at La Maternité, in Paris, in 1829, that the puerperal fever then raging prevailed in its greatest degree during cold dry weather, during dry temperate weather, and during moist temperate weather, while at other seasons of the year the same kinds of weather did not render the disease more frequent than usual. Its malignity at one time, as reported by Dr. John Clarke, has been checked by a sharp frost, while more commonly warm weather has favoured its decline, and cold weather its increase.

Nor does the consideration of coexistent epidemics throw much more light on the subject. Undoubtedly, on many occasions, as at Aberdeen, Leeds, and Aylesbury, and more frequently in London and Dublin, erysipelas has prevailed, at the same time as puerperal fever. Still more frequently have epidemic fevers, of a typhoid type, been the concomitant diseases; but yet so many instances have occurred, especially in country practice, in which either malady has existed independently of

the other, that we cannot venture to deduce any certain practical conclusions from their occasional coexistence. When the cases Dr. Hughes has now reported occurred, low typhoid fever was prevalent in the neighbourhood of Mold, and some few cases of erysipelas of the head and face, but of no serious character, had come under treatment. In the year 1847 he attended very many more, ill of both diseases, and experienced no injurious effects in his midwifery practice.

As the preceding observations are intended to show that the causes which favour the origin of puerperal fever from the contagion of erysipelas are as yet obscure, he can only repeat the recommendation to medical men when attending such cases, carefully to watch the recoveries in the lying-in room, and at the same time strictly to observe those precautions which their duties, and the well-doing of their patients, peremptorily demand of them.

Mold, June 12, 1850.

ON THE STRUCTURE OF THE MEMBRANA TYMPANI IN THE HUMAN EAR.*

By JOSEPH TOYNBEE, Esq., F.R.S.

FELLOW OF THE ROYAL COLLEGE OF SURGEONS IN ENGLAND, AND SENIOR
SURGEON TO ST. GEORGE'S AND ST. JAMES'S DISPENSARY, LONDON.

IN this paper the membrana tympani is described as consisting of the following layers, which are quite distinct from each other, both as regards their structure and functions:—

1. Epidermis.
2. The proper fibrous layer, composed of,—
 - a The lamina of radiating fibres.
 - b The lamina of circular fibres.
3. Mucous membrane.

One of the principal objects of the paper is to describe the structure and functions of the fibrous laminae. Since the time of Sir Everard Home, who pronounced the layer of radiating fibres to be muscular, anatomists have differed in their views of the nature of the fibrous element of the membrana tympani. The lamina of radiating fibres, the outer surface of which is covered by the epidermis, is described as continuous with the periosteum of the external meatus. With the exception of the uppermost fibres which, on account of their being somewhat flaccid, have been considered as a separate tissue, under the name of "membrana flaccida," the radiate layer is composed of fibres which extend from the circular cartilaginous ring to the malleus, and they interlace in their course. These fibres are from 4000 to 5000 parts of an inch in breadth.

The lamina of circular fibres consists of circular fibres which are firm and strong towards the circumference, but very attenuated towards the centre. These

* The paper, of which a short abstract is here published, was read at the Royal Society, on the 20th of June.