Studies on Attenuated Measles-Virus Vaccines in Canada *

F. P. NAGLER,¹ A. R. FOLEY,² J. FURESZ³ & G. MARTINEAU⁴

This paper describes the results of a study of live attenuated measles vaccines (one in a series of WHO-sponsored field trials) carried out in children 6-33 months old at an orphanage in Quebec City. The Enders Edmonston B vaccine alone and the same vaccine administered with gamma-globulin were compared with the Schwarz further-attenuated vaccine. The over-all seroconversion rates were found to be 96.9 %, 98.1 % and 98.8 % respectively. Severe clinical reactions, except for high fever, were not observed in any of the groups. Rectal temperatures over 103°F (39.5°C) were noted in 16.2 % of the children given Schwarz vaccine, in 59.2 % of the children receiving the Enders Edmonston B vaccine alone and in 27.8 % of the children inoculated with the Enders Edmonston B vaccine plus gamma-globulin. The high incidence of mild pharyngitis following inoculation of these vaccines was not observed in the group of children who had received vaccine plus gamma-globulin. No significant differences were noted in the frequency of other symptoms, such as cough, coryza, conjunctivitis and diarrhoea, between vaccinated and control groups.

A measles vaccine for routine use should provide effective protection without producing undesirable reactions such as high fever, rash or involvement of the central nervous system. Live measles vaccine developed by Enders produces excellent antibody response and protection in humans but causes clinical reactions in most of those vaccinated (Enders, 1960). These reactions can be reduced, but not eliminated, by the simultaneous injection of gamma-globulin with the vaccine. Recently, several further-attenuated measles vaccines have been

developed, and the results of preliminary studies on these vaccines have been very encouraging (Andelman et al., 1963; Hendrickse et al., 1964). The WHO Scientific Group on Measles Vaccine Studies, whose report was published in 1963, recommended in July of that year that the World Health Organization should stimulate, co-ordinate and support a series of field trials on the further-attenuated measles vaccines in a number of countries with different climates. The present paper describes the results of one of these trials, which was carried out in Canada in 1964.

METHODS AND MATERIALS

PLAN OF STUDY

The study was carried out in healthy infants and young children in an orphanage at Quebec City. The first part of the study was undertaken in March and April 1964 (spring trial), and the second part in September and October 1964 (fall trial).

Spring trial

In all, 193 children, 8-33 months of age, were chosen for vaccination. The children were divided at random into five groups. Group I received Enders Edmonston B vaccine with gamma-globulin, Group II received Schwarz further-attenuated

* This investigation was carried out under WHO auspices and was supported mainly by a grant from the Department of National Health and Welfare, Ottawa, Canada (Grant No. 604-7-452), and in part by a grant from the National Institutes of Health of the United States Public Health Service, Bethesda, Md. (Grant No. AI-03817) to the World Health Organization.

¹ Chief, Virus Laboratories, Laboratory of Hygiene, Department of National Health and Welfare, Ottawa, Canada.

² Chief Epidemiologist, Ministry of Health, Province of Quebec, and Professor of Epidemiology, University of Montreal, Canada.

³ Medical Officer, Virus Laboratories, Laboratory of Hygiene, Department of National Health and Welfare, Ottawa, Canada.

⁴ Assistant Epidemiologist, Ministry of Health, Province of Quebec, Canada.

vaccine, Group III was given Enders Edmonston B alone, Group IV received a placebo and Group V received Beckenham 20 further-attenuated vaccine.

Fall trial

Altogether 165 children, 7-27 months of age, were chosen for vaccination. The children were divided at random into four groups. Group I received Enders Edmonston B vaccine alone, Group II the Schwarz vaccine, Group III Enders Edmonston B plus gamma-globulin and Group IV a placebo injection. In addition to these groups, a group of 29 infants 6 months old received Schwarz vaccine. Beckenham 20 vaccine free of avian leucosis virus was not available for this trial.

In both trials the children were bled one week before and one month after vaccination. Of the 193 children vaccinated in the spring trial 7 had prevaccination antibodies to measles virus. Of the 165 children vaccinated in the fall trial 6 had prevaccination antibodies. The postvaccination sera of these children showed no increase in antimeasles titres and the children were excluded from both studies. To eliminate bias, one team of medical officers allocated the children to the different groups and performed their inoculation, while another team of medical officers (paediatricians) carried out the follow-up observation. Only children in good health and with a rectal temperature of under 100°F (37.8°C) were vaccinated. After vaccination the children were examined daily by the paediatricians for a period of 30 days; the body temperature was taken rectally every day between 4 p.m. and 6 p.m. The daily observations were documented carefully in individual follow-up record cards designed by WHO. Four children in the spring trial left the orphanage some days before the end of the trial and were not included in the clinical follow-up. However, these children were brought back to the orphanage at the end of the trial, and blood samples were taken to establish their serological response.

VACCINES AND GAMMA-GLOBULINS

The Enders Edmonston B vaccine (Lot No. 91055/70005) was supplied by Merck, Sharp & Dohme Laboratories, West Point, Pa., USA. The Schwarz further-attenuated vaccine (Batch No. 18 for experimental use, derived from the Edmonston A strain and passed 85 additional times in chick-embryo tissue cultures at 32°C) was received from Glaxo Laboratories, Greenford, Middlesex, England.

The Beckenham 20 vaccine (Lot No. VO2820—a descendant of the Edmonston B strain, passed 77 additional times in chick-embryo tissue cultures at 32°C) was supplied by Parke, Davis Laboratories, Detroit, Mich., USA. The bulk material of each vaccine was tested in the Ottawa laboratories for safety in animals and for potency in FL human amnion and BS-C-1 African green monkey kidney tissue cultures. The ampouled material of the Enders Edmonston B and Schwarz (Glaxo) vaccines was also tested for potency and was found to contain approximately 500 TCID₅₀ and 100 000 TCID₅₀ per 0.5 ml respectively. The shipment of the ampouled material of the Beckenham 20 vaccine (Parke, Davis) was received only a few days before the spring trial commenced. The viability of this vaccine had been destroyed during the process of ampouling, but this fact became known only after the children had been vaccinated. The bulk material of this vaccine had not suffered any measurable loss of viable virus on storage for a subsequent period of six months.

The gamma-globulins used in both trials were supplied by Merck, Sharp and Dohme Laboratories and were tested for potency in the Ottawa laboratories. Lot No. 56831B-1 (Poliomyelitis Immune Globulin, Human), used in the spring trial, showed a measles neutralizing antibody titre of 1:388/0.1 ml when tested in FL human amnion tissue cultures. Lot No. 74332B (Gammagee), employed in the fall trial, had a measles neutralizing antibody titre of 1:850/0.1 ml.

The placebo material consisted of physiological saline supplied by the Connaught Medical Research Laboratories, Toronto.

ADMINISTRATION OF THE VACCINE

Before shipment of the vaccines and placebo to Quebec City the original labels of all vials were replaced by labels showing only a code letter. Vaccines were stored at -20°C prior to use. Enders Edmonston B and Schwarz vaccines were supplied in lyophilized form and the vaccines were reconstituted immediately before use with distilled water supplied with the vaccine by the manufacturers. The vaccines were administered subcutaneously in 0.5-ml amounts into the left deltoid area using individual disposable syringes and needles. Children receiving Enders Edmonston B vaccine plus gamma-globulin were inoculated with 1.0 ml of vaccine (on the recommendation of the manufacturer) into the

left arm and 0.01 ml of gamma-globulin/lb (0.45 kg) body-weight by separate syringe into the right arm. The placebo was inoculated in 0.5-ml amounts into the left arm of each child in the control groups.

SERUM ANTIBODY TITRATIONS

Prevaccination and postvaccination sera were simultaneously titrated using the haemagglutination-inhibition (HI) and serum-neutralization tests. The HI test was performed by Rosen's method (Rosen, 1961) adapted to Takátsy's micro-technique (Takátsy, 1955; Sever, 1962). Since none of the 400 prevaccination sera tested contained non-specific haemagglutination inhibitors, the kaolin treatment of the sera was omitted. All prevaccination and

postvaccination sera were tested for non-specific haemagglutinins using African green monkey red blood cells. Sera containing such haemagglutinins were adsorbed with *Cercopithecus* erythrocytes.

In the serum-neutralization test, FL tissue cultures were used. Neutralization of the "wild" Edmonston strain (30-150 TCID₅₀) was carried out for 2 hours at 27°C and the serum-virus mixtures were inoculated in 0.4-ml amounts into two culture tubes containing 1.6 ml of fresh M150 medium (Morgan, Campbell & Morton, 1955) plus 2% calf serum. The inoculated tube cultures (kept stationary) were incubated at 37°C, and serum titres were calculated from the 12th-day reading by Kärber's method (Kärber, 1931). All pre- and postvaccination sera were titrated in 1 : 4 and 1 : 16 dilutions.

RESULTS

SPRING TRIAL

Clinical reactions

A comparison of the frequencies of fever in the five groups (Table 1) indicates that the Enders

Edmonston B vaccine (alone) produced a significantly higher incidence than the Schwarz vaccine. The gamma-globulin administered with the former vaccine reduced but did not eliminate the pyrexial response of the children. The Beckenham 20

TABLE 1
FEBRILE REACTIONS OBSERVED IN VACCINATED INFANTS, SPRING TRIAL

Observations	Enders Edmonston B alone		Enders Edmonston B plus gamma-globulin		Schwarz		Beckenham 20 ^a		Placebo	
	Age-group (months)		Age group (months)		Age-group (months)		Age-group (months)		Age-group (months)	
	8-12	13-27	8-12	13-28	8-12	13-33	8-12	13-27	8-12	13-23 ^b
Total number in groups	14	18	13	21	18	21	18	24	10	25
Temperature (% distribution) ^c :										
Under 101°F (38.3°C)	7.2	0.0	38.4	19.1	27.8	33.3	89.0	87.5	100	80.0
101°F-102.9°F (38.3°C-39.4°C)	35.7	38.9	30.8	38.1	66.6	42.9	5.5	12.5	0	20.0
103°F (39.5°C) and over	57.1	61.1	30.8	42.8	5.6	23.8	5.5	0	0	0
Mean maximum temperature 1-6 days after vaccination (°F(°C))	99.9 (37.72)	100.5 (38.05)	99.8 (37.67)	100.2 (37.89)	100.2 (37.89)	100.2 (37.89)	99.8 (37.67)	100.1 (37.83)	100.4 (38.00)	100.1 (37.83)
Mean maximum temperature 7-14 days after vaccination (°F(°C))	103.3 (39.61)	103.2 (39.56)	101.7 (38.72)	102.4 (39.11)	101.4 (38.56)	101.6 (38.67)	100.3 (37.94)	100.4 (38.00)	100.0 (37.78)	100.2 (37.89)
Mean duration of fever, 101°F (38.3°C) and over (days) ^d	2.7 (1-4)	2.5 (1-5)	2.3 (1-4)	2.2 (1-4)	1.7 (1-3)	1.7 (1-3)	1.5 (1-2)	1.0	—	1.0
Mean onset of fever, 101°F(38.3°C) and over (days) ^d	8.7 (8-9)	8.7 (8-10)	10 (8-13)	8.4 (7-11)	8.3 (7-10)	8.1 (7-9)	10.5 (8-13)	10.3 (9-12)	—	10 (7-12)

^a Vaccine contained no live virus.

^b One child was 39 months old.

^c Rectal.

^d Range shown in parentheses.

TABLE 2
CLINICAL SYMPTOMS OBSERVED IN INFANTS 7-14 DAYS AFTER VACCINATION, SPRING TRIAL

Observations	Enders Edmonston B alone		Enders Edmonston B plus gamma-globulin		Schwarz		Beckenham 20 ^a		Placebo	
	Age-group (months)		Age-group (months)		Age-group (months)		Age-group (months)		Age-group (months)	
	8-12	13-27	8-12	13-28	8-12	13-33	8-12	13-27	8-12	13-23
Total number in groups	14	18	13	21	18	21	18	24	10	25
Percentage distribution:										
Rash	50.0	22.2	23.1	23.8	33.3	23.8	5.5	0	0	0
Cough	35.7	16.7	30.8	9.5	22.3	14.3	27.7	16.6	10.0	12.0
Coryza	21.4	72.2	15.4	38.1	33.3	14.3	27.7	8.3	30.0	8.0
Conjunctivitis	28.6	11.1	15.4	0	16.6	4.8	2.2	0	30.0	12.0
Pharyngitis	57.1	38.8	7.7	19.0	33.3	23.8	11.1	8.3	0	4.0
Mean duration (days):										
Rash	2.3	1.8	2.7	2.2	2.0	1.6	1.0	—	—	—
Conjunctivitis	1.8	2.0	3.3	—	1.3	1.0	3.0	—	1.7	2.3
Pharyngitis	1.3	1.0	2.0	1.5	1.5	1.0	1.0	1.0	—	1.0
Mean onset (days):										
Rash	10.4	11.3	11.0	10.6	9.5	10.8	9.0	—	—	—
Conjunctivitis	9.5	10.0	9.5	—	8.7	10.0	8.5	—	9.7	10.7
Pharyngitis	9.9	9.1	9.0	8.3	9.2	9.0	9.0	8.5	—	9.0
Convulsions (number of cases)	—	1	—	—	—	—	—	—	—	—

^a Vaccine contained no live virus.

vaccine contained no live virus, consequently the incidence of fever in this group was similar to that of the control group. The means of the maximum temperatures taken 1-6 days after vaccination indicate that the vaccines did not produce fever during the first 6 days in any of the groups. However, fever appeared on the 7th-8th day and persisted for 1-5 days in the vaccinated groups. The means of the maximum temperatures taken 7-14 days after vaccination and the mean duration of fever in days were the lowest in the group that had received the Schwarz vaccine.

Additional clinical symptoms observed in infants 7-14 days after vaccination are summarized in Table 2. The frequency of rash was approximately the same in the three groups receiving live virus vaccines, with the exception of the group of children aged 8-12 months given Enders Edmonston B alone,

which showed an incidence of 50%. The mean onset of rash varied from 9.5 to 11.3 days in these groups. The rash appeared to be much milder than the rash associated with natural measles, with a limited distribution on the face, neck and trunk. None of the children showed Koplik's spots. Other symptoms, such as cough, coryza and conjunctivitis, appeared with about equal frequency in all five groups. Enders Edmonston B vaccine (alone) produced a somewhat higher incidence of mild pharyngitis (without follicular tonsillitis) than the Schwarz vaccine. The administration of gamma-globulin greatly reduced the incidence of pharyngitis.

Convulsions were observed in only one child, who had been vaccinated with Enders Edmonston B vaccine. This 14-month-old boy developed a fever of 103.4°F (39.6°C) on the 8th day after vaccination and his temperature rose to 105°F (40.5°C) the

TABLE 3
ANTIBODY RESPONSE OF INFANTS TESTED ONE MONTH AFTER VACCINATION, SPRING TRIAL

Observations	Enders Edmonston B alone	Enders Edmonston B plus gamma-globulin	Schwarz	Beckenham 20	Placebo
Total number in groups (susceptible to measles)	32	35	39	43	37
Number of children showing post-vaccination serum-neutralization titre:					
< 1 : 4	2	1	1	43	37
1 : 4 to 1 : 16	0	2	1	0	0
≥ 1 : 32	30 (93.9 %)	32 (91.4 %)	37 (94.9 %)	0	0
Serological conversion (% distribution, ≥ 1 : 4)	93.8	96.2	97.6	0	0

following day. A transient convulsion lasting for half an hour was observed at that time. On the 10th day his temperature returned to normal and the child showed no signs of illness. An electroencephalogram taken three days after his convulsions showed no abnormalities.

Antibody response to the vaccines

The results of the trial (which are summarized in Table 3) indicated that the Enders Edmonston B vaccine, administered alone or with gamma-globulin, and the Schwarz vaccine produced antibody responses in, respectively, 93.8%, 96.2% and 97.6% of children having no prevaccination antibodies. Good antibody response (serum-neutralization titre ≥ 1 : 32) occurred in, respectively, 93.8%, 91.4% and 94.9%. Children inoculated with the Beckenham 20 vaccine or with the placebo material showed no antibody response to measles virus.

FALL TRIAL

Clinical reactions

The frequency of fever in the four groups is shown in Table 4. The highest incidence of fever (103°F (39.5°C) and over) occurred again in the Enders Edmonston B group. The incidence of high fever was approximately the same in the group receiving Schwarz vaccine and the group given Enders Edmonston B vaccine plus gamma-globulin. Infants less than 12 months old showed higher pyrexial responses than those older than 12 months. This difference was not seen in the spring trial. The means of maximum temperatures taken 7-14 days

after vaccination in each group were similar to the mean temperatures observed in the corresponding groups during the spring trial.

The incidence of rash in each group of the fall trial was significantly lower than that observed at the spring trial (Table 5). Other symptoms, such as cough, coryza and conjunctivitis, revealed only insignificant differences between control and vaccinated groups. The administration of gamma-globulin reduced the incidence of pharyngitis to about the same level as that noted in the control group.

Antibody response to the vaccines

Each vaccinated group showed a serological conversion rate of 100% (Table 6). However, serum-neutralization titres of ≥ 1 : 32 were observed in a smaller proportion of children at the fall trial than at the spring trial.

Vaccination of 6-month-old infants with Schwarz vaccine

In the fall trial an additional group of 29 6-month-old infants was inoculated with the Schwarz vaccine. The prevaccination sera of these children were tested for measles antibodies using the HI and serum-neutralization tests. Antibodies were revealed in 4 sera by the HI test and in 6 sera by the neutralization test. However, when the remaining 23 prevaccination sera were retested by the neutralization test using a very small dose of virus (3-10 TCID₅₀), 5 additional sera were found to contain antibodies to measles virus. Thus, 11 of 29 infants appeared to have maternal antibodies at the time of vaccination.

TABLE 4
FEBRILE REACTIONS OBSERVED IN VACCINATED INFANTS, FALL TRIAL

Observations	Enders Edmonston B alone		Enders Edmonston B plus gamma-globulin		Schwarz		Placebo	
	Age-group (months)		Age-group (months)		Age-group (months)		Age-group (months)	
	7-12	13-24	7-12	13-25 ^a	7-12	13-25	7-12	13-27
Total number in groups	20	17	19	18	24	19	24	18
Temperature (% distribution): ^b								
Under 101°F (38.3°C)	10	17.7	31.6	66.7	45.9	31.6	79.2	77.8
101°F-102.9°F (38.3°C-39.4°C)	30	23.5	42.1	22.2	29.1	57.9	12.5	22.2
103°F (39.5°C) and over	60	58.8	26.3	11.1	25.0	10.5	8.3	0
Mean maximum temperature, 1-6 days after vaccination (°F(°C))	100.1 (37.83)	100.5 (38.05)	100.1 (37.83)	100.4 (38.00)	100.6 (38.11)	100.8 (38.22)	100.5 (38.05)	100.2 (37.89)
Mean maximum temperature 7-14 days after vaccination (°F(°C))	103.0 (39.44)	102.8 (39.33)	101.9 (38.83)	100.9 (38.27)	101.4 (38.56)	101.6 (38.67)	100.5 (38.05)	100.4 (38.00)
Mean duration of fever, 101°F (38.3°) and over (days) ^c	2.8 (1-4)	3.1 (2-6)	2.1 (1-4)	2.5 (1-4)	2.0 (1-5)	1.8 (1-3)	2.6 (1-5)	1.8 (1-3)
Mean onset of fever, 101°F (38.3°C) and over (days) ^c	8.8 (7-11)	8.1 (7-9)	10 (8-12)	9.0 (7-11)	7.8 (7-11)	8.6 (7-10)	10.6 (7-14)	12.0 (10-13)

^a One child was 45 months old.

^b Rectal.

^c Range shown in parentheses.

The clinical symptoms observed in 28 of these infants are summarized in Table 7. One infant left the orphanage 10 days after vaccination but returned at the end of the trial for his postvaccination blood sample. In order to compare the results obtained in this group with those obtained in infants aged 7-12 months who had been vaccinated with the same vaccine, the clinical symptoms of the latter group and those of the control group (Tables 4 and 5) are also included in Table 7. The frequency of high fever (103°F (39.5°C) and over) following vaccination was about the same in the infants aged 6 months as in those aged 7-12 months having no prevaccination antibodies. However, 70% of the 6-month-old infants with maternal antibodies showed a fever of 101°F-102.9°F (38.3°C-39.4°C) 7-14 days after vaccination. It may be assumed that the fever was due to the vaccination, since only 20.8% of the placebo group showed fever over 101°F (38.3°C) and only a very few children with slightly elevated temperatures were found in any

of the groups during the first week after vaccination (see Table 7). The onset of fever was delayed in the 6-month-old group with maternal antibodies.

The antibody response to Schwarz vaccine in the 6-month-old infants is compared with that in the infants aged 7-12 months in Table 8. Of the 11 infants having prevaccination antibodies to measles virus, 8 showed no change in serum antibody titre one month after vaccination, while 3 showed no antibodies after vaccination, although all three had prevaccination antibodies. This is probably due to the normal decline of maternal antibodies in the second 6 months of life.

Of the 18 6-month-old infants without maternal antibodies, 17 showed an antibody response to the vaccine (94.4% conversion rate). Only 16.6% of this group had a good antibody response (serum neutralization titre $\geq 1:32$) as compared with 66.7% of the group aged 7-12 months. This may have been due to the presence of low-level maternal antibodies in these infants not detectable by the tests employed in this study.

TABLE 5
CLINICAL SYMPTOMS OBSERVED IN INFANTS 7-14 DAYS AFTER VACCINATION, FALL TRIAL

Observations	Enders Edmonston B alone		Enders Edmonston B plus gamma-globulin		Schwarz		Placebo	
	Age-group (months)		Age-group (months)		Age-group (months)		Age-group (months)	
	7-12	13-24	7-12	13-25	7-12	13-25	7-12	13-27
Total number in groups	20	17	19	18	24	19	24	18
Percentage distribution:								
Rash	5.0	0	0	0	4.2	10.5	0	0
Cough	20.0	23.5	26.3	22.2	33.3	31.6	20.8	16.6
Coryza	15.0	41.2	26.3	38.9	33.3	31.6	12.5	16.6
Conjunctivitis	30.0	17.6	5.3	16.6	12.5	15.8	12.5	11.1
Pharyngitis	35.0	41.2	10.5	16.6	33.3	31.6	12.5	16.6
Mean duration (days):								
Rash	4.0	—	—	—	5.0	3.0	—	—
Conjunctivitis	2.7	2.7	1.0	1.2	3.7	1.3	2.7	1.5
Pharyngitis	1.1	1.0	1.0	1.0	1.5	1.0	2.7	2.7
Mean onset (days):								
Rash	10.0	—	—	—	9.0	11.0	—	—
Conjunctivitis	8.8	10.0	9.0	7.2	7.7	7.7	13.0	7.5
Pharyngitis	8.7	8.6	9.0	9.0	8.9	8.8	8.0	9.7
Convulsions (number of cases)	—	—	—	—	—	—	—	—

TABLE 6
ANTIBODY RESPONSE OF INFANTS TESTED ONE MONTH AFTER VACCINATION, FALL TRIAL

Observations	Enders Edmonston B alone	Enders Edmonston B plus gamma-globulin	Schwarz	Placebo
Total number in groups (susceptible to measles)	39	37	42	45
Number of children showing postvaccination serum-neutralization titre:				
<1:4	0	0	0	45
1:4 to 1:16	5	17	10	0
≥1:32	34 (87.2 %)	20 (54.1 %)	32 (76.2 %)	0
Serological conversion (% distribution, ≥1:4)	100	100	100	0

TABLE 7
COMPARISON OF CLINICAL SYMPTOMS OBSERVED IN INFANTS AGED 6 MONTHS
AND INFANTS AGED 7-12 MONTHS VACCINATED WITH SCHWARZ VACCINE, FALL TRIAL

Observations	Schwarz			Placebo
	Infants aged 6 months		Infants aged 7-12 months	Infants aged 7-12 months
	With prevaccination antibodies ^a	Without prevaccination antibodies ^a		
Total number in groups	10	18	24	24
Temperature (% distribution):				
Under 101°F (38.3°C)	30	33.3	45.9	79.2
101°F-102.9°F (38.3°C-39.4°C)	70	38.9	29.1	12.5
103°F (39.5°C) and over	0	27.8	25.0	8.3
Mean maximum temperature, 1-6 days after vaccination (°F(°C))	100.3 (37.94)	100.5 (38.05)	100.6 (38.11)	100.5 (38.05)
Mean maximum temperature, 7-14 days after vaccination (°F(°C))	101.6 (38.67)	101.8 (38.78)	101.4 (38.56)	100.5 (38.05)
Mean duration of fever, 101°F (38.3°C) and over (days) ^b	2.0	2.3	2.0	2.6
Mean onset of fever, 101°F (38.3°C) and over (days) ^b	12.0 (10-14)	9.7 (7-14)	7.8 (7-11)	10.6 (7-14)
Rash (% distribution)	0	0	4.2	0
Pharyngitis (% distribution)	0	22.2	33.3	12.5

^a Serum samples were obtained one week before vaccination.

^b Range shown in parentheses.

TABLE 8
COMPARISON OF ANTIBODY RESPONSES IN INFANTS AGED 6 MONTHS
AND INFANTS AGED 7-12 MONTHS, FALL TRIAL

Observations	Schwarz			Placebo
	Infants aged 6 months		Infants aged 7-12 months	Infants aged 7-12 months
	With pre-vaccination antibodies ^a	Without pre-vaccination antibodies ^a		
Total number in groups	11	18	24	24
Number of children showing post-vaccination serum-neutralization titre:				
< 1:4	3	1	0	24
1:4 to 1:16	8	14	8	0
≥ 1:32	0	3 (16.6 %)	16 (66.7%)	0
Serological conversion (% distribution, ≥ 1:4)	0	94.4	100	0

^a Serum samples were obtained one week before vaccination

DISCUSSION

The present study on Enders Edmonston B vaccine and the Schwarz further-attenuated vaccine (the results obtained with the Beckenham 20 further-attenuated vaccine had to be disregarded owing to the absence of viable virus in the vaccine) indicates that severe clinical reactions, except for high fever, were uncommon in all vaccinated groups. Only one 14-month-old infant had convulsions, which, however, appeared to be related to the high fever and not to any specific involvement of the central nervous system.

The combined results of the spring and fall trials show that both Enders Edmonston B vaccine (with and without gamma-globulin) and Schwarz vaccine (without gamma-globulin) had a pronounced immunogenic effect. Of the 119 children in the control group (including also the 42 children of the spring trial who received the inactive Beckenham 20 vaccine), 14.4% had a temperature rise to more than 101°F (38.3°C). This figure is in sharp contrast to the rates of 91.2%, 61.1% and 65.3% found in the groups that had received the Enders Ed-

monston B vaccine alone, the Enders Edmonston B vaccine plus gamma-globulin and the Schwarz vaccine, respectively. The Schwarz vaccine caused significantly fewer cases of high fever (103°F (39.5°C) and over) (16.2%) than the Enders Edmonston B vaccine alone (59.2%) or the Enders Edmonston B vaccine plus gamma-globulin (27.8%). Schwarz (1962) reported in the preliminary tests of his vaccine that high fever (103°F and over) occurred in 5.7% of the vaccinated children. This figure varied from 2.5% (Andelman et al., 1963) to 4.6% (Morley et al., 1964) and to 11.2% (Schwarz, 1964) in later studies. Krugman et al. (1963) found that, using the Schwarz vaccine, the percentage incidence of high fever in infants between 1½ and 2 years of age was 35%—a rate significantly higher than the 15% febrile reaction rate observed in older children. An analysis of the pyrexial responses included in Tables 1 and 4 reveals that 5.6% and 25% of infants less than 12 months old and 23.8% and 10.5% of infants older than 12 months showed febrile reactions in the spring and fall trials, respectively. The cause

TABLE 9
SUMMARY OF CLINICAL AND SEROLOGICAL RESPONSES IN VACCINATED INFANTS,
SPRING AND FALL TRIALS COMBINED

Observations	Enders Edmonston B alone	Enders Edmonston B plus gamma- globulin	Schwarz	Placebo ^a
	Age-group (months)	Age-group (months)	Age-group (months)	Age-group (months)
	7-27	7-28	7-33	7-27
Total number in groups	69 (71) ^b	71 (72) ^b	82 (81) ^b	119 (125) ^b
Temperature (% distribution):				
Under 101°F (38.3°C)	8.8	38.9	34.7	85.6
101°F-102.9°F (38.3°C-39.4°C)	32.0	33.3	49.1	12.1
103°F (39.5°C) and over	59.2	27.8	16.2	2.3
Mean maximum temperature (°F-(°C))	103.1 (39.50)	101.7 (38.72)	101.5 (38.61)	100.3 (37.94)
Rash (% distribution)	19.3	11.7	18.0	0.9
Conjunctivitis (% distribution)	21.8	9.4	12.4	11.3
Pharyngitis (% distribution)	43.0	13.5	30.5	13.1
Serological conversion (% distribution)	96.9	98.1	98.8	0

^a The placebo group in this table includes also the group vaccinated with Beckenham 20 vaccine, which contained no live virus.

^b First figure represents the number of infants observed clinically; figure in parentheses represents the number of paired sera tested serologically.

of this variation in febrile reaction rates between age-groups is not clear. However, the over-all rate (16.2%) was higher in these studies than in those reported by other workers (Morley et al., 1964; Schwarz, 1964). Similarly, the febrile reaction rates were also higher in the groups receiving Enders Edmonston B vaccine alone (over-all rate, 59.2%) than the corresponding rates (20%-40%) reported in earlier studies (WHO Scientific Group on Measles Vaccine Studies, 1963). When gamma-globulin was administered with the vaccine, the incidence of high fever was significantly reduced (over-all rate, 27.8%). It is noteworthy that this reduction was more apparent in the fall than in the spring trial. This was most probably due to the higher potency of the gamma-globulin used in the fall trial. The gamma-globulin used in the spring trial was compared with the US National Institutes of Health reference measles serum and was found to contain antibodies for measles virus below the minimum permissible level required by the US Public Health Service Regulations (United States Public Health Service, 1963).

The frequency of rash was significantly lower in the fall trial (0-10.5%) than in the spring trial (22.2%-50%). The highest percentage occurred in the group given Enders Edmonston B vaccine alone, while the lowest occurred in the group receiving

Enders Edmonston B vaccine plus gamma-globulin.

There was no significant difference in the frequency of other symptoms, such as cough, coryza, conjunctivitis or diarrhoea, between control and vaccinated groups. While the administration of gamma-globulin with Enders Edmonston B vaccine reduced the rate of mild pharyngitis to the level found in the control group (see Table 9—13.5% compared with 13.1% of the control group), the percentage distribution in the groups given Enders Edmonston B vaccine alone and Schwarz vaccine was found to be high (43% and 30.5% respectively).

Of the 29 infants 6 months old inoculated with Schwarz vaccine, 11 had maternal antibodies prior to vaccination. These infants did not show an antibody response to the vaccine but 7 reacted with a temperature of 101°F-102.9°F (38.3°C-39.4°C) 7-14 days after vaccination. Of the 18 6-month-old infants having no maternal antibodies, 17 revealed a serological response to the vaccine. However, only 16.6% of this group showed a good antibody response (serum neutralization titre $\geq 1:32$), as compared with 66.7% of the group aged 7-12 months given the same vaccine (Table 8). The frequencies of high fever and mild pharyngitis were approximately the same in infants of both age-groups having no maternal antibodies.

ACKNOWLEDGEMENTS

The authors are indebted to WHO for advice and assistance in planning this study and for the supply of record forms. The excellent co-operation of the following persons on the staff of La Crèche, St Vincent de Paul, Quebec, is gratefully acknowledged: Dr D. Lapointe,

Medical Director; Dr E. Déchène and Dr A. LaRue, Paediatricians; Marie de Jessé, Director of Nursing; and Miss M. Bergeron, Head-Nurse. The valuable technical assistance of Mrs C. R. Green, Mr R. A. J. Lapointe and Mr G. Labelle is greatly appreciated.

RÉSUMÉ

Dans le cadre d'une série de recherches patronnées par l'OMS, les vaccins antirougeoleux vivants atténués ont fait l'objet de deux enquêtes à Québec (Canada) au cours de l'année 1964.

La première a eu lieu en mars-avril: au total, 193 enfants âgés de 8-33 mois, répartis au hasard en cinq groupes, ont été vaccinés au moyen de l'une des préparations suivantes: vaccin Enders de souche Edmonston B avec gamma-globuline, vaccin Schwarz, vaccin Enders seul, vaccin Beckenham 20; le cinquième groupe a reçu du placebo. On observa un changement du sens de la

séroréaction chez respectivement 93,8%, 96,2% et 97,6% des enfants vaccinés par le vaccin Enders seul, le vaccin Enders avec gamma-globuline et le vaccin Schwarz. En dehors d'une fièvre élevée, on ne décéla de réaction clinique grave dans aucun des groupes. L'administration du vaccin Beckenham 20 ne détermina aucune réponse en anticorps pour le virus morbilleux.

Au cours de la seconde enquête en septembre-octobre de la même année, 165 enfants âgés de 7-27 mois reçurent à leur tour, soit du vaccin Enders seul, soit du vaccin Enders avec gamma-globuline, du vaccin Schwarz ou du

placebo. Le vaccin Enders sans gamma-globuline entraîna les réactions fébriles les plus intenses. Dans les autres groupes, les réactions fébriles post-vaccinales furent moins fortes et d'une intensité sensiblement égale. La vaccination suscita très généralement l'apparition d'anticorps, mais les titres élevés furent moins nombreux que lors de la vaccination effectuée au printemps.

Les résultats globaux des deux enquêtes montrent que les réactions fébriles importantes (plus de 39,5°C) sont moins fréquentes (16,2%) après vaccination par le vaccin Schwarz qu'après utilisation du vaccin Enders seul (59,2%), ou avec gamma-globuline (27,8%). Le vaccin Schwarz provoqua une éruption légère chez 18% des

enfants vaccinés, les chiffres correspondants, pour les vaccins Enders seul ou avec gamma-globuline étant de 19,3% et 11,7% respectivement. Chez les enfants recevant de la gamma-globuline en même temps que le vaccin Enders, on ne nota qu'une faible proportion (13,5%) de cas de pharyngite; en comparaison, les enfants vaccinés par le vaccin Enders sans gamma-globuline et par le vaccin Schwarz furent atteints de cette complication post-vaccinale dans la proportion de 43% et 30,5% respectivement.

Il n'y eut guère de différences quant la fréquence et la gravité des autres réactions (toux, coryza, conjonctivite, diarrhée) en fonction des diverses préparations vaccinales utilisées.

REFERENCES

- Andelman, S. L., Schwarz, A., Andelman, M. B. & Zackler, J. (1963) *J. Amer. med. Ass.*, **184**, 721
 Enders, J. F. (1960) *New Engl. J. Med.*, **263**, 153
 Hendrickse, R. G., Montefiore, D., Sherman, P. M. & Van der Wall, H. M. (1964) *Brit. med. J.*, **1**, 470
 Kärber, G. (1931) *Naunyn-Schmiedeberg's Arch. exp. Path. Pharmac.*, **162**, 480
 Krugman, S., Giles, J. P., Jacobs, A. M. & Friedman, H. (1963) *Pediatrics*, **31**, 919
 Morgan, J. F., Campbell, M. E. & Morton, H. J. (1955) *J. nat. Cancer Inst.*, **16**, 557
 Morley, D. C., Woodland, M., Krugman, S., Friedman, H. & Grab, B. (1964) *Bull. Wld Hlth Org.*, **30**, 733
 Rosen, L. (1961) *Virology*, **13**, 139
 Schwarz, A. J. F. (1962) *Amer. J. Dis. Child.*, **103**, 386
 Schwarz, A. J. F. (1964) *Ann. paediat. (Basel)*, **202**, 241
 Sever, J. L. (1962) *J. Immunol.*, **88**, 320
 Takátsy, G. (1955) *Acta microbiol. Acad. Sci. hung.*, **3**, 191
 United States Public Health Service (1963) *Biological products*, Washington, D.C., US Government Printing Office (Publication No. 437)
 WHO Scientific Group on Measles Vaccine Studies (1963) *Wld Hlth Org. techn. Rep. Ser.*, **263**