shoulder and hip joints; in the former, surrounding the glenoid cavity; in the latter, the acetabulum, with the exception of the cotyloid notch.

The third class of fibro-cartilages are those placed between the bones of the vertebrae, acting as elastic cushions, in preventing concussion, and thus preserving the delicate spinal marrow from those shocks which must of necessity be continually occurring were it not for this contrivance. The arrangement of their structure is different from that of the other fibro-cartilages; they are thickest in the middle. The fibrous tissue is disposed in concentric layers, each layer being formed of interlacing fibres, having their interstices filled with cartilaginous substance; towards the centre the fibrous structure gradually disappears, leaving a soft pulpy mass, adapting itself to every motion of the vertebral column. These fibro-cartilages are firmly attached to the bones immediately above and below them, and also to the connecting ligamentous fibres; and although from this arrangement but very little, if any, motion is admitted between any two of the bones, still there results in the aggregate an elastic flexible column, in every way fitted to the very important duties it has to discharge in the animal economy. These pads or cushions are not only very elastic, but they possess a remarkable power of resistance to pressure, as you may see in this preparation which I now show you.

The patient from whom it was taken was the porter of this hospital, and had suffered for several years from aneurism of the descending aorta, which ultimately destroyed him. You may perceive the bones have been absorbed to a considerable extent, but the intervertebral substances remain entire, having resisted that pressure which destroyed the harder and less yielding tissue.

In our next lecture we will resume the consideration of the tissues entering into the formation of joints.

CASE OF EXCISION OF THE GREATER PORTION OF THE ULNA.

By WILLIAM MC. EWEN, Esq., House Surgeon to the Chester Infirmary, and Surgeon to Chester Castle, &c.

(Read before the Provincial Medical and Surgical Association, August 8, 1844.)

Thomas Griffiths, aged 19, apprentice to a marble mason, states that in the beginning of February, 1843, three of the men were lowering a large block of stone, and he was assisting them with a crow-bar. The stone being too heavy, came down suddenly, and knocked him down, his right arm being under the crow-bar, from which he could not get released until the stone was removed, which was done as speedily as possible.

He was able however to continue his employment for a fortnight after, though complaining much of his arm, which was very painful and a little swelled. He applied to a surgeon, who ordered leeches to his arm, and afterwards cold applications, with rest. This treatment was followed up for a week. The leech bites became a little inflamed, but after soothing applications got better, the arm being much relieved. His eyes then became inflamed, and the attention of

his medical attendant was chiefly directed to them, the arm being allowed to rest.

In May, his eyes gradually got better, but his arm again became very painful, red, and swelled. Blistering and other topical applications were used until August, when he was recommended to try sea bathing; after a time however, he returned home, without having received any benefit.

Shortly after Christmas an abscess on the outer side of the arm burst, and discharged freely, poultices and fomentations being applied daily.

In the beginning of March, 1844, he applied to me to know whether anything could be done; his general health was giving way rapidly, as evinced by night perspirations, loss of appetite, and muscular power, under the irritation and discharge from the arm, which presented the following appearance:—

About an inch above the right wrist there was a large sloughing ulcer, extending half way round the arm, exposing the shaft of the ulna to the extent of rather more than an inch, the exposed portion of bone having a whitish, worm-eaten appearance. The skin and cellular tissue for a considerable space around the ulcer were of a dull livid colour, and the muscles so matted together as to prevent motion of the wrist and hand, which was half closed. The margin of the ulcer was ragged, and had an irritable and angry appearance, and was extremely sensitive and painful. The ulcerative process was rapidly extending itself round the arm. Several sinuses communicated with the ulcer: in one the probe could be passed along the bone near to the olecranon; another passed down close to the articulation of the wrist.

The discharge from the ulcer was extremely offensive, and of a thick whitish consistence.

Before adopting any operative procedure, the system required to be put in order. The digestive organs were carefully attended to, and quinine administered with sulphuric acid, which improved his appetite and reduced the night perspirations; still it was evident the system could only be temporarily benefitted so long as the source of irritation remained.

After mature deliberation I proposed to remove the diseased portion of bone, which he willingly submitted to. Before doing so, I had written to my friend, Professor Fergusson, of King's College, as to the propriety of removing the bone at its articulation at the wrist, if it should be found that the disease had not extended so low. He, in reply, advised me to remove the articulating surface, as there might be a possibility, if that small portion only was left, that it might die for want of nutrition.

Coinciding in his opinion, on the 15th of March I proceeded in the following manner:—

The patient being seated in an arm chair, an assistant held the arm semiflexed, and with a common scalpel I carried an incision from the upper part of the ulcer along the outer edge of the ulna, to within three inches of the point of the olecranon. The bone being my guide, I dissected away the attachments of the muscles first along the outer, and then along the inner edge of the bone, making in fact two flaps.

The diseased portion of bone being now exposed to the extent of three inches, from the upper end of the ulcer, the periosteum was here found very much

thickened, and peeled easily off the bone. I therefore carried my incision a little higher up, until it appeared perfectly healthy. The next step was to divide the bone. This I attempted to do with a pair of Liston's bone forceps, having first made a notch in the bone with a small saw; unfortunately however in my attempt, one of the blades of the forceps gave way. I had immediate recourse to one of Hey's crown saws, which was at hand, an assistant holding down the flaps; the bone was with some difficulty sawn through. The next step was to disarticulate the bone at the wrist; this there was little difficulty in doing. An incision was carried down from the lower end of the ulcer in a direct line with that made up the arm to the head of the metacarpal bone of the little finger, the tendons of the flexor and extensor muscles being carefully separated from their attachments on the anterior and posterior aspect. The knife was then thrust perpendicularly betwixt the radius and ulna, and carried round the articulating surface, completely dividing the sacciform ligament, and the attachments of the triangular fibro-cartilage. There was now little left but the attachments of the interosseous ligament.

A copper spatula having been passed under the disarticulated end of the bone, by gently raising it there was more freedom allowed for carefully dissecting the piece out, which was done without wounding any large artery; one small vessel only was divided, a branch from the interosseal, but the hæmorrhage from it was so trifling as not to require ligature. Several portions of very thickened periosteum were dissected out of the bottom of the wound.

The edges of the incision at the wrist were brought together with adhesive plaster. As there was reason to suppose from the tolerably healthy state of parts that union would take place by the first intention, the upper portion of the wound was filled up with dry lint. The patient was then put to bed with his arm pronated, and resting upon a high pillow, and cold water dressing applied. He bore the operation very well; little more than three ounces of blood were lost.

8 p.m. on the day of operation. Has been tolerably easy, and slept at intervals in the afternoon, rather thirsty, pulse 80. To have fifty drops of laudanum immediately.

March 16th, 10 a.m. Has slept between three and four hours in the early part of the night, and has been dosing this morning. Does not complain of pain in the arm; pulse 90, regular, rather full; face a little flushed; tongue moist, and a little furred in the centre, which has been so for some months. No thirst; bowels not open. To have tea and bread and butter. Arrowroot if he wants it. Continue wet cloths.

8 p.m. Has been easy during the day; no great pain in the arm, no swelling; pulse 100; slept none; bowels not open; is cheerful, and in good spirits. To have fifty drops of tincture of opium in the night if he does not sleep, and cold spirit lotion applied.

17th, 10 p.m. Last night about half-past ten, he complained of griping pain in his bowels; the draught was administered, which procured him immediate relief, and a good night's rest, having slept six hours. Pulse 100, but regular and sharp. Heat of skin natural; no great thirst; bowels not open; does not complain of much pain in the arm; no swelling;

slight discharge of matter from the edges of the wound. To have half an ounce of castor oil immediately.

8 p.m. Has had his bowels opened twice with the castor oil; arm easy; has slept two hours in the afternoon; was a little chilly when he awoke. Pulse 102, and sharp. To have forty-five drops of tincture of opium in the night, if he does not sleep.

18th. He took the draught in the night, and slept nearly eight hours. Perspired a little in the morning. Pulse 100. The dressings were for the first time removed; a considerable quantity of pus followed; wound looks healthy; no hæmorrhage. Dressed with dry lint as before, brought round with a few strips of plaster.

8 p.m. Has slept two hours in the afternoon; arm easy. To have fifty drops of tincture of opium at bed time.

March 19th. Has had a good night, and feels very comfortable. No swelling or uneasiness in the arm. Pulse 100. Perspired freely in the night; dressing removed; no hæmorrhage; wound looks healthy, and granulations springing up around the edge of the old ulcer. Dressed with dry lint.

20th. Has not had a good night, having been without his draught; bowels open; dressing removed; free discharge of pus of a healthy character; here and there slight union had taken place betwixt the lips of the wound, these were broken up in order to allow of the free escape of matter. To be dressed with dry lint as before, and covered over with soap plaster.

21st. Going on favourably.

23rd. Has had two good nights without any draughts. Bowels open; pulse 88; no perspirations; tongue clean and moist; no tenderness or swelling of the arm. The granulations are healthy, and the discharge moderate. The incision at the wrist nearly filled up. To be strapped as before. Got up to day to have his bed changed.

25th. Sleeps well; no perspirations; wound looks very healthy; no uneasiness in the arm. To have meat diet, and a pint of porter daily.

26th. Sleeps well; bowels open; tongue clean and moist; no perspirations; wound at the wrist nearly healed. There is a little discharge from the end of the bone, which appears more prominent than usual, but has plenty of covering. The arm was freely extended, which gave him no pain.

27th. Only feels a little stiff. The arm can be put in a state of pronation or supination without any uneasiness. The arm to be placed in the same semiflex position, but a little more extended. Continue his meat diet, and porter.

28th. Pulse 84; slept well; has had some pain along the front part of the arm, from the altered position and movement yesterday. To be strapped as before, but without any lint, and the arm to be more flexed.

April 1. Wound looks healthy; complains of pain at the end of the bone. Got up to-day for several hours, sleeps well, and takes his food well.

4th. Bottom of the wound filling up, and very much contracted. The edges of the old ulcer appear irritable and painful. To have them touched with the nitrate of silver; can pronate and supinate the hand

himself. The fingers are rather stiff. The incision at the wrist perfectly healed. Health very good.

7th. Wound continues improving; the edges of the old ulcer still painful, but look more healthy. Nitrate of silver applied to them. Has perspired a little the last two nights. Complains a little of sore throat; is a little feverish: bowels open. To steam his throat with hot water frequently.

9th. Sore throat better; the edges of the ulcer appear irritable; the discharge is unhealthy, and the ulceration is extending over the skin in the under surface of the arm. To have it fomented frequently with warm water, and linseed poultice applied for a few days.

12th. Since the poultice has been applied, the wound has discharged freely. The edges of the ulcer look clean and more healthy. Poultice to be discontinued, and the wound strapped as before.

14th. Ulcer healing rapidly. Surface very much contracted, and new skin forming. Says he feels in better health.

20th. Has been walking out daily since last report, and came down to the surgery to have it dressed with plaster. The granulations are now nearly on a level with the edges of the wound, which has to day an irritable and erysipelatous appearance, brought on, I imagine, from some injudicious friend giving him wine. To have a poultice applied, and to take some aperient mixture.

From this period the edges and parts around the old ulcer alternately progressed and retrograded. During the whole progress of the case the pus secreted from the ulcer was very similar to what is seen in opening a scrofulous abscess of the neck.

Until the beginning of June, a variety of topical applications were used without much benefit. Amongst these I may mention solutions of sulphate of zinc, sulphate of copper, black wash, and the nitrate of silver in pencil, with a view of altering the character of the sore, attention at the same time being directed to the digestive functions, with generous diet; strapping with soap plaister, and bandaging the arm, were found on the whole most efficacious.

The incisions made with the knife being at this time healed, he was sent to the sea side, and ordered to bathe the arm in salt water. There he remained five weeks, and returned with the sore quite healed, and in perfect health.

REMARKS.

I think it is pretty evident from the early history of the case, that the bone had received serious injury, and although he was able to follow his employment for some time after the accident, no doubt there was a considerable degree of inflammation of the bone and periosteum going on. We are well aware that bone generally speaking is very slow in taking on diseased action, but there was here, besides the injury, a scrofulous diathesis to combat with, which merely required some external agency to produce its development. We see too, how in such a constitution diseased action may suspend itself, and attack a more distant organ, and after a time return to the seat of the original mischief.

With regard to the operation, I may remark that the difficulty attending the division of the bone was, inde-

pendent of its enlargement, much increased, owing to the blade of the forceps giving way, and though it would have materially shortened the operation, I much question if I could have accomplished my object with them. I found Hey's saws answer every purpose, though tedious in their operation; the only danger likely to accrue from their use, is when having nearly divided the whole thickness of bone, they might, if caution were not used, suddenly pass through and divide the ulnar artery which is lying close to it. It is better therefore to finish the division with the forceps.

Velpeau had a case in which he was anxious to remove the radius for disease entirely restricted to that bone, but the patient would not submit to the operation, and had her arm amputated. In his book, (*Nouv. Elem. de Med. Oper.*, Tome 1,) he recommends a chain saw to be placed under the bone, by means of a director, and the bone to be sawed through, and if there was not room for its use by the longitudinal incision, that a transverse one should be made. I have never used the chain saw, nor seen it used, though much has been written about its applicability in such cases, but I should much question the propriety of its use in any case.

The removal of portions of the shafts and extremities of bones extending into joints, is now a matter of common occurrence, but the resection of the whole thickness of any of the long bones for necrosis is extremely rare. I am not aware that any case has presented itself, in this country at least, nor can I find any case on record in which so much bone has been removed.

The disease, as you will perceive in the specimen before you, is extensive,* and extends close to the articular extremity, and though the heads of long bones are slow in taking on diseased action, it would in this instance soon have been affected, and though the head of the bone is not anatomically connected with the wrist joint, there is every reason to fear that sooner or later the limb would have required amputation.

You will be able to judge of the present appearance of the limb by this cast. The arm is small certainly, but the muscles, although wasted very much, have a very considerable degree of power, indeed the youth has been daily exercising it in rowing about on the river: he can pronate and supinate the hand, can flex and extend the fingers readily, and has free motion at the wrist.

I may mention that in operating upon a case of a similar kind, I would strongly recommend that after the excision of bone, the hard and irregular edges of the ulcer should be carefully pared: it would contribute materially in hastening a cure, and I am convinced had this been done in the case just recorded, that a cure would have been sooner effected.

* The preparation and a cast of the limb taken after the restoration to health were exhibited.