resistance to the closure of these tubes; for they appear to follow closely upon each other even when their interval is magnified to the utmost extent. The direction of their current is determined by the known forces which maintain the circulation; but those forces are aided materially by an agency which tends to adapt the diameter of the capillaries to that of the globules of blood; to diminish it, when the circulation is arrested, in a direct ratio to the quantity of globules left; and, thus, to render the force of the heart, which is in an inverse ratio to the diameter of the vessels, available to the utmost extent during returning circulation.

If a drop of blood be placed on glass at the margin of a thin plate of talc, it will be attracted towards, and, as is well known, will creep into the capillary space. When the spontaneous arrangement of the blood, thus disposed of, is observed with a microscope, it will be noticed that the globules do not rest upon the same level, but one upon another. By pressing upon the talc with a needle, and changing the focus, it will be found that the capillary space may be cleared of the globules around the spot where the talc is compressed; that by relaxing the pressure gently, a single layer of globules will rush into view, and by removing it entirely, a wave of blood, two or three globules deep, will roll over the first layer.

When blood corpuscles are thus compressed, they are more flattened than natural, but resume their former shape when set free. The exercise of this elastic property of the globules is the cause of the ascent of the talc in the preceding experiment, and of the return of these bodies to their former arrangement. So far the globules of the blood evince sufficient elasticity to resist the attraction which opposite surfaces exert.

Two surfaces made to adhere by means of a saline fluid, are so closely in contact as scarcely to admit blood-globules. Talc and glass thus made to adhere, and a drop of blood placed at the margin of the talc, the blood-globules penetrate in small numbers, comparatively, and only in a single layer, of which the globules are far apart, and have only advanced a line or two; in fact the globules do not penetrate with the same facility as when the interval of the glass and talc is dry. When capillaries are once empty, the apposition of their walls excludes the blood-globules to a great extent, much in the same manner as above.

A few words may be added on the lymphatic circulation. The presence of valves in a vessel implies the existence of some opposing force, calculated to impede the movement of its contents. The valves of the veins afford an instance of this, and are so formed as to neutralize the effect of gravity, and render the contractions of the muscles available, as an auxiliary force, to impel the venous blood in the direction of the heart only; a force which would otherwise distribute it equally in that and the opposite direction. But the lymphatics are capillary tubes, the force of gravity therefore can exercise no influence upon their contents. The valves which they contain must consequently have a different use to the valves of the veins. It has been remarked above that the heart determines the course of the current, but that the pressure of the sides of capillaries, by adapting the diameter of the vessels to their contents, increases its rapidity. This reasoning applies equally to the movement of the lymph, except

that in the former instance the heart directs the current by impulsion, in the latter by suction.

The effect of this capillary force, exerted by the delicate surface of the lymphatics upon itself, without the interposition of valves, would be to create an oscillating movement throughout the length of a vessel, whenever the volume of its contents was diminished by the suction of the heart; whereas, the presence of valves, by putting a check to retrograde movement, causes the force of the pressure to take effect only in the direction of the heart.

The preceding remarks will assist the reader to acquire a clearer understanding of the physiology of congestion, as relates to the lessened diameter of capillary tubes which succeeds to arrest of circulation.

Congestion as above defined (VII.), might be more correctly distinguished by the name of sub-inflammation, as it is the invariable antecedent of the true inflammatory process. Such a distinction would serve to separate it from other forms of capillary disorder which are likewise called congestion, and all of which acknowledge a different pathogeny.

Recent observers have corrected the ideas, but left the language of science in an unsettled state; if they have introduced fresh terms, the new have but inadequately responded to the wants of knowledge. The difficulty of defining a state of disease by means of a single term, is insurmountable: a full sentence, even is unequal to the purpose. Hence the necessity of retaining words, which by usage, have attached themselves in so comprehensive a manner to states of disease, as really to mean the states themselves in their most ample and recent developments. These remarks apply particularly to inflammation, the philological value of which term, in no wise concerns pathology.

The term sub-inflammation, applied instead of congestion, to the process (VII.) which immediately precedes inflammation, would be free from objection, inasmuch as it embraces all that it is intended to signify; while *erubescence*, or *vital turgescence*, is sufficiently expressive to denote the class of facts which comes under definition VI. Even this normal phenomenon is sometimes called congestion, as in speaking of the turgescence of erectile tissue and analogous facts.

VIII.—*When irritation subsides, the velocity of the circulation is restored to its natural state; exhalation is re-established, the usual diameter of the capillaries is regained, and the globules recover their elasticity.*

The consequences of this form of congestion (sub-inflammation) may thus undergo a process of cure by resolution, which consists in the separate steps of functional disturbance being retraced, as defined above; or may advance into inflammation.

(To be continued.)

REPORT OF MIDWIFERY CASES OCCURRING IN PRIVATE PRACTICE.

By KENRICK WATSON, Esq., F.R.C.S., Stourport.

TO THE EDITOR OF THE PROVINCIAL MEDICAL AND SURGICAL JOURNAL.

SIR,

At the present crisis it appears to me to be very desirable that the attention of the profession and of the public should be directed to the necessity of all

persons, both male and female, who engage in the practice of midwifery, being more or less educated. This, perhaps, would be done most effectually, by practitioners in different parts of the kingdom publishing the result of their practice, with the view of showing the proportion of cases in which the life of both the mother and the child depended upon immediate assistance. With this intention I beg to offer for publication the result of 800 cases of midwifery, of which I have kept an account. With reference to the unusual number of dangerous cases, I would merely observe that they constituted a large proportion of the difficult cases occurring in the neighbourhood, many of them residing several miles from my house.

I am, Sir, yours very truly,

KENRICK WATSON.

Stourport, Nov. 5, 1844.

The following is the result of 800 cases of midwifery of which I kept an account during a period of eighteen years:—

Premature labours	12
Twin cases	18
Face presentations	14
Breech ditto	11
Presentations of arm, &c.	11
Dangerous flooding	17
Convulsions during labour	3
Ditto after birth of the child	1
Rupture of the vagina	3
Ditto of an artery in the labia pudendi	1
Cases where it was necessary to open the head	12
Severe flooding after extraction of the placenta	2
	105
Natural Labours	695
	800

Of the premature births several were occasioned by one or both parents being affected with syphilis.

In two cases in which the arm presented in the vagina, the placenta had been over the os uteri, but had been forced into the vagina by the protrusion of the arm. The flooding in the first instance, in both cases, had been violent, but had ceased when the placenta was forced into the vagina. The women in both cases were strong and healthy, and the uterus strongly contracted round the body of the child.

Of the seventeen cases of severe flooding, fifteen arose from the adhesion of the placenta over the os uteri; and in two cases the placenta was not so situated, and the flooding ceased after the membranes were ruptured. Of the fifteen cases one died before I got to the house, and one soon after she was delivered. She had been flooding two or three days before I saw her. In these seventeen cases four children were saved.

In one of the cases in which the head was opened the woman was dropsical. She had been in labour many hours; the uterus was so thin that the limbs of the child could be distinguished by the touch through the parietes of the abdomen. The uterus was dilated, the head was large, and the bones detached. After the head was punctured, about a quart of fluid was discharged, as the bones were so loose that they afforded no hold to the crotchet. I passed my hand by its side, and brought down the feet. No

contraction of the uterus followed; the re-introduction of my hand produced no effect, the uterus continued like an empty bag, with the placenta attached to its fundus. I gave her brandy and ammonia, poured cold water on the abdomen, and then injected cold water into the uterus without the least effect. I then applied a thick compress upon the abdomen, and a flannel roller round the body. There was no discharge of blood at the termination of two hours. I left her in the charge of her medical attendant, and in the course of the night she died.

The case in which a rupture of the labial artery took place, was attended by a midwife. Two hours after delivery, she complained of pain and swelling of the right labium, bearing-down, and discharge of blood from the rectum. She died on the third day. Upon examination, thirty hours after death, the perineum and surrounding parts were found injected with blood; the integuments of the abdomen emphysematous, a cavity in the labium lined with coagulated blood, and communicating with the rectum, through a laceration about half an inch in extent.

In two cases the placenta was diseased. In one, the maternal part was left firmly adhering; great irritative fever supervened, with very offensive discharges, but by injections into the uterus, bark and acids, the woman, who was nearly fifty, recovered. In the other, a discharge of water took place in the fourth month, and continued to recur from time to time till the seventh month, when labour commenced. At this time the uterine tumour was divided into two parts. When the os uteri was sufficiently dilated, I passed my hand and found the foetus, with the membranes entire, occupying one half of the uterus, and the placenta and a quantity of grumous blood, the other half. So far as I could ascertain, the putrid mass was formed by blood and hydatids. The woman recovered.

In one case there was a knot on the umbilical cord, and in another, a rupture of the umbilical artery. In a third case the child was apparently seized with convulsions before labour commenced, and died. The woman when at work, ironing, in a very hot room, became faint from the violent motion of the child; it shortly became still, and never moved again. She was delivered on the fourth day afterwards, and the child was in a state of decomposition.

Several cases required the use of the lever or the forceps, and in no case, unless the death of the child was evident, was the head opened till one or other of them had been tried.

Out of the 800 cases seven proved fatal,—three from rupture of the vagina; two from flooding; one from rupture of an artery in the labia pudendi; and one, in which the head was opened and the child putrid, where peritonitis had either preceded, or occurred during, labour.

EFFECTS OF MIASMA.

TO THE EDITOR OF THE PROVINCIAL MEDICAL AND SURGICAL JOURNAL.

SIR,

As the etiology of disease has always excited the attention of the medical practitioner, perhaps you will deem the following contribution, abstracted from my