PROVINCIAL MEDICAL & SURGICAL JOURNAL.

OBSERVATIONS ON AGUE.

By RICHARD CHAMBERS, M.D., Physician to the Essex and Colchester Hospital.

John Copsey, aged 18, a sailor, applied as an out-patient of the Essex and Colchester Hospital, November 20th, 1845. He stated that he had been fifteen months ill with ague, which at first was a tertian, but latterly has assumed the quartan type. He was under the care of three different medical men, and had also applied to a lady in his neighbourhood; but from none of them did he derive the slightest benefit. I could not learn any particulars of the treatment that had been adopted, except what might be concluded from a message that he brought me, to say that he could not bear the use of quinine. At present he feels weak and exhausted, and looks anæmic; pulse 60; bowels confined, motions dark; urine scanty; tongue furred, at the base; and he has a bitter taste in his mouth. He complains of heaviness in the hypochondria; the feet and ankles swell towards night, and there is a degree of puffiness of face,—in a word there is congestion of all the viscera. The action of the heart is regular, but extremely laboured, it appears as if it is overloaded.

Fiat venesection ad oz. x.

R. Quinæ Sulph., gr. x.; Acid. Sulph. Dil., m. ij.; Aquæ Menthæ Pip., oz. j. M. Fiat haustus quam primum sumendus.

R. Extract. Colocynth. Comp., gr. v.; Extract. Hyoscyami, Hydrargyri Chloridi, utrq., gr. iiss. M. Fiant pilulæ duæ tertiis noctibus sumendæ.

R. Quinæ Sulph., gr. ij.; Acid Sulph. Dil., m. viij.; Tinct. Cardam. Comp., dr. ss.; Aquæ Menthæ Pip., oz. j. M. Fiat haustus ter quotidie sumendus.

December 1st. He has not had a paroxysm since the 19th, that is, the day previous to his admission on the hospital books. He feels much better; pulse 72; tongue clean; bowels regular.

R. Extract. Colocynth. Comp., gr. iij.; Pil. Hydrarg., gr. ½; Pil. Scillæ Co., gr. iss. M. Fiat pilula omni nocte sumenda.

Cont. Haustus.

He continued this medicine for a fortnight, and having remained free from the ague, he discontinued further attendance, and up to this moment has been perfectly well.

Although the treatment of ague by quinine partakes in some measure of empiricism, we shall find, on examination, that it is based upon correct principles; and that in those cases which prove refractory to this parti-

cular treatment, the blame is attributable, not to the remedy, but to our inattention to those principles which we recognize, (and which are deemed so essential,) in the management of other diseases. It is true that quinine will alone, in the larger proportion of cases, be adequate to the removal of the disease; but in cases of long standing, we meet with congestions of the internal organs, which interfere with, or even altogether forbid its employment; and if these congestions are not removed, we will be unable to cure the disease, although we may arrest it temporarily.

In a former communication on this disease, which is inserted in No. 44 of the last volume of the Journal, I stated that I had experienced unexceptionable success from the use of ten-grain doses of quinine, followed up by smaller doses; and that I had not met with a single case in which, through idiosyncrasy, or other causes, I had been prevented giving the quinine. As the case which I have just detailed could not bear the quinine before he came under my care, I resolved on giving the matter a fair trial. But first let me again mention an outline of the case:—

There was congestion, and consequent inactivity, of all the excretory and secretory organs; the blood was increased in quantity, but deteriorated in quality; there existed that state which Dr. Turnbull, in his lectures in the *Lancet*, alludes to as anæmic plethora. There was mental and physical depression, in consequence of the long duration of the disease, and its resistance to treatment. It was evident, then, that relief to the congestion was of primary importance; yet his feeble pulse, and his weak and pallid appearance, almost forbade having recourse to depletion; however, all doubt as to its propriety was quickly removed on referring to the heart, from the laboured action of which it was evident that the feebleness of the pulse was merely the feebleness of oppression. Accordingly he was bled to ten ounces, and took ten grains of quinine before he left the hospital, and in addition I directed him to take a glass of gin-and-water, as he had a long journey to go, and the day was cold. The quinine was continued for three weeks, and during the whole period it did not produce the least unpleasant effect. It is, then, I think, apparent from the result, that the obstinacy of the disease, and the disagreement of the quinine, arose from non-attention to the co-existent visceral congestion.

But to me the case appears to derive its chief interest as an illustration of a rule of practice applicable to this and several other diseases, and one not sufficiently attended to. As an example, I will take neuralgia. It must be familiar to us all to see a case that has been

aggravated by iron in the hands of one practitioner, cured by the same remedy in the hands of another, with this difference, that the latter may have prescribed a few blue pills and a saline aperient; nay, more, a remedy may be borne to a certain point, congestion may then arise, the remedy disagrees, and without any effort to remove the congestion, the remedy is condemned, and of course discontinued. Quinine and iron afford daily illustrations of this fact.

In the last number of the *Dublin Journal*, there is a paper by Dr. Graves, on the relapsed period of ague. The paper is one of great interest, and will be read with that attention which everything emanating from one who has so earnestly and so successfully devoted himself to the advancement of clinical medicine is entitled to. It does, however, appear to me, that the practical deductions he has drawn are not warranted by the facts of the case he details. A brief outline of the case will be necessary before passing it in review; my space is necessarily too limited to transfer the whole, I beg, therefore, to refer my readers to the original for full details.

"The case I am about to detail, possesses likewise several features of practical interest, and serves to show that a very obstinate species of ague may be cured by the use of quinine alone, and that very large quantities of that powerful medicine may, under such circumstances, be taken, not only with impunity, but with advantage. A boy, of good constitution, and eleven years of age, had been at a boarding school in Kent during the spring and summer of 1842, and remained in perfect health all that time. In autumn he was very imprudently allowed to bathe daily in a pond of stagnant water, and he frequently continued in the water for more than an hour. In the November following, feverish symptoms exhibited themselves, and he was several times an inmate of the school Infirmary; his disease was considered to a frequent return of feverish attacks, from cold and indigestion; and accordingly he was treated by confinement and low diet, with mercurial and saline purgatives. Notwithstanding those remedies the disease frequently recurred, nor was its true nature ever suspected by the medical attendant. He arrived in Dublin on the evening of the 16th December, and the moment I saw him I concluded from his complexion that he had ague."

On the next day the ague shewed itself, and continued to recur, (with the exception of a few intervals,) for a period of two years, during which time he took the enormous quantity of 1680 grains of quinine, in doses varying from five to ten grains.

"With respect to the manner in which quinine was used, the following observations may be made:—At first I gave it in the usual manner, until the particular series of fits ceased, and persisted in its use for ten days or a fortnight, gradually decreasing the quantity taken. This is the method generally recommended by authors, and it is founded on the notion that it is necessary where the medicine is given in large doses, not to omit its use abruptly, lest the system should feel the loss of the powerful tonic. My experience in this and other cases leads me to doubt the accuracy of the reasoning upon which this treatment is founded, and I am convinced, that in following this rule, we are defeating our own object, by accustoming the constitution to the medicinal effects of the quinine, at a time

when the ague fit is absent. The quinine is the proper antiperiodic of the fit, and while the fits require the medicine, it is borne well by the constitution. On the contrary, when the fits are absent, its curative effects appear to be diminished, and the constitution becomes so accustomed to it, that when the disease again requires it, the medicine no longer exerts its anti-ague influence. We have an analogous example in the case of mercury, of which moderate quantities, judiciously exhibited, are sufficient to cure the venereal disease, provided the mercury is given when venereal symptoms are present, and only in the quantity necessary to control these symptoms. If it be given by way of prevention, when these symptoms are not present, or in too great quantity when they are, the system in either case becomes saturated with the mineral, but it is not protected from the further ravages of the venereal disease. The second mode of treatment which I adopted was calculated to avoid the inconvenience already pointed out. This method consisted of giving quinine for four successive days, and intermitting it for the six following days, thus embracing the interval comprehended in three fits. By these means it was hoped to keep the system sufficiently under the curative influence of quinine, while we avoided rendering the constitution too familiar with the medicine, the six-day interval preventing it from being saturated by the quinine. This method of treatment seems to have been eminently successful, and under its influence the disease abated in violence, the frequency of attacks decreased, and the long interval of 136 days was at last obtained. Finding however, that though it had broken the violence of the disease it had not extinguished it, I tried another on the third plan, which was to give no quinine until a well-marked fit or shadow of a fit occurred, and then at once to use the medicine in large doses, so as to stop the fits as soon as possible. The moment this object was accomplished the medicine was omitted, and was not again given until the paroxysms recurred, when they were similarly treated. This, on the whole, appears the best method, as it stops the paroxysms speedily, and keeps the medicine in reserve till they re-appear."

Here, then, was a case which lasted for two years, and did not yield till 1680 grains of quinine had been taken; and yet it is wished, from the result of this case, to lay down practical rules for the general treatment of this "generally" manageable disease.

Is quinine a specific for ague? It is: but only to the extent of being the most powerful vegetable tonic that we possess. Arsenic is just as powerful an antiperiodic, but from its tendency to injure the constitution, it is very properly less resorted to. Iron, zinc, gentian, and several other remedies, denominated tonics, have been used effectually in the cure of ague, and even change of air by reason of its tonic influence has done it. Our object in using quinine in ague is, to obtain its tonic influence on the vascular system. One large dose may effect this, but in consequence of the distended state of the capillaries, we find it necessary to follow it up with repeated small doses, not as Dr. Graves supposes, lest the system should feel its abrupt discontinuance, but to keep up a stimulus to the capillaries till they had recovered their tone. It is a fallacy to suppose, with Dr. Graves, that we ought to discon-

tinue the quinine for fear of weakening its effects, if subsequently needed; for I contend that, if proper precautions are observed in the administration of the medicine, its continuance is the only sure and expeditious mode of overcoming the disease. He resorts to the analogy of mercury, in the cure of the venereal, to support his views; but it strikes me that a closer analogy exists between the administration of wine in the debility of fever, or the exhaustion of hæmorrhage; and let me ask, who would think of discontinuing the wine on the first appearance of reaction in either instance, lest, by continuing its use, it would lose its influence if again required; would they not rather continue it in diminished doses, or at longer intervals, till, by its influence, aided by the necessary auxiliary treatment, the probability of relapse had passed away.

Piorry states that the paroxysm of ague forms but the second link in the concatenation, the first being splenic engorgement; and unquestionably there exists a very close connection between ague and affections of the spleen. And where a person who has resided for a time in a malarious district, removes to a healthy, and yet becomes attacked with ague, is it not more than probable that it is by an impression made upon the spleen at first, that the disease is subsequently manifested, if aided by depressing causes. We know, from observation, that the blood of those who have imbibed the malarious poison, becomes deteriorated in quality, and is prone to accumulate, (probably in consequence of a diminution of its ordinary stimulant quality,) in the internal viscera, more especially the spleen. We also know that this splenic engorgement is best treated by tonics; nay, that it is aggravated by the ordinary antiphlogistic treatment.

The error that appears to me, on the perusal of the case under consideration, is the too exclusive reliance that was placed on quinine, no efforts having been made to remove the necessarily attendant visceral congestion. It is even only incidentally mentioned that the spleen was engorged; or, as the report states, "evident tumefaction of the spleen had commenced." *Quere?*

It is important to remark, that after the disease had been absent at one period for three weeks; it recurred again, in consequence of an exposure to a cold breeze, while on a boating excursion. Now, under ordinary circumstances, we would expect to find internal congestion from exposure to cold, and is it not right to infer that the relapse in this instance was caused by internal congestion? It is not pretended that any malaria was in operation. We can readily perceive that the severity of this case arose from the oft-repeated and peculiarly powerful manner in which the malarious poison was applied, namely, by frequent and long immersions in stagnant water. As to the relapsed periods of ague that were observed in this case, I think that we ought more properly to view the intervals as the masked periods, the disease subsequently showing itself when the influence of the quinine had passed off.

In the case that I have myself detailed, quinine could not be borne before it came under my care, in consequence of the neglect in not removing the great congestion that existed, and in the case that I have quoted, I think I am justified in stating that like causes produced analogous results.

TABLE OF DISEASES TREATED BETWEEN THE 24TH OF FEBRUARY, 1843, AND THE 31ST OF JANUARY, 1846.

By GEORGE FIFE, M.D., Physician to the Sunderland and Bishop-Wearmouth Infirmary.

(Continued from page 120.)

In my last communication my remarks on the diseases, contained in the table, extended to rheumatism and its complications; in the present, it is my intention to continue the same plan of proceeding, i.e., taking the diseases according to their numerical importance; and when this has been done, to enter upon the more particular consideration of such as appear to possess most practical interest.

The disease which now calls for notice is phthisis, which in the table may be regarded as rather a small number, forming not more than a twenty-third and a fraction of the gross number of cases treated. That the bills of mortality of the district would, if looked into, afford a different average, I have not the slightest doubt, nor would such result constitute a real discrepancy. To explain this, it is again to be remembered, that the patients visited me at my own house, and that, consequently, most of the cases came under my notice at a comparatively early stage; besides which I was most scrupulous in setting down any as phthisis, which did not bear the test of physical examination, as well as the assurance of general symptoms. It is also more than probable, that not having inserted any cases as phthisis where any doubt existed, many may have eventuated in the disease, which were differently regarded by me. This uncertainty could not, of course, be avoided, where many of the cases were lost sight of, after a few visits. Be this as it may, I would not regard phthisis as being, by any means, as prevalent in this town and its immediate neighbourhood as it is in other localities. In this opinion I derive support, both from my private practice and from my books at the Infirmary. From minute attention to the history of many of the cases, and from careful observation of the course which they ran, I feel fully justified in the opinion that inadequate nutriment is a most fertile source of tubercular disease, and that it is so, even where no decided family predisposition exists, so far as it was possible to ascertain the fact; and no cases, which have presented themselves to my notice, have run their course more rapidly to a fatal termination than those which have been most manifestly dependent for their origin on this cause. It being my intention to enter more fully on the investigation of this subject, I shall not extend the present remarks.

Herpetic eruptions and cutaneous affections may now be briefly disposed of. Most of these cases partook of the herpetic character, and evidently originated in the common constitutional causes. Many of them, however, occurred under circumstances which justified the belief in a syphilitic or mercurial origin. The effect of treatment also tended to the confirmation of such view. In several instances the most rapid improvement took place under the employment of the bichloride of mercury, where all other means, local and general,