Structure and Expression of Class II Defective Herpes Simplex Virus Genomes Encoding Infected Cell Polypeptide Number 8

HILLA LOCKER,^{1†} NIZA FRENKEL,^{1*} AND IAN HALLIBURTON²

Department of Biology, The University of Chicago, Chicago, Illinois 60637,¹ and Department of Microbiology, University of Leeds, Leeds, England²

Received 12 February 1982/Accepted 13 May 1982

Defective genomes present in serially passaged virus stocks derived from the *tsLB2* mutant of herpes simplex virus type 1 were found to consist of repeat units in which sequences from the U_L region, within map coordinates 0.356 and 0.429 of standard herpes simplex virus DNA, were covalently linked to sequences from the end of the S component. The major defective genome species consisted of repeat units which were 4.9×10^6 in molecular weight and contained a specific deletion within the U_L segment. These *tsLB2* defective genomes were stable through more than 35 sequential virus passages. The ratios of defective virus genomes to helper virus genomes present in different passages fluctuated in synchrony with the capacity of the passages to interfere with standard virus replication. Cells infected with passages enriched for defective genomes overproduced the infected cell polypeptide number 8, which had previously been mapped within the U_L sequences present in the *tsLB2* defective genomes. In contrast, the synthesis of most other infected cell polypeptides was delayed and reduced. The abundant synthesis of infected cell polypeptide number 8 followed the β regulatory pattern, as evident from kinetic studies and from experiments in which cycloheximide, canavanine, and phosphonoacetate were used. However, in contrast to many β (early) and γ (late) viral polypeptides, the synthesis of infected cell polypeptide number 8 was only minimally reduced when cells infected with serially passaged *tsLB2* were incubated at 39°C. The *tsLB2* mutation had previously been mapped within the domains of the gene encoding infected cell polypeptide number 4, the function of which was shown to be required for β and γ viral gene expression. It is thus possible that the *tsLB2* mutation affects the synthesis of only a subset of the β and γ viral polypeptides. An additional polypeptide, 74.5×10^3 in molecular weight, was abundantly produced in cells infected with a number of *tsLB2* passages. This polypeptide was most likely expressed from truncated gene templates within the most abundant, deleted repeats of *tsLB2* defective virus DNA.

The standard DNA genomes of herpes simplex virus types 1 and 2 (HSV-1 and HSV-2) are approximately 100×10^6 in molecular weight (MW) and consist of two covalently linked components, L and S (38, 42, 50). The L component consists of unique sequences U_L (MW, 67×10^6) bound by the inverted repeats *ab* and *b'a'* (MW, 5.8×10^6). The S component consists of unique sequences U_S (MW, 9×10^6) bound by the inverted repeats *ac* and *c'a'* (MW, 4.1×10^6). The sequence *a* is present in one or few copies at the L and S termini and at the L-S junctions (11, 19, 22, 26, 30, 38, 49, 51). The L and S components invert relative to each other, forming four

equimolar isomers which have been designated P, I_S , I_L , and I_{SL} (5, 14, 38, 55).

Whereas plaque-purified HSV stocks contain predominantly standard HSV genomes, virus preparations obtained through serial undiluted passaging have been shown to contain variable proportions of defective virus genomes in which the bulk of the parental DNA sequences have been deleted and substituted by tandemly arranged repeat units consisting of limited sets of the viral DNA sequences.

On the basis of the DNA sequences contained in their repeat units, the defective HSV genomes which have been characterized thus far fall into two distinct classes. The class I defective genomes derive their DNA sequences either exclusively from the S component of standard HSV

[†] Present address: Department of Genetics, The Hebrew University, Jerusalem, Israel.

DNA or from the L-S junctions (7, 9, 12, 21, 25). The most common species of this class contain repeat units ranging in MW from 5×10^6 to 6×10^6 and consisting of the entire *ac* inverted repeat sequences and a small set of the adjacent sequences from *U_S*. Because these DNA sequences have a high guanine-plus-cytosine content (5, 14), the defective HSV genomes of this type have a buoyant density which is higher than that of standard virus DNA. The high-density (HD) class I defective genomes were initially described by Bronson et al. (1) and have since been identified in series derived from several different HSV-1 and HSV-2 strains (for review, see reference 7).

The second class (class II) of defective HSV genomes derive the majority of their DNA sequences from the *U_L* region of standard HSV DNA and exhibit a buoyant density similar to that of standard virus DNA. Such defective HSV genomes were first recognized by Schröder et al. (40) in series derived from the ANG strain of HSV-1 (21, 22) and have since been found to occur in a number of additional HSV-1 series (for review, see reference 7). As shown previously, the class II defective virus genomes contain, in addition to the *U_L* sequences, sequences derived from the S terminus of standard HSV DNA (10, 21, 22). However, the exact organization of the *U_L* and S sequences within the repeat units of the defective genomes has not yet been fully documented. In the present communication, we describe our studies of the fine structure of class II defective HSV genomes which have evolved in the course of undiluted propagation of the temperature-sensitive mutant LB2 of the HSV-1 strain HFEM (13). In addition, we describe the pattern of viral gene expression in cells infected with *ts*LB2 virus stocks containing variable proportions of class II defective virus genomes.

(Portions of these studies were previously reviewed [7, 10].)

MATERIALS AND METHODS

Cells and viruses. Human epidermoid 2 (HEp-2) cells and African green monkey kidney (Vero) cells were obtained from the American Type Culture Collection. The derivation and properties of the *ts*LB2 mutant of HSV-1 (HFEM) were previously described (13, 15). The plaque-purified (PO) *ts*LB2 stock was derived by three sequential plaque purifications in Vero cells, followed by two sequential passages in HEp-2 cells at 10^{-4} PFU per cell and one additional passage at 10^{-2} PFU per cell. Passage 1 of the *ts*LB2 series was generated by infection of 2×10^8 HEp-2 cells with 1 PFU of PO virus per cell. Each of the subsequent passages was derived by infection of 2×10^8 cells with one-fourth of the virus stock constituting the preceding passage. The derivation and properties of the HSV-1 (Justin) series were previously described (8, 9, 25).

Preparation of viral DNA. Vero cells infected with 0.5 to 4 PFU of PO or serially passaged virus stocks per cell were labeled from 4 to 24 h postinfection with either [3 H]thymidine or 32 P_i as previously described (26). Briefly, the infected cells were treated with lysis solution containing proteinase K and sodium dodecyl sulfate (SDS), and viral DNA was prepared from the resultant cell lysates by equilibrium density centrifugation in CsCl gradients.

Restriction enzyme and hybridization analyses. Restriction enzymes were purchased from New England Biolabs and from Bethesda Research Laboratories, Inc. Enzyme digestions were carried out as specified by these companies. Electrophoresis of restriction enzyme digests for analytical purposes was done in 0.4 to 0.5% SeaKem ME agarose (Marine Colloids, FMC) gels. For the preparation of fragments to be redigested with additional restriction enzymes, electrophoresis was done in 0.5% low-melting-point agarose (Bethesda Research Laboratories), and the resultant gels were stained with ethidium bromide and visualized with UV light. Gel slices containing DNA bands of interest were excised, melted at 63°C, mixed with the appropriate restriction enzyme reagents, and incubated with an excess of the second restriction enzyme for 2 to 4 h. The resultant digests were then electrophoresed in 0.5% SeaKem ME agarose gels as described above. Transfer of viral DNA fragments to nitrocellulose strips (43), preparation of restriction enzyme fragment probes by electroelution and nick translation, and hybridization to nitrocellulose strips were performed as previously described (24, 25).

Calculations of map coordinates. For the calculation of map coordinates within the *U_L* sequences of the *ts*LB2 defective genomes, the *Bgl*III site, located at the junction between *Bgl*III fragments I and D, was assumed to be at 41.15×10^6 MW from the left end of the genome (26; G. S. Hayward, T. Buchman, and B. Roizman, unpublished data). The locations of *Bam*I, *Kpn*I, and *Sal*I sites relative to this *Bgl*III site were taken from previously published (26) restriction enzyme maps of HSV-1 (Justin). The relevant fragments generated from the *Bgl*III-I fragment of *ts*LB2 PO DNA comigrated with those of HSV-1 (Justin) DNA. When converting to map units, we assumed that HSV-1 DNA was 96×10^6 in MW.

Analyses of ICPs. Confluent HEp-2 cells in 96-well culture dishes (6×10^4 cells per well) were infected with PO or serially passaged virus stocks at the multiplicities of infection (MOIs) specified in the text. Infection was in medium 199 (KC Biologicals, Inc.) supplemented with 1% heat-inactivated calf serum (199-V). After 1 h of virus adsorption, the inocula were removed and the cells were overlaid with 199-V. For labeling of infected cell polypeptides (ICPs), cell monolayers were rinsed three times with medium 199 lacking leucine, isoleucine, and valine and containing 1% dialyzed calf serum. The cell monolayers were then overlaid with labeling medium which consisted of medium 199 containing 10% of the concentration of unlabeled leucine, isoleucine, and valine; 1% dialyzed calf serum; and 2 μ Ci (each) of 14 C-labeled leucine, isoleucine, and valine per ml.

In experiments performed at the nonpermissive temperature, the tissue culture plates were immersed under water heated to the appropriate temperature in a circulating bath which was placed in a 37°C environ-

mental room to obtain rapid temperature equilibration during the overlay, rinsing, and labeling procedures. After the pulse, the infected cells were rinsed with ice-cold phosphate-buffered saline and lysed in the dish by the addition of a disruption mixture consisting of 0.05 M Tris-hydrochloride (pH 7.0), 2% SDS, 5% β -mercaptoethanol, 3% sucrose, and 20 μ g of bromophenol blue per ml. The cell lysates were sonicated for 30 s and heated to 100°C for 2 min before loading on SDS-polyacrylamide gels. Electrophoresis of ICPs was in 20-cm-long slab gels basically as described by Morse et al. (31), with some modifications. Specifically, 9.2% separation gels contained 0.375 M Tris-hydrochloride (pH 8.8), 0.1% (wt/vol) SDS, 0.08% (wt/vol) ammonium persulfate, 9.2% (wt/vol) acrylamide, 0.238% (wt/vol) *N,N'*-diallyltartardiamide, and 0.08% (vol/vol) *N,N,N',N'*-tetramethylethylenediamine. The 12% gels contained 12% (wt/vol) acrylamide and 0.31% (wt/vol) *N,N'*-diallyltartardiamide. The stacker gel contained 0.122 M Tris-hydrochloride (pH 6.8), 0.1% (wt/vol) SDS, 0.2% (wt/vol) ammonium persulfate, 3% (wt/vol) acrylamide, 0.078% (wt/vol) *N,N'*-methylenebisacrylamide, and 0.08% (vol/vol) *N,N,N',N'*-tetramethylethylenediamine. All reagents were obtained from Bio-Rad Laboratories. Electrophoresis was at 10 mA for 16 to 18 h, using electrode buffer containing 0.025 M Tris, 0.192 M glycine, and 0.1% SDS (pH 8.5). After electrophoresis, the gel was fixed and stained for 30 min in 25% isopropanol–10% acetic acid–0.03% (wt/vol) Coomassie brilliant blue and then destained (2 $\times$ 15 min) in 10% isopropanol–10% acetic acid. The gel was then rinsed in water, dried, and exposed for autoradiography. ICP designations were based on measured molecular weights, following the nomenclature of Morse et al. (31).

RESULTS

Infectious virus production by *tsLB2* passages. As detailed above, the *tsLB2* series was derived from PO virus by sequential propagation of virus stocks at low (1:4) dilutions. Figure 1 shows the yield of infectious virus produced per passage during propagation of the series for 39 passages. The serial undiluted propagation of the *tsLB2* series was accompanied by large fluctuations in the amount of infectious virus produced per passage reflecting the cyclic accumulation of defective interfering virus particles (18, 48).

Physical mapping of the *tsLB2* defective virus genomes. Analyses of viral DNA extracted from cells infected with various *tsLB2* passages revealed that each of the serially passaged virus DNA preparations consisted of a mixture of standard helper virus and defective virus genomes with grossly altered structures.

Although these defective virus genomes exhibited a buoyant density indistinguishable from that of standard virus DNA in CsCl gradients (data not shown), they clearly differed from standard virus DNA in that they were resistant to cleavage by a number of restriction enzymes,

FIG. 1. Infectious virus yields in the course of serial undiluted propagation of *tsLB2* virus. The yield of infectious virus per passage was calculated from the titers of the various *tsLB2* stocks on Vero cells.

including *HindIII*, *EcoRI*, *HpaI*, *XbaI*, and *BglII*, and exhibited a distinct fragmentation pattern when digested with *KpnI*, *BamI*, and *SalI* (Fig. 2).

For physical mapping of defective *tsLB2* genomes, viral DNA prepared from cells infected with *tsLB2* passage 17 (P17) was doubly digested with *HindIII* and *EcoRI*, and enzyme-resistant, defective virus DNA (*Hin/EcoR* DNA; approximately 100×10^6 in MW) was recovered free of helper virus DNA (which was sensitive to these enzymes) by either electrophoresis in agarose gels or sedimentation in 10 to 30% sucrose velocity gradients (8). The structure of P17 *Hin/EcoR* defective virus DNA molecules was then determined by various restriction enzyme analyses, including single and double digestion with *KpnI*, *BamI*, and *SalI* (Fig. 2); partial cleavages with these enzymes (Fig. 2); sequential digestion of individual fragments generated by one enzyme with a second restriction enzyme (Fig. 3); and determination of sequence relatedness by blot hybridizations (Fig. 3). The results of these analyses yielded the physical maps shown in Fig. 4 and can be summarized as follows. (i) The *tsLB2* P17 *Hin/EcoR* DNA molecules consisted of repeat units arranged in a head-to-tail tandem array. (ii) Digestion of these reiterated defective virus DNA molecules with *KpnI*, *BamI*, and *SalI* yielded, in each case, both major and minor fragments which were mapped within abundant and less-abundant types of defective virus genomes. The major type of the *Hin/EcoR* DNA molecules comprised 80% of the defective virus DNA in the *tsLB2* P17-infected cells and consisted of repeat units (type I, Fig. 4) 4.9×10^6 in MW. Two additional types of

FIG. 2. Restriction enzyme analyses of PO and serially passaged *tsLB2* virus DNA. Lanes 1 through 10, Autoradiograms of electrophoretically separated fragments of ^{32}P -labeled PO and P17 *tsLB2* DNAs digested with the enzymes shown. R, DNA resistant to the restriction enzyme. Lanes 11 through 16, ^{32}P -labeled P17 *tsLB2* DNA (17) or purified P17 *Hin/EcoR* DNA (R) digested with *Bam*I, *Kpn*I, and *Sal*I. Numbers represent MW $\times 10^{-6}$. Lanes 17 through 19, Partial digestion of P17 *Hin/EcoR* DNA with *Bam*I (B), *Kpn*I (K), and *Sal*I (S). $\circ$, Complete digestion products; $\bullet$, partial digestion products corresponding to one or two full-sized repeat units of molecule I (PI), molecule II (PII), or molecule III (PIII) shown in Fig. 4. These partial digest products were generated by cleavage every second or fourth restriction site along the concatemeric defective virus DNA molecules and are identical in the *Bam*I, *Kpn*I, and *Sal*I patterns. $\square$, Partial digestion products unique to each of the restriction enzymes described above; digestion occurred every third site along the concatemeric defective virus DNA molecules. Lanes 20 through 22, Digestion of *Hin/EcoR* DNA with combinations of *Bam*I (B), *Sal*I (S), and *Kpn*I (K).

defective *tsLB2* genomes comprised 10 and 6% of the P17 *Hin/EcoR* DNA and consisted of repeat units (types II and III, Fig. 4) 5.1×10^6 and 6.3×10^6 , respectively, in MW. (iii) The three types of repeat units (I, II, and III) shared the majority of their DNA sequences, as evident from both the arrangement of restriction enzyme sites (see Fig. 4) and hybridizations of ^{32}P -labeled individual *Kpn*I/*Bam*I and *Bam*I/*Sal*I fragments of P17 *Hin/EcoR* DNA to nitrocellulose strips (43) containing the homologous, unlabeled DNA fragments (Fig. 3). As shown in Fig. 4, the variability between the three types of repeat units could be localized within two regions. First, repeat unit II contained a *Bam*I/*Sal*I fragment, 1.44×10^6 in MW, which shared sequences with the equivalent *Bam*I/*Sal*I fragment, 1.26×10^6 in MW, present in repeat units I and III. Second, repeat unit III contained a *Sal*I/*Kpn*I fragment, 1.70×10^6 in MW, in place

of a fragment, 0.34×10^6 in MW, present in the smaller repeats I and II. (iv) Experiments involving the partial digestion of purified defective virus DNA with *Kpn*I, *Bam*I, and *Sal*I (Fig. 2; additional data not shown) have revealed that the defective *tsLB2* genomes consisted of long stretches of identical repeat units (representing homopolymers), with minimal intermixing of the different-sized repeat units within the same defective virus DNA molecule. (v) The *tsLB2* P17-infected cells contained several additional *Hin/EcoR* defective virus DNA species which were present in very low abundance and which were not further characterized.

Origin of *tsLB2* defective genomes. Several sets of studies were done to identify the standard virus DNA sequences contained within repeat units of *tsLB2* defective genomes. These included hybridizations of ^{32}P -labeled P17 *Hin/EcoR* DNA and individual restriction enzyme frag-

FIG. 5. Origin of *tsLB2* defective genomes. (A) Hybridization of ³²P-labeled PO (control), Justin P15 HD (P15 Justin), and *tsLB2* P17 *Hin*/*EcoR* (P17 LB2) DNAs and the individual *Bam*I/*Sal*I fragments of *tsLB2* P17 *Hin*/*EcoR* DNA to nitrocellulose strips containing unlabeled *Kpn*I (lanes 1 through 3) and *Bam*I (lanes 4 through 11) fragments of PO DNA and to the unlabeled *Bam*I/*Sal*I fragments (25) of Justin P15 HD DNA (lanes 12 and 13). (B) *Bam*I and *Kpn*I restriction enzyme maps of standard HSV-1 (Justin) DNA (26). The horizontal lines indicate the PO DNA fragments which show homology to the various probes described above. Heavy horizontal lines, thin lines, and dotted lines represent relative intensities of hybridization bands.

ments derived thereof to nitrocellulose strips containing unlabeled restriction enzyme fragments of standard virus DNA and, conversely, hybridizations of ^{32}P -labeled fragments of standard virus DNA to nitrocellulose strips containing fragments of *tsLB2* P17 *Hin/EcoR* DNA. We also defined by hybridizations the regions of homology between *tsLB2* defective genomes and class I HD defective genomes derived from HSV-1 (Justin). Autoradiographic exposures of the hybridized strips are shown in Fig. 5 through 7. The results of these hybridizations, in conjunction with data presented above regarding the size and sequence relatedness of the three types of repeat units of the *tsLB2* defective genomes,

have revealed the following. (i) The majority of the *tsLB2* *Hin/EcoR* DNA sequences were homologous to standard virus DNA sequences located within the U_L region between map coordinates 0.356 and 0.423 (defined by strong hybridization to *KpnI* fragments N, P, and V; *BamI* fragments G and V; and *SalI* fragments X, R, and N; Fig. 5, lanes 3 and 6 through 11, and Fig. 6, lanes 4 and 5). A minority of the defective virus DNA sequences showed homology to the S component sequences within map coordinates 0.978 and 1.00 or 0.824 and 0.846 (*KpnI* fragments C, D, I, and K; *BamI* fragments K and R; and *SalI* fragments A and J). These S end sequences included at least a portion of the

FIG. 6. Lack of detectable homology between the U_L sequences within map coordinates 0.322 and 0.429 and the S end sequences of *tsLB2* PO DNA. (A) Hybridizations of ^{32}P -labeled probes to nitrocellulose strips containing the separated *SalI* fragments of *tsLB2* PO DNA. The probes used were PO *tsLB2* DNA (lane 1); *BglIII* fragment I of PO *tsLB2* PO DNA, spanning map coordinates 0.322 and 0.429 (lane 2); *EcoRI* fragment K of *tsLB2* PO DNA spanning map coordinates 0.963 and 1.00 (lane 3); *tsLB2* P17 *Hin/EcoR* DNA (lane 4); and the *SalI* 1.15 $\times 10^6$ -MW fragment of *tsLB2* P17 *Hin/EcoR* DNA (lane 5). (B) Schematic diagram showing the locations of *EcoRI*-K and *BglIII*-I and the corresponding *SalI* fragments within standard HSV-1 (Justin) DNA. Horizontal lines indicate PO DNA fragments which hybridize with the probes shown.

FIG. 7. Organization of U_L DNA sequences within the repeat units of P17 *Hin/EcoR* DNA. (A) Hybridization of ^{32}P -labeled probes to the unlabeled *BamI/SalI* fragments of *tsLB2* P17 *Hin/EcoR* DNA which were immobilized on nitrocellulose strips. The probes used were P17 *tsLB2 Hin/EcoR* DNA (LB2 R), Justin P15 HD DNA (JUS HD), and *tsLB2 PO SalI* fragments N, R, and X located between map coordinates 0.354 and 0.429 of *tsLB2 PO* DNA. (B) Schematic diagram of map coordinates 0.354 to 0.429 of *PO* DNA showing the locations of *SalI* (S) fragments N, R, and X and *BamI* (B) fragments V and G. Numbers above fragment designations represent $\text{MW} \times 10^{-6}$. (C) Diagrammatic representation of the sequence homology between the *SalI* fragments of *PO* DNA and the *BamI/SalI* fragments of the three types of *tsLB2* repeat units. Also shown are the areas of homology between *tsLB2* P17 *Hin/EcoR* DNA and Justin P15 HD DNA.

terminal reiteration *a* sequences (11, 38), since the *tsLB2* P17 *Hin/EcoR* DNA probes also exhibited low levels of hybridization to the L end fragments. On the basis of the relative intensities of the hybridization bands corresponding to the U_L and S fragments, we estimated that the major 4.9×10^6 -MW repeat units of the *tsLB2* defective genomes contained approximately 4.4×10^6 -MW DNA sequences homologous to U_L DNA sequences and 0.5×10^6 -MW sequences homologous to terminal portions of the *ac* inverted repeat sequences of the S component. (ii) The observed hybridizations of the *tsLB2* defective virus DNA probes to the terminal sequences of the S component could not be attributed to an existing homology in *PO* DNA between the corresponding U_L and S end sequences. Thus, no cross-hybridization could be detected among fragments derived from these regions of stan-

dard virus DNA (Fig. 6, lanes 2 and 3). Therefore, the *tsLB2* defective genome repeat units must have evolved as a result of recombinational events leading to covalent linkage between these separate sets of HSV DNA sequences. (iii) In agreement with the above data, class I and II defective HSV genomes, represented by Justin HD and *tsLB2 Hin/EcoR* DNAs, respectively, shared only those DNA sequences corresponding to the terminal portions of the S component of standard HSV-1 DNA (Fig. 5, lane 13). (iv) The internal arrangement of the U_L and S end sequences within the three types of *tsLB2* repeat units is summarized in Fig. 8. The majority of the DNA sequences within repeat unit III (5.1×10^6 of the 6.3×10^6 -MW repeat) were colinear with the U_L sequences located between the *SalI* site at map coordinate 0.370 and the *BamI* site at map coordinate 0.423 of standard HSV-1 DNA

FIG. 8. Arrangement of *ac* and U_L DNA sequences within *tsLB2* defective genomes. (Top) Schematic representation of standard virus (parental) DNA showing the arrangement of *Bam*I (B), *Kpn*I (K), and *Sal*I (S) restriction enzyme sites within the corresponding U_L sequences. (Bottom) Representation of the arrangement of the U_L and *ac* sequences in three classes (I, II, and III) of *tsLB2* *Hin*/*Eco*R DNA molecules. The U_L sequences represented by the thin line are deleted in repeat units I and II.

(Fig. 8). This was apparent from both the patterns of hybridizations (Fig. 5 through 7) and the similar sizes of the corresponding restriction enzyme fragments generated from repeat unit III DNA and from the above-described U_L region of standard virus DNA. Repeat units I and II contained a 1.35×10^6 -MW deletion within the 2.5×10^6 -MW *Sal*I fragment R of PO DNA. Thus, the two *Sal*I sites bounding this fragment at map coordinates 0.370 and 0.396 were conserved in repeat units I and II but were now bounding a *Sal*I fragment, 1.15×10^6 in MW, which hybridized solely to the 2.5×10^6 -MW *Sal*I fragment R of *tsLB2* PO DNA (Fig. 6, lane 5). As already noted above, this deletion could be more precisely localized within the *Sal*I/*Kpn*I 1.7×10^6 -MW fragment of repeat unit III (i.e., to within map coordinates 0.370 and 0.388 of standard virus DNA), which in repeat units I and II was replaced by a 0.34×10^6 -MW fragment.

Finally, the junction between the U_L and S end sequences could be localized within the 1.26×10^6 -MW *Bam*I/*Sal*I fragment of repeat units I and III and within the 1.44×10^6 -MW *Bam*I/*Sal*I fragment of repeat unit II. As discussed above, we estimated that approximately 0.5×10^6 of the 1.26×10^6 -MW fragment corresponded to DNA

sequences arising from the S component, whereas the remaining portions of this fragment corresponded to sequences extending to the left of the *Sal*I site at coordinate 0.370 and to the right of the *Bam*I site at coordinate 0.423 of standard virus DNA. Indeed, the *tsLB2* P17 *Hin*/*Eco*R probes hybridized at low levels to the corresponding *Sal*I fragment X (Fig. 6, lane 4, and Fig. 7, lane 4) and the *Bam*I fragment Q (Fig. 5, lane 9) of standard HSV DNA.

Comparison of viral DNA from different *tsLB2* passages. Analyses of the cleavage patterns of 32 P-labeled viral DNAs prepared from cells infected with passages 0 through 35 of the *tsLB2* series are shown in Fig. 9. The majority of defective virus DNA molecules present in the various passages of the series corresponded to the three types of defective virus genomes described above for P17 virus. However, later passages appeared to contain lower proportions of the defective genome species consisting of the largest (6.3×10^6 -MW) repeats. New species of defective genomes, if such were generated de novo, were not significantly amplified in the course of this prolonged serial propagation.

Quantitative estimates for the overall fraction of defective genomes produced in cells infected

FIG. 9. *KpnI* restriction enzyme patterns of viral DNA from cells infected with different *tsLB2* passages. ^{32}P -labeled viral DNA was prepared from cells infected with the *tsLB2* passages at the MOI shown. Numbers to the left represent $\text{MW} \times 10^{-6}$. P = 4.9, Partial digest product corresponding to monomers of repeat unit I (i.e., sum of 2.1×10^6 and 2.8×10^6 uncleaved fragments).

with various *tsLB2* passages were obtained by scanning adequate autoradiographic exposures of gels containing the *KpnI* (Fig. 9) and *BamI* (data not shown) digests of serially passaged virus DNAs. These analyses (Fig. 10) revealed

cyclic fluctuations in the fraction of defective virus genomes produced in cells infected with the various passages.

Correlation between the fraction of defective virus genomes and reduced yields of infectious

FIG. 10. Correlation between autointerference and the fraction of defective genomes. For interference assays, replicate HEP-2 cell cultures were infected with *tsLB2* PO and passages 7 through 35 of the *tsLB2* series. After 24 h of infection, the yield of infectious virus from each culture was determined by titration on Vero cells. The MOI of passages 7 through 35 used in the interference assays were the same as those employed in the preparation of ^{32}P -labeled viral DNA shown in Fig. 9. The MOI of PO infections was varied accordingly. Interference (●—●) denotes the total reduction in infectious virus produced during 24-h infections with the various passages as compared with virus yield in cells infected with *tsLB2* PO at the matching MOI. The fractions of defective genomes (★—★) were calculated from scans of autoradiograms of gels containing *BamI* and *KpnI* (Fig. 9) digests of serially passaged virus DNAs. Different autoradiographic exposures were used for different passages to allow adequate scanning. The graph points denote averages of the values obtained from the *BamI* and the *KpnI* data.

virus. The undiluted propagation of the *ts*LB2 virus stocks resulted in a typical von Magnus-type interference effect (18, 48), with infectious virus yield per culture (passage) changing in a cyclic pattern (Fig. 1). To determine whether these observed fluctuations in the yield of infectious virus correlated with the abundance of defective genomes in the various serially passaged virus stocks, we assayed the extent of autointerference exhibited by the various *ts*LB2 passages under constant and relatively high input MOIs. Such analyses were necessary because, in the course of the serial (1:4 dilution) propagation, the input MOI used to generate the various *ts*LB2 passages varied from a low of 10^{-3} PFU per cell to a high of 5 PFU per cell. In consequence, the observed fluctuations in the yield of infectious virus per passage reflected not only the differential ability to interfere but also variable numbers of infected cells per culture. In the present autointerference experiment, we compared the production of infectious virus in cells infected with serially passaged virus stocks (P1 through P35) with the yield of

infectious virus in cells infected with an equal number of PFU of the standard (PO) virus stock per cell.

The results of these autointerference assays are shown in Fig. 10, along with the estimated fractions of defective genomes produced in the course of infections with the corresponding passages. The capacity of the various passages to produce infectious progeny fluctuated in synchrony with the fluctuations in the fraction of defective virus genomes produced by the corresponding virus stocks (Fig. 10). Thus, the level of interference exhibited by the various *ts*LB2 virus stocks correlated well with the abundance of the class II defective virus genomes.

Polypeptide synthesis in cells infected with *ts*LB2 passages. To determine whether the presence of the class II defective HSV genomes had an effect on the pattern of viral gene expression in infected cells, replicate HEP-2 cell cultures were infected at 34°C with 0.5 to 5 PFU of selected *ts*LB2 passages per cell. After incubation of the infected cells with ^{14}C amino acids at 9 to 15 or 10 to 12 h postinfection, the cells were

FIG. 11. ICP synthesis in cells infected with serially passaged *ts*LB2 stocks. HEP-2 cells in 96-well culture dishes were mock infected or infected at 34°C with PO and the *ts*LB2 passages as shown, at input PFU per cell as indicated. Virus adsorption was done in a total volume of 50 μl per well, except for passages 11, 12, and 21, in which 200 μl was used (owing to the low titers of these virus stocks). The cells were pulsed-labeled with ^{14}C -amino acids at the times shown. ICP nomenclature is that of Morse et al. (31).

harvested, and the ICPs were analyzed by electrophoresis in SDS-polyacrylamide gels. The results of these studies (Fig. 11 and 12) can be summarized as follows. (i) Analyses of cell cultures infected with passages 3 through 22 and labeled at 9 to 15 h postinfection (Fig. 11) revealed that infections with virus stocks containing relatively high proportions of defective genomes (notably P9, P18, and P19) resulted in the abundant synthesis of a protein 128×10^3 in MW. This overproduced protein was identified as the standard virus-ICP number 8 (ICP8) on the basis of its electrophoretic mobility and the pattern of its regulation in the serially passaged virus-infected cells, as outlined below. Furthermore, the gene encoding ICP8 had previously (2, 29, 31) been mapped to the U_L sequences contained within the $tsLB2$ defective genomes, thus supporting the hypothesis that the overproduced polypeptide was expressed from multiple ICP8 gene templates within the defective $tsLB2$ genome repeat units. (ii) Analyses of cell cultures infected with selected passages from P8 through P36 and labeled at 10 to 12 h postinfection (Fig. 12) revealed that cells exposed to highly defective virus passages (notably P9, P18, and P33) produced, in addition to abnormally

high quantities of ICP8, a polypeptide of estimated MW 74.5×10^3 (in 9% gels; 77.5×10^3 in 12% gels). We designated this polypeptide defective virus-induced ICP 20.5 (DICP20.5) because it migrated in 9% gels slightly faster than the 77×10^3 -MW polypeptide produced in standard HSV-1-infected cells and designated by Morse et al. (31) as ICP20. As discussed below, it is at present unclear whether DICP20.5 is also produced during standard virus infections. Additional polypeptides appeared to be abnormally made in cells infected with a more limited subset of passages (notably P33). The nature of these polypeptides is currently under investigation. (iii) The overproduction of ICP8 was clearly apparent in both the 10- to 12-h and the 9- to 15-h pulse-labeled samples. In contrast, the synthesis of DICP20.5 appeared to be more pronounced in the infected cell samples which were labeled during the shorter (10- to 12-h) pulse. As seen below, these different patterns reflected, at least in part, differences in the stabilities of these polypeptides in the serially passaged virus-infected cells. (iv) Cells infected with a number of passages appeared to synthesize reduced amounts of many of the ICPs. However, in some cases, this generalized reduction in the synthesis

FIG. 12. ICP synthesis in cells infected with serially passaged *tsLB2* stocks. HEp-2 cells in 96-well dishes were mock infected or infected at 34°C with the *tsLB2* virus passages as shown. Virus adsorption with passages 9 through 20 was done in a total volume of $50 \mu\text{l}$, except for passage 11, in which $200 \mu\text{l}$ was used. ICPs were analyzed in 9% SDS-polyacrylamide gels. Virus adsorption with passages 23 through 36 was done in a total volume of $20 \mu\text{l}$, except for P27, P28, and P29, for which 200 , 150 , and $30 \mu\text{l}$, respectively, were used. P31 was used at 50 PFU per cell, owing to an error. ICP analysis done was in a 12% gel. $\circ$, β polypeptide; $\bullet$, γ polypeptide. The designations of Morse et al. (31) were followed.

of ICPs might have resulted, at least in part, from the use of a lower effective input MOI which was dictated by the low titers of the corresponding virus passages. Reduced synthesis of ICPs was observed also when the MOI of standard virus infection was lower than 1 PFU per cell (Fig. 12, lane 13).

Regulation of ICP8 production in serially passaged virus-infected cells. HSV ICPs were previously classified into three general regulatory classes, α , β , and γ , differing with respect to the kinetics of their syntheses and the requirements for their production in infected cells (16, 17). Previous analyses of viral gene expression in cells infected with HSV-1 temperature-sensitive mutants of the 1-2 complementation group (39), including *tsLB2*, showed that the sequential transition from the α (immediate early) to the β (early) and γ (late) stages of expression was dependent, at least in part, upon the activity of the α ICP4 (=VP175) shown to be affected by the 1-2 mutations (3, 4, 6, 23, 31, 34, 36, 37, 45). Thus, infections of cells at the nonpermissive temperature with various members of the 1-2 complementation group, including *tsLB2*, resulted in overproduction of the α viral polypeptides and in the reduced or unapparent synthesis of many of the later β and γ ICPs (4, 6, 13, 15, 23, 28, 36). Moreover, transcriptional analyses and studies of viral polypeptides produced in temperature shift protocols led to the conclusion that the function of the 1-2 gene product (i.e., ICP4) was continuously required for the synthesis of β and γ mRNA, inasmuch as viral transcription under nonpermissive conditions was limited to α genes (6, 35-37, 52, 53).

ICP8 had been previously assigned to the β kinetic class of viral polypeptides (16), the synthesis of which was dependent on the presence of a functional ICP4. It was therefore of interest to determine whether ICP8 synthesis during serially passaged *tsLB2* infections followed the β regulatory pattern at the permissive temperature and whether the mutation in the *tsLB2* helper virus affected the expression of ICP8 from defective virus genomes at the nonpermissive temperature. Thus, in the experiment shown in Fig. 13, we followed the kinetics of ICP8 synthesis in cells infected at 34°C with 15 PFU of either PO or *tsLB2* P19a per cell. P19a was a relatively concentrated virus stock containing more than 90% defective genomes (Fig. 13) and was derived by high-scale propagation of virus starting from P16 of the original *tsLB2* series. The PO and P19a-infected cells were pulse-labeled with ^{14}C -amino acids at 2 to 4, 6 to 8, 10 to 12, and 14 to 16 h postinfection. The infection of cells with P19a virus resulted in the pronounced synthesis of ICP8 and in significant reduction and delay in the synthesis of all other

viral polypeptides (Fig. 13). The synthesis of ICP8 was maximal at 10 to 12 h postinfection and thereafter declined, resembling the delayed synthesis and decline of other β class viral polypeptides (e.g., ICPs 6, 39, and 40) in the P19a-infected cells.

In a second experiment (Fig. 14), HEp-2 cells were infected at 34 and 39°C with 15 PFU of standard PO or P19a *tsLB2* virus stocks per cell and labeled with ^{14}C -amino acids at 3 to 11 h postinfection. Parallel cultures were similarly infected with PO or P19a viruses in the presence of 500 μg of canavanine (an analogue of arginine) per ml. Canavanine was shown previously to affect the synthesis of many β and γ viral polypeptides while allowing efficient synthesis of the α and a subset of β ICPs in standard HSV-infected cells (17, 33). A third set of cultures was infected at both temperatures in the presence of 300 μg of phosphonoacetate per ml, which was shown previously (27) to inhibit the replication of viral DNA and to reduce the synthesis of a subset of the late γ viral polypeptides (20).

The results of these analyses (Fig. 14) can be summarized as follows. (i) In agreement with previous reports, infection of cells with *tsLB2* PO virus under nonpermissive conditions resulted in most abundant synthesis of the α polypeptides (e.g., ICPs 4, 0, and 27) and in reduced synthesis of many β and γ polypeptides (e.g., ICPs 5, 20, 25, 39, and 40). (ii) Cells infected with P19a virus at 39°C also produced abnormally high amounts of the α polypeptides (ICPs 4, 0, and 27). However, contrary to expectations, the P19a-infected cells synthesized substantial amounts of ICP8 at the nonpermissive temperature. Thus, the *tsLB2* mutation within the helper virus genomes did not significantly affect the synthesis of ICP8 under nonpermissive conditions. (iii) The addition of canavanine to cells infected at the permissive temperature with the PO virus stock resulted in the expected reduction in the synthesis of many of the β and γ viral polypeptides. The addition of canavanine to cells infected with P19a virus at the permissive or the nonpermissive temperature resulted, in both cases, in a pronounced inhibition of ICP8 synthesis. Thus, the expression of ICP8 in the P19a-infected cells required the presence of one or more functional ICPs. (iv) Incubation of infected cells in the presence of phosphonoacetate resulted in an expected reduction in the synthesis of many of the late viral polypeptides (e.g., ICPs 5, 15, 20, 25, 43, and 44). The drug, however, did not significantly reduce the synthesis of ICP8 in P19a-infected cells. Thus, the overproduced ICP8 was mostly expressed from parental defective virus genomes.

Stability of overproduced ICP8 and DICP20.5. As discussed above, the 128×10^3 -MW poly-

FIG. 13. Time course of ICP synthesis in cells infected with PO and P19a *ts*LB2 virus. (A) *Bam*I restriction enzyme patterns of 32 P-labeled viral DNAs prepared from cells infected with 1 PFU of PO and P19a viruses per cell. P19a-infected cells synthesized more than 90% defective virus genomes. (B) Hep-2 cells in 96-well dishes were mock infected or infected at 34°C with PO and P19a viruses at 15 PFU per cell. The cells were harvested immediately after the 14 C-amino acid pulses at the times shown. Electrophoresis was performed in a 9% gel.

peptide which we have identified as ICP8 appeared to be overproduced in infected cells labeled during both the shorter (10- to 12-h) and longer (9- to 15-h) pulses, whereas the abundant production of DICP20.5 was apparent only in the cultures labeled for the shorter time intervals. To determine whether these different labeling patterns reflected differences in the rates of turnover of the two ICPs, we examined the stability of the overproduced ICP8 and DICP20.5 during pulse-chase protocols. Tests for the stability of the abundant ICP8 in serially passaged virus infections at 39°C were of added interest because they could confirm the identification of the overproduced 128×10^3 -MW polypeptide as ICP8 rather than as the α polypeptide ICP0, which migrated in 9% SDS-polyacrylamide gels in close proximity to ICP8 and which was overproduced in cells infected at 39°C with *ts*LB2 defective virus stocks. Thus, it was previously shown by Wilcox et al. (54) for

standard virus infections that the α ICP0 was characteristically less stable than the β ICP8.

Duplicate Hep-2 cell cultures were infected at 39°C with 10 PFU of *ts*LB2 P9 virus per cell in the absence of added drugs or in the presence of 500 μ g of canavanine per ml. After a 14 C-amino acid pulse at 6 to 8 h postinfection, one set of cultures was harvested immediately after the pulse, whereas the parallel set of cultures was further incubated from 8 to 15 h postinfection in medium containing excess unlabeled amino acids. Two additional cultures were similarly infected at 39°C with P9 virus in the presence of 50 μ g of cycloheximide per ml. At 5 h postinfection, the cycloheximide was removed, and the cells were labeled for 1 h with 14 C-amino acids and either harvested immediately after the pulse or incubated for an additional 6 h in the presence of excess unlabeled amino acids. As originally shown by Honess and Roizman (16), only α polypeptides (including ICP0) are synthesized

FIG. 14. ICP synthesis in cells infected with *tsLB2* virus stocks at the permissive and the nonpermissive temperature in the absence and presence of drugs. HEP-2 cells in 96-well dishes were mock infected (M) or infected at 34 or 39°C with the PO and P19a *tsLB2* stocks. Infection was in the absence of drugs (—), in the presence of 500 μ g of canavanine (Can.) per ml, or in the presence of 300 μ g of phosphonoacetate per ml. The cells were labeled at 3 to 11 h postinfection and harvested immediately after the pulse. Proteins are designated as α ($\square$), β ($\circ$), and γ ($\bullet$) polypeptides according to the designations of Morse et al. (31). The bottom part of the figure is a higher magnification of the ICP8/0 region of the autoradiogram shown above.

immediately after the reversal of cycloheximide. Thus, the stability of the α ICP0 synthesized in cycloheximide-treated cells could be compared with that of the overproduced ICP8 in infected cells which were not treated with the drug.

The results of these analyses (Fig. 15) revealed a relatively rapid turnover of D1CP20.5 in cells infected in the absence of added drugs. In contrast, ICP8, which was synthesized during the 2-h pulse in the absence of drugs, appeared

FIG. 15. Stability of ICP8 and DICP20.5. Lanes 1 through 3, HEp-2 cells mock infected or infected with P9 *tsLB2* virus in the presence of 50 μ g of cycloheximide per ml. At 5 h, the drug was removed and the cells were pulsed with 14 C-amino acids from 5 to 6 h postinfection (lanes 1 and 3) or from 5 to 6 h postinfection and then chased from 6 to 12 h (lane 2) postinfection. Lanes 4 through 8, HEp-2 cells mock infected or infected with 10 PFU of P9 virus per cell in the presence or absence of canavanine. The cells were either pulsed with 14 C-amino acids for 6 to 8 h (lanes 4, 6, and 8) or pulse-labeled for 6 to 8 h and then chased with excess unlabeled amino acids at 8 to 15 h postinfection (lanes 5 and 7). In the canavanine-treated cell cultures, the drug was present throughout the pulse and chase periods.

to be stable throughout the subsequent chase period and, in this respect, differed significantly from the unstable ICP0, which was synthesized in the cultures treated initially with cycloheximide. As was the case with P19a virus infections, the addition of canavanine prevented the overproduction of ICP8. Moreover, canavanine treatment also inhibited the synthesis of DICP20.5, indicating that the synthesis of this polypeptide required the presence of one or more functional ICPs.

DISCUSSION

We have described in this paper our studies concerning the structural features of defective virus genomes present in serially propagated virus populations derived from the *tsLB2* mutant of HSV-1 (HFEM). Several of these structural features merit further discussion. First, like their class I counterparts, the *tsLB2* defective genomes, as well as other class II defective genomes propagated from different HSV strains (21, 22, 40; G. S. Hayward, personal communication; R. Spaete, L. Diess, N. Frenkel, and B. Murray, unpublished data), derive their sequences from separate regions of standard virus DNA. We have previously proposed (7, 10, 25, 46, 47) that this specific incorporation of defined subsets of viral DNA sequences within defective genome repeat units reflected the existence of two separate recognition signals which were needed for the propagation of the defective virus genomes: an origin for DNA replication located in the U_L region within map coordinates 0.356 to 0.423 and a site specifying the cleavage of viral DNA concatemers and their packaging into nucleocapsids. This last recognition site has been localized to the end of S, within the sequences shared by both classes of the defective HSV genomes (25, 46, 47). Direct support for this hypothesis has come more recently from our studies (R. Spaete and N. Frenkel, manuscript in preparation) showing that both U_L and S end sequences are required for the successful propagation of the class II defective genomes in serially passaged virus stocks.

Second, although the nature of recombinational events leading to the formation of the class II defective genome repeat units is at present unknown, the hybridization studies reported above revealed no detectable homology between the sequences located in the vicinity of the U_L and S regions which become covalently associated within the *tsLB2* repeat units. Thus, it seems unlikely that the initial linkage between the U_L and S end sequences occurred by a recombinational process involving the recognition of relatively large stretches of homologous DNA sequences.

Finally, the studies reported above have shown that the junctions between the S and U_L sequences within the *tsLB2* repeat units which we studied were located at the ends rather than at internal positions within the stretch of U_L sequences. Moreover, the recombining S end region contained the terminal reiteration α sequences shown previously to play a role in recombinational events leading to the generation of the S and L inversions of standard virus DNA (30, 38). When taken together, these structural features might imply that the generation of the

first class II defective genome repeat unit from the parental standard virus DNA is but a low-probability extension of the processes leading to the generation of the four isomeric forms of standard virus DNA. Specifically, it is possible that the S (and L) ends of standard virus DNA undergo random recombinations at low efficiencies with other, nonhomologous regions of the viral genome and that only the resultant "viable" recombinational products, containing a replication origin linked to the cleavage-packaging site become amplified and packaged using the helper virus trans replication functions. By extension of previously proposed models for standard virus DNA (30, 38), such a process could involve the folding of the S end sequences over nonhomologous U_L DNA sequences to serve either as an intermediate in an intramolecular crossover event or, alternatively, as a primer for the copying of the corresponding stretch of the U_L DNA sequences into the S terminus, followed by excision of the resultant chimeric sequences from the parental DNA.

Expression of the τ LB2 defective virus genomes. We have previously shown that infections of cells with HSV-1 (Justin) populations containing high proportions of class I defective genomes results in a reduced synthesis of many of the ICPs and in the overproduction of the α ICP4 encoded within the *ac* inverted repeat sequences which form part of the Justin defective genome repeat units (8, 10). Furthermore, our analyses of viral transcription in the serially passaged Justin-infected cells have shown that these changes in viral polypeptide synthesis reflected the increased transcription of defective genomes at the expense of transcription from helper virus DNA (7, 10; H. Locker and N. Frenkel, manuscript in preparation). Overproduction of ICP4 was also observed by Murray et al. (32) in cells infected with serially passaged HSV-1 (Patton) populations containing HD (class I) defective virus DNA.

Our studies described above concerning the effect of class II defective HSV genomes on viral gene expression have similarly revealed the reduced synthesis of many ICPs in cells infected with τ LB2 virus populations containing high proportions of defective genomes. However, as expected on the basis of the different structural features of the class I and class II defective genomes, the τ LB2-infected cells overproduced ICPs which were different from those overproduced in cells infected with the class I defective virus populations.

The identification of the overproduced polypeptides as specific viral proteins synthesized in standard virus-infected cells is, at present, based on previous information regarding standard virus polypeptides encoded by the DNA se-

quences which are represented within defective genome repeat units (2, 29, 31). Of particular relevance in this respect has been the previous report by Conley et al. (2) that the in vitro translation of infected cell RNA selected by hybridization to the *Sall/BamI* fragment spanning map coordinates 0.396 to 0.409 (see Fig. 7B) yielded two polypeptides, 128×10^3 (equal to that of ICP8) and 25×10^4 in MW, whereas transcripts selected by hybridization to the entire *BamI* fragment G (map coordinates 0.356 to 0.409) directed the synthesis of the 128×10^3 - and 25×10^3 -MW polypeptides as well as two additional polypeptides 114×10^3 and 105×10^3 in MW.

Because the τ LB2 defective genomes contain multiple copies of the sequences within map coordinates 0.370 and 0.423, i.e., overlapping a portion of the *BamI* fragment G, we concluded that the 128×10^3 -MW polypeptide which was overproduced with β kinetics in cells infected with τ LB2 passages corresponded to the standard virus ICP8. Furthermore, because ICP8 was most abundantly synthesized in τ LB2-infected cells, it is reasonable to assume that the entire ICP8 gene is contained within the most abundant repeat unit (type I; Fig. 8) and must therefore be located to the right of the deleted region in this repeat unit, i.e., within map coordinates 0.384 and 0.423.

The identification of the smaller overproduced polypeptide which we designated DICP20.5 is, on the other hand, not definitive. Specifically, this 745×10^3 -MW polypeptide appeared to be uniquely produced in the serially passaged virus-infected cells. Furthermore, it was not present among the translational products reported by Conley et al. (2) to be synthesized in vitro from standard virus mRNA homologous to the corresponding U_L viral DNA sequences. Assuming that DICP20.5 is expressed from defective virus genomes, one possible explanation of these observations takes into account the structural features of the major defective genome repeat units in the τ LB2 series. Thus, it is possible that DICP20.5 represents either a truncated polypeptide produced from the sequences flanking the 1.35×10^6 -MW deletion in the major τ LB2 repeat units or a translational product of a fusion transcript read through the junctions between adjacent repeat units. Transcriptional studies to test these models are currently in progress.

It is of special interest that the viral genes which are located within the defective genome repeat units remained responsive to the sequentially regulated scheme of viral gene expression in the infected cell. Thus, the overproduced ICP8 exhibited properties of the β class of viral polypeptide, similarly to its standard virus-encoded counterpart with regard to both the time

course of its synthesis and its sensitivity to the drugs affecting the β HSV-1-encoded polypeptides. This observation is all the more striking in that the synthesis of many viral ICPs was greatly suppressed in the serially passaged virus-infected cells. Thus, it would appear that small amounts of the regulatory gene product(s) involved in the turning on and off of the β class of viral polypeptides are sufficient for the regulation of expression of multiple copies of β gene templates within the defective genomes.

In this light, it is puzzling that the abundant synthesis of ICP8 was not significantly reduced in cells infected with the P19a virus stock at 39°C, i.e., at a temperature which was not permissive for the helper virus. Because ICP8 migrated in SDS-polyacrylamide gels in close proximity to the α ICP0 (which was overproduced at the nonpermissive temperature in *ts*LB2 infections), it is not clear from the available data whether the cells infected with PO virus indeed did not produce ICP8 under nonpermissive conditions. In contrast, ICP8 could be readily distinguished from ICP0 in the P19a virus infections because of its pronounced overproduction. Furthermore, the polypeptide which was overproduced during defective-virus infections at 39°C in the absence of drugs was clearly absent from infected cells treated with canavanine or reversed from cycloheximide treatment (Fig. 14 and 15). Thus, the presence of multiple ICP8 gene templates within the *ts*LB2 defective genomes could have resulted in at least partial compensation for the inefficiency of the mutated ICP4 effector protein. Alternatively, the abundant ICP8 synthesis during defective-virus infections at the nonpermissive temperature truly reflected the absence of a significant effect of the *ts*LB2 mutation on ICP8 synthesis. Of relevance to this last argument could be the previous reports by a number of investigators (6, 15, 28, 37) that different temperature-sensitive mutants in the 1-2 complementation group exhibited variable degrees of "leakiness" in the synthesis of certain β and γ transcripts and ICPs. Thus, the *ts*LB2 mutant could exhibit one such selective phenotype, allowing the synthesis of ICP8 while more tightly affecting the synthesis of other β and γ polypeptides. Because different members of the 1-2 complementation group have been shown to contain lesions in widely separated locations within the ICP4 gene (6, 23, 37), these observations might imply that different active sites or different conformational requirements are involved in the interactions of ICP4 with different β and γ gene templates and that these are affected to different degrees by various mutations in the ICP4 gene.

A final point regarding the biological properties of serially passaged *ts*LB2 populations con-

cerns the quantitative relation among interference, viral gene expression, and the fraction of defective genomes within the virus populations. Specifically, Schröder et al. (41) and Stegman et al. (44) previously reported the lack of quantitative correlation between the level of interference and the proportions of restriction enzyme-resistant virus DNA in serially passaged HSV-1 (ANG) populations. Based on their observation that passages which appeared to contain low proportions of enzyme-resistant defective genomes exhibited high levels of interference, they made the distinction between defective and interfering particles. In contrast, the results of our studies revealed a quantitative correlation between the level of interference exhibited by 33 different *ts*LB2 passages and the fraction of class II defective genomes produced in cells infected with these passages. It is at present not clear whether the discrepancy between these two sets of results reflects true differences between the two viral series or differences in the methodologies used to quantitate the fraction of defective virus DNA. Although no direct transcriptional data are available at present for either series, it is strongly suggested from our data that interference by class II defective HSV genomes, like that exhibited by class I Justin defective genomes (7, 10), is brought about by a reduced synthesis of the majority of the infected-cell gene products, most likely owing to transcriptional competition between amplified genes encoded within the class II defective genome repeats and all other nonamplified viral genes encoded by the standard helper virus genomes. Transcriptional analyses are currently in progress to further test this hypothesis.

ACKNOWLEDGMENTS

We thank Barbara Burckart and Glynis McCray for excellent technical assistance, Jeffrey M. Leiden for helpful discussions, Lawrence S. Morse for assistance in initial protein analyses, and an anonymous reviewer for useful comments on the manuscript.

These studies were supported by Public Health Service research grants AI-15488 and CA-19264 from the National Cancer Institute and by National Science Foundation grant PCM-78-16298.

LITERATURE CITED

1. Bronson, D. L., G. R. Dreesman, N. Biswal, and M. Benyesh-Melnick. 1973. Defective virions of herpes simplex viruses. *Intervirology* 1:141-153.
2. Conley, A. J., D. M. Knipe, P. C. Jones, and B. Roizman. 1981. Molecular genetics of herpes simplex virus. VII. Characterization of a temperature-sensitive mutant produced by in vitro mutagenesis and defective in DNA synthesis and accumulation of γ polypeptides. *J. Virol.* 37:191-206.
3. Courtney, R. J., and M. Benyesh-Melnick. 1974. Isolation and characterization of a large molecular-weight polypeptide of herpes simplex virus type 1. *Virology* 62:539-551.

4. Courtney, R. J., P. A. Schaffer, and K. L. Powell. 1976. Synthesis of virus specific polypeptide by temperature sensitive mutants of herpes simplex virus type 1. *Virology* 75:306-318.
5. Delius, H., and J. B. Clements. 1976. A partial denaturation map of herpes simplex virus type 1 DNA: evidence for inversion of the unique DNA regions. *J. Gen. Virol.* 33:125-133.
6. Dixon, R. A. F., and P. A. Schaffer. 1980. Fine-structure mapping and functional analysis of temperature-sensitive mutants in the gene encoding the herpes simplex virus type 1 immediate early protein VP175. *J. Virol.* 36:189-203.
7. Frenkel, N. 1981. Defective interfering herpes viruses, p. 91-120. In A. J. Nahmias, W. R. Dowdle, and R. S. Schinazy (ed.), *The human herpesviruses—an interdisciplinary perspective*. Elsevier/North-Holland Publishing Co., New York.
8. Frenkel, N., R. G. Jacob, R. W. Honess, G. S. Hayward, H. Locker, and B. Roizman. 1975. Anatomy of herpes simplex virus DNA. II. Characterization of defective DNA molecules and biological properties of virus population containing them. *J. Virol.* 16:153-167.
9. Frenkel, N., H. Locker, W. Batterson, G. S. Hayward, and B. Roizman. 1976. Anatomy of herpes simplex virus DNA. VI. Defective DNA originates from the S component. *J. Virol.* 20:527-531.
10. Frenkel, N., H. Locker, and D. Vlazny. 1980. Defective herpes simplex viruses. *Ann. N.Y. Acad. Sci.* 354:347-370.
11. Grafstrom, R. H., J. C. Alwine, W. L. Steinhart, C. W. Hill, and R. W. Hyman. 1975. The terminal repetition of herpes simplex virus DNA. *Virology* 67:144-157.
12. Graham, B. J., Z. Bengali, and G. F. Vande Woude. 1978. Physical map of the origin of defective DNA in herpes simplex virus type 1 DNA. *J. Virol.* 25:878-887.
13. Halliburton, I. W., R. E. Randall, R. A. Killington, and D. H. Watson. 1977. Some properties of recombinants between type 1 and type 2 herpes simplex virus. *J. Gen. Virol.* 36:471-484.
14. Hayward, G. S., R. J. Jacob, S. C. Wadsworth, and B. Roizman. 1975. Anatomy of herpes simplex virus DNA: evidence for four populations of molecules that differ in the relative orientation of their long and short components. *Proc. Natl. Acad. Sci. U.S.A.* 72:4243-4247.
15. Honess, R. W., A. Buchan, I. W. Halliburton, and D. H. Watson. 1980. Recombination and linkage between structural and regulatory genes of herpes simplex virus type 1: study of the functional organization of the genome. *J. Virol.* 34:716-742.
16. Honess, R. W., and B. Roizman. 1974. Regulation of herpesvirus macromolecular synthesis. I. Cascade regulation of the synthesis of three groups of viral proteins. *J. Virol.* 14:8-19.
17. Honess, R. W., and B. Roizman. 1975. Regulation of herpesvirus macromolecular synthesis: sequential transition of polypeptide synthesis requires functional viral polypeptides. *Proc. Natl. Acad. Sci. U.S.A.* 72:1276-1280.
18. Huang, A. S. 1973. Defective interfering viruses. *Annu. Rev. Microbiol.* 27:101-117.
19. Hyman, R. W., S. Burke, and L. Kudler. 1976. A nearby inverted repeat of the terminal sequence of herpes simplex virus DNA. *Biochem. Biophys. Res. Commun.* 68:609-615.
20. Jones, P. C., and B. Roizman. 1979. Regulation of herpesvirus macromolecular synthesis. VIII. The transcription program consists of three phases during which both extent of transcription and accumulation of RNA in the cytoplasm are regulated. *J. Virol.* 31:299-314.
21. Kaerner, H. C., I. B. Maichle, A. Ott, and C. H. Schröder. 1979. Origin of two different classes of defective HSV-1 Angelotti DNA. *Nucleic Acids Res.* 6:1467-1478.
22. Kaerner, H. C., A. Ott-Hartmann, R. Schatten, C. H. Schröder, and C. P. Gray. 1981. Amplification of a short nucleotide sequence in the repeat units of defective herpes simplex virus type 1 Angelotti DNA. *J. Virol.* 39:75-81.
23. Knipe, D. M., W. T. Ruyechan, B. Roizman, and I. W. Halliburton. 1978. Molecular genetics of herpes simplex virus: demonstration of regions of obligatory and non-obligatory identity within diploid regions of the genome by sequence replacement and insertion. *Proc. Natl. Acad. Sci. U.S.A.* 75:3896-3900.
24. Leiden, J., N. Frenkel, and F. Rapp. 1980. Identification of the herpes simplex virus DNA sequences present in six herpes simplex virus thymidine kinase-transformed mouse cell lines. *J. Virol.* 33:272-285.
25. Locker, H., and N. Frenkel. 1979. Structure and origin of defective genomes contained in serially passaged herpes simplex virus type 1 (Justin). *J. Virol.* 29:1065-1077.
26. Locker, H., and N. Frenkel. 1979. *Bam*I, *Kpn*I, and *Sal*I restriction enzyme maps of the DNAs of herpes simplex virus strains Justin and F: occurrence of heterogeneities in defined regions of the viral DNA. *J. Virol.* 32:429-441.
27. Mao, J. C., and E. E. Robishaw. 1975. Mode of inhibition of HSV DNA polymerase by PAA. *Biochemistry* 14:5475-5479.
28. Marsden, H. S., I. K. Crombie, and J. H. Subak-Sharpe. 1976. Control of protein synthesis in herpesvirus-infected cells: analysis of the polypeptides induced by wild-type and sixteen temperature sensitive mutants of strain 17. *J. Gen. Virol.* 31:347-372.
29. Marsden, H. S., N. D. Stow, V. G. Preston, M. C. Timbury, and N. M. Wilkie. 1978. Physical mapping of herpes simplex virus-induced polypeptides. *J. Virol.* 28:624-642.
30. Mocarski, E. S., and B. Roizman. 1981. Site-specific inversion sequence of the herpes simplex virus genome: domain and structural features. *Proc. Natl. Acad. Sci. U.S.A.* 78:7047-7051.
31. Morse, L. S., L. Pereira, B. Roizman, and P. A. Schaffer. 1978. Anatomy of herpes simplex virus (HSV) DNA. X. Mapping of viral genes by analysis of polypeptides and functions specified by HSV-1 $\times$ HSV-2 recombinants. *J. Virol.* 26:389-410.
32. Murray, B. K., N. Biswal, J. B. Bookout, R. E. Lanford, R. J. Courtney, and J. C. Melnick. 1975. Cyclic appearance of defective interfering particles of herpes simplex virus and the concomitant accumulation of early polypeptide VP175. *Intervirology* 5:173-184.
33. Pereira, L., M. H. Wolff, M. Fenwick, and B. Roizman. 1977. Regulation of herpesvirus macromolecular synthesis. V. Properties of α polypeptides made in HSV-1 and HSV-2 infected cells. *Virology* 77:733-749.
34. Post, L. E., S. Mackem, and B. Roizman. 1981. Regulation of α genes of herpes simplex virus: expression of chimeric genes produced by fusion of thymidylate kinase with α gene promoters. *Cell* 24:555-565.
35. Preston, C. M. 1979. Control of herpes simplex virus type 1 mRNA synthesis in cells infected with wild-type virus or the temperature-sensitive mutant *tsK*. *J. Virol.* 29:275-284.
36. Preston, C. M. 1979. Abnormal properties of an immediate early polypeptide in cells infected with herpes simplex virus type 1 mutant *tsK*. *J. Virol.* 32:357-369.
37. Preston, V. G. 1981. Fine structure mapping of herpes simplex virus type 1 temperature-sensitive mutations within the short repeat region of the genome. *J. Virol.* 39:150-161.
38. Roizman, B. 1979. The structure and isomerization of herpes simplex virus genomes. *Cell* 16:481-494.
39. Schaffer, P. A., V. C. Carter, and M. C. Timbury. 1978. Collaborative complementation study of temperature-sensitive mutants of herpes simplex virus types 1 and 2. *J. Virol.* 27:490-504.
40. Schröder, C. H., B. Stegmann, H. F. Lauppe, and H. C. Kaerner. 1975/76. An unusual defective genotype derived from herpes simplex virus strain ANG. *Intervirology* 6:270-284.
41. Schröder, C. H., and G. Urbaczka. 1978. Excess of

- interfering over infectious particles in herpes simplex virus passaged at high m.o.i. and their effect on single-cell survival. *J. Gen. Virol.* 41:493-501.
42. **Sheldrick, P., and N. Berthelot.** 1974. Inverted repetitions in the chromosome of herpes simplex virus. *Cold Spring Harbor Symp. Quant. Biol.* 39:667-678.
43. **Southern, D. M.** 1975. Detection of specific sequences among DNA fragments separated by gel electrophoresis. *J. Mol. Biol.* 98:503-517.
44. **Stegmann, B., H. Zentgraf, A. Ott, and C. H. Schröder.** 1978. Synthesis and packaging of herpes simplex virus DNA in the course of virus passages at high multiplicity. *Intervirology* 10:228-240.
45. **Stow, N. D., and N. M. Wilkie.** 1978. Physical mapping of temperature-sensitive mutants of herpes simplex virus type 1 by intertypic marker rescue. *Virology* 90:1-11.
46. **Vlazny, D. A., and N. Frenkel.** 1981. Replication of herpes simplex virus DNA: localization of replication recognition signals within defective virus genomes. *Proc. Natl. Acad. Sci. U.S.A.* 78:742-746.
47. **Vlazny, D. A., A. Kwong, and N. Frenkel.** 1982. Site specific cleavage/packaging of herpes simplex virus DNA and the selective maturation of nucleocapsids containing full length viral DNA. *Proc. Natl. Acad. Sci. U.S.A.* 79:1423-1427.
48. **Von Magnus, P.** 1954. Incomplete forms of influenza virus. *Adv. Virus Res.* 2:59-79.
49. **Wadsworth, S., G. S. Hayward, and B. Roizman.** 1976. Anatomy of herpes simplex virus DNA. V. Terminally repetitive sequences. *J. Virol.* 17:503-512.
50. **Wadsworth, S., R. J. Jacob, and B. Roizman.** 1975. Anatomy of herpes simplex virus DNA. II. Size, composition, and arrangement of inverted terminal repetitions. *J. Virol.* 15:1487-1497.
51. **Wagner, M. J., and W. C. Summers.** 1978. Structure of the joint region and the termini of the DNA of herpes simplex virus type 1. *J. Virol.* 27:374-387.
52. **Watson, R. J., and J. B. Clements.** 1978. Characterization of transcription deficient temperature-sensitive mutants of herpes simplex virus type 1. *Virology* 91:364-379.
53. **Watson, R. J., and J. B. Clements.** 1980. A herpes simplex virus type 1 function continuously required for early and late virus RNA synthesis. *Nature (London)* 285:329-330.
54. **Wilcox, K. W., A. Kohn, E. Sklyanskaya, and B. Roizman.** 1980. Herpes simplex virus phosphoproteins. I. Phosphate cycles on and off some viral polypeptides and can alter their affinity for DNA. *J. Virol.* 33:167-182.
55. **Wilkie, N. M., and R. Cortini.** 1976. Sequence arrangement in herpes simplex virus type 1 DNA: identification of terminal fragments in restriction endonuclease digests and evidence for inversions in redundant and unique sequences. *J. Virol.* 20:211-221.