Published in final edited form as:

Mol Nutr Food Res. 2008 July ; 52(7): 772–779. doi:10.1002/mnfr.200700081.

Gauging the clinical significance of P-glycoprotein-mediated herb-drug interactions: Comparative effects of St. John's wort, echinacea, clarithromycin, and rifampin on digoxin pharmacokinetics

Bill J. Gurley¹, Ashley Swain², D. Keith Williams³, Gary Barone⁴, and Sunil Kumar Battu⁵

¹University of Arkansas for Medical Sciences, College of Pharmacy, Department of Pharmaceutical Sciences, 4301 W. Markham St., Slot 522, Little Rock, AR 72205

²University of Arkansas for Medical Sciences, College of Medicine, General Clinical Research Center, 4301 W. Markham St., Little Rock, AR 72205

³University of Arkansas for Medical Sciences, Department of Biometry, 4301 W. Markham St., Little Rock, AR 72205

⁴University of Arkansas for Medical Sciences, Department of Surgery, 4301 W. Markham St., Little Rock, AR 72205

⁵University of Mississippi, School of Pharmacy, Department of Pharmaceutics, University, MS 38677

Abstract

Concomitant administration of botanical supplements with drugs that are P-glycoprotein (P-gp) substrates may produce clinically significant herb-drug interactions. This study evaluated the effects of St. John's wort and *Echinacea* on the pharmacokinetics of digoxin, a recognized P-gp substrate. Eighteen healthy volunteers were randomly assigned to receive a standardized St. John's wort (300 mg three times daily) or *Echinacea* (267 mg three times daily) supplement for 14 days, followed by a 30-day washout period. Subjects were also randomized to receive rifampin (300 mg twice daily, 7 days) and clarithromycin (500 mg twice daily, 7 days) as positive controls for P-gp induction and inhibition, respectively. Digoxin (Lanoxin® 0.25 mg) was administered orally before and after each supplementation and control period. Serial digoxin plasma concentrations were obtained over 24 hours and analyzed by chemiluminescent immunoassay. Comparisons of AUC₍₀₋₃₎, AUC₍₀₋₂₄₎, T_{1/2}, and C_{max}, were used to assess the effects of St. John's wort, *Echinacea*, rifampin, and clarithromycin on digoxin disposition. St. John's wort and rifampin both produced significant reductions ($p < 0.05$) in AUC₍₀₋₃₎, AUC₍₀₋₂₄₎, and C_{max}, while clarithromycin increased these parameters significantly ($p < 0.05$). *Echinacea* supplementation did not affect digoxin pharmacokinetics. Clinically significant P-gp-mediated herb-drug interactions are more likely to occur with St. John's wort than with *Echinacea*.

INTRODUCTION

Over the last eight years a considerable body of work has been published about herb-drug interactions, including several comprehensive reviews [1-10]. Interest in this topic is easily understood given the growing popularity of alternative medicine, the myriad botanicals

available worldwide as dietary supplements, the multitude of unique phytochemicals present in these products and the paucity of knowledge regarding their pharmacology. More importantly, almost 25% of all prescription drug users take herbal medicines concomitantly with conventional medications [11-13].

From a clinical perspective, the take home message from herb-drug interaction studies can oftentimes be confusing, especially when the results of in vitro studies utilizing human tissue or cell lines are not supported by human in vivo studies [14]. While in vitro studies provide insight as to which botanicals may affect drug pharmacokinetics, only in vivo studies can provide a definitive means for determining the clinical importance of pharmacokinetic herb-drug interactions. However, statistically significant effects observed among human in vivo herb-drug interaction studies might not always translate into clinical significance. For instance, Gurley et al noted that goldenseal supplementation produced a statistically significant increase (14%) in digoxin C_{max} values; however, when compared to the 95% increase produced by clarithromycin, the goldenseal-digoxin interaction appears clinically insignificant [15]. Other recent studies utilizing well-recognized modulators of cytochrome P-450 3A (CYP3A) and P-glycoprotein (P-gp) activity in vivo, demonstrated that milk thistle and black cohosh had no clinically significant effects on human CYP3A-mediated drug metabolism [16], or P-glycoprotein-mediated drug efflux [17]. Taken together, these examples illustrate that, when possible, in vivo effects of botanicals should be compared to appropriate controls in order to gauge the clinical relevance of herb-drug interactions.

P-glycoprotein (P-gp, ABCB1), an ATP binding cassette protein coded by the *ABCB1* gene, plays a prominent role in the disposition of many xenobiotics through its action as a drug efflux pump. Phytochemical-mediated alterations in P-gp activity may give rise to herb-drug interactions by altering drug absorption, distribution, and elimination [5]. In this report we describe the effects of St. John's wort and, for the first time in humans, *Echinacea* supplementation on the pharmacokinetics of digoxin, a putative P-gp substrate that exhibits a narrow therapeutic index. As a means of gauging the clinical relevancy of botanical-mediated P-gp interactions, we also compare supplement effects to those of clarithromycin, an inhibitor of P-gp activity [18] and rifampin, an inducer of P-gp expression [19].

MATERIALS AND METHODS

Study subjects

This study protocol was approved by the University of Arkansas for Medical Sciences Human Research Advisory Committee (Little Rock, AR) and all participants provided written informed consent before commencing the study. Eighteen young adults (9 females) (age, mean $\pm$ SD = 30 ± 5.4 years; weight, 77.5 ± 16.5 kg) participated in the study and all subjects were in good health as indicated by medical history, routine physical examination, electrocardiography, and clinical laboratory testing. All subjects were nonsmokers, ate a normal diet, were not users of botanical dietary supplements, and were not taking prescription (including oral contraceptives) or nonprescription medications. All female subjects had a negative pregnancy test at baseline. All subjects were instructed to abstain from alcohol, caffeine, fruit juices, cruciferous vegetables, and charbroiled meat throughout each two-week phase of the study. Adherence to these restrictions was further emphasized five days before digoxin administration. Subjects were also instructed to refrain from taking prescription and nonprescription medications during supplementation periods, and any medication use during this time was documented. Documentation of compliance to these restrictions was achieved through the use of a food/medication diary.

Supplements and supplementation/medication regimens

The effect of St. John's wort, *Echinacea*, rifampin and clarithromycin on digoxin oral absorption was evaluated individually on four separate occasions in each subject. This was an open-label study randomized for supplementation/medication sequence. ("Supplementation/medication" refers to either St. John's wort, *Echinacea*, rifampin, or clarithromycin.) Each supplementation phase (St. John's wort or *Echinacea*) lasted 14 days while each medication phase (rifampin or clarithromycin) was of 7 days duration. Each supplementation/medication phase was followed by a 30-day washout period. This randomly assigned sequence of supplementation/medication followed by washout was repeated until each subject had received all four products. Single lots of St. John's wort (lot # 530812, standardized extract WS 5572) and *Echinacea* (lot #A05551200) were purchased from Nature's Way Products, (Springville, UT) and Gaia Herbs, Inc. (Brevard, NC), respectively. (Both companies are recognized leaders in the botanical supplement industry for providing products of high quality and consistency.) Rifampin (Rifadin[®], Aventis Pharmaceuticals, Kansas City, MO.) and clarithromycin (Biaxin[®], Abbott Laboratories, North Chicago, IL) were utilized as positive controls for P-gp induction and inhibition, respectively. Product labels were followed regarding the recommended dosing of St. John's wort extract (300 mg, three times daily, standardized to contain 3% hyperforin); *Echinacea* extract (*E. purpurea* aerial, root, and seed parts 195 mg, and *E. augustifolia* root parts 72 mg [equivalent to 2600 mg crude herb], three times daily, standardized to contain 2.2 mg isobutylamides per capsule,); clarithromycin (500 mg, twice daily) and rifampin (300 mg, twice daily). Rifampin, however, was not administered on the day of digoxin administration [38]. Telephone and electronic mail reminders were used to facilitate compliance, while pill counts and supplementation usage records, were used to verify compliance.

Digoxin administration

Following an overnight fast, subjects reported to the University of Arkansas for Medical Sciences General Clinical Research Center for digoxin administration and blood sampling. Prior to digoxin administration, subjects were weighed and questioned about their adherence to the dietary and medication restrictions. Female subjects were administered pregnancy tests and only those with negative test results were allowed to participate. Following the placement of a 20 gauge indwelling catheter into a peripheral vein of the forearm, an oral dose of digoxin (0.25 mg, Lanoxin[®], GlaxoSmithKline, Research Triangle Park, NC) was administered with 240 ml of water. Throughout the study, digoxin doses were administered 24 hours before the start of each supplementation/medication phase (baseline) and again on the last day of each phase. Serial blood samples were obtained before and at 0.33, 0.67, 1, 1.5, 2, 3, 4, 6, 8, 12, and 24 hours after digoxin administration. Each subject's blood pressure, heart rate, and respiration rate was monitored at 1, 2, and 6 hours post digoxin administration. Four hours after digoxin administration, subjects received identical meals consisting of a turkey sandwich, potato chips, carrot sticks, and water.

Determination of digoxin serum concentrations

Digoxin serum concentrations were determined by an automated chemiluminescent immunoassay system (ACS:180 Digoxin, Chiron Diagnostics Corp., West Walpole, MA). Calibrations were performed in the range of 0.1–5.0 ng/mL. Serum concentrations greater than 5 ng/mL were diluted and reassayed. The lower limit of quantitation was 0.1 ng/mL. The interday accuracy for digoxin at 0.58, 1.77, and 3.48 ng/mL was 5.4, 3.7, and 2.9%, respectively. The interday precision for digoxin at 0.49, 0.98, and 1.97 ng/mL was 7%, 6%, and 2% respectively.

Supplement analysis

The phytochemical content of each supplement was independently analyzed for specific “marker compounds” by ChromaDex Inc. (Clearwater, FL). Quantitative determination of hyperforin, hypericin, and various flavonoids in St. John's wort was achieved using a proprietary isocratic HPLC method similar to that described by Liu et al [21]. *E. purpurea* was analyzed for various phenolic acids (e.g. chicoric acid, echinacoside, and caftaric acid) and isobutylamide content using a proprietary gradient HPLC method similar to that described by Molgaard et al. [22].

Pharmacokinetic analysis

Digoxin pharmacokinetics were determined using standard noncompartmental methods with the computer program WinNonlin (version 2.1; Pharsight, Mountain View, CA). Area under the plasma concentration time curves from zero to 24 hours ($AUC_{(0-24)}$) and zero to 3 hours ($AUC_{(0-3)}$) were determined by use of the trapezoidal rule. Rifampin, clarithromycin, and other P-gp modulators have significant effects on digoxin pharmacokinetics during the absorption phase [18,19], which was the reason for evaluating $AUC_{(0-3)}$. The terminal elimination rate constant (k_e) was calculated using the slope of the log-linear regression of the terminal elimination phase and the elimination half-life ($T_{1/2}$) was calculated as $0.693/k_e$. Peak digoxin concentrations (C_{max}) and the times when they occurred (T_{max}) were derived directly from the data.

Disintegration tests

An absence of botanical-mediated changes in CYP phenotype could stem from products exhibiting poor disintegration and/or dissolution characteristics. To address this concern, each product was subjected to disintegration testing as outlined in the United States Pharmacopeia 27 [23]. The disintegration apparatus consisted of a basket-rack assembly operated at 29-32 cycles per minute with 0.1 N HCl (37°C) as the immersion solution. One dosage unit of each supplement was placed into each of the six basket assembly tubes. The time required for the complete disintegration of six dosage forms was determined. This process was repeated with an additional six dosage units to assure accuracy. Since there are no specifications for the disintegration time of the botanical supplements used in this study, the mean of six individual dosage forms was taken as the disintegration time for that particular product. A product (soft gelatin capsule) was considered completely disintegrated if the entire residue passed through the mesh screen of the test apparatus, except for capsule shell fragments, or if the remaining soft mass exhibited no palpably firm core.

Statistical Analysis

A repeated measures ANOVA model was fit for each pharmacokinetic parameter using SAS Proc Mixed software (SAS Institute, Inc. Cary, N.C., USA). Since pre- and post-supplementation/medication pharmacokinetic parameters were determined in each subject for all four study phases, a covariance structure existed for measurements within subjects. Sex, supplement/medication, and supplement/medication-by-sex terms were estimated for each parameter using a Huynh-Feldt covariance structure fit. If supplement/medication-by-sex interaction terms for a specific parameter measure were significant at the 5% level, the focus of the post-supplementation/medication minus pre-supplementation/medication response was assessed according to sex. If the supplement/medication-by-sex interaction was not statistically significant, responses for both sexes were combined. Additionally, a power analysis was performed to estimate the ability to detect significant post- minus presupplementation/medication effects. All four models obtained at least 80% power at the 5% level of significance to detect a Cohen effect size of 1.32 to 1.71 standard deviation units [24].

RESULTS

General Experimental Observations

All eighteen subjects completed each phase of the study. Neither spontaneous reports from study participants or their responses to questions asked by study nurses regarding supplement/medication usage revealed any serious adverse events. While 2 subjects taking St. John's wort noted drowsiness, no significant side effects were reported for *Echinacea*. Nausea, indigestion, and complaints of a metallic taste were frequently noted during clarithromycin phases. Mild indigestion and reddish discoloration of the urine were common conditions reported with rifampin use. No clinically significant changes in blood pressure, heart rate, or respiratory rate were observed after digoxin administration. Examination of pill counts and food/medication diaries revealed no significant deviations from the study protocol.

Effect of Supplementation on Digoxin pharmacokinetics

The effects of clarithromycin, rifampin, St John's wort, and *Echinacea* on serum digoxin concentration versus time profiles are depicted in Figure 1. Statistically significant increases ($p < 0.05$) in digoxin $AUC_{(0-3)}$ (68%), $AUC_{(0-24)}$ (46%), elimination half-life (73%) and C_{max} (75%) were observed after 7 days of clarithromycin administration (Fig. 1; Table 1). Statistically significant reductions ($p < 0.05$) in digoxin $AUC_{(0-3)}$ (−30%), $AUC_{(0-24)}$ (−25%), and C_{max} (−38%) were noted following rifampin administration (Fig. 1; Table 1). St. John's wort also produced statistically significant reductions in digoxin $AUC_{(0-3)}$ (−28%), $AUC_{(0-24)}$ (−23%), and C_{max} (−36%) (Fig. 1; Table 1). No significant effects on digoxin disposition were observed as a result of *Echinacea* supplementation. Digoxin T_{max} (1 hour) was not significantly affected by any of the treatments. In addition, no sex-related changes in digoxin pharmacokinetics were noted for any of the supplement/medication interventions.

Phytochemical Content and Disintegration Testing

Results of phytochemical analyses and disintegration testing for St. John's wort and *Echinacea* are presented in Table 2.

DISCUSSION

Considerable evidence supports the premise that prolonged supplementation with St. John's wort can induce P-gp-mediated drug efflux, resulting in clinically significant reductions in the bioavailability of P-gp substrates like digoxin [25-27], fexofenadine [28-30], and cyclosporine [31-33]. St. John's wort appears to induce P-gp expression through the actions of hyperforin, a phytochemical unique to *Hypericum* species and potent ligand for the orphan nuclear receptor, SXR (steroid-xenobiotic-receptor; or human pregnane xenobiotic receptor, hPXR) [34,35]. When ingested, hyperforin is taken up by intestinal enterocytes and hepatocytes binding SXR in the nucleus of these cells. In turn, SXR-hyperforin complexes form heterodimers with the retinoid-X-receptor that bind to the drug response element of the *ABCB1* gene upregulating its expression [4]. Recent evidence, however, suggests that the magnitude of St. John's wort-mediated drug interactions is a function of hyperforin dose [27,33,36]. Administration of St. John's wort for at least 14 days at hyperforin doses less than 5 mg/day produced no significant changes in digoxin pharmacokinetics, whereas significant reductions in AUC and C_{max} were noted for daily doses in excess of 10 mg [27,36]. In the present study, subjects exposed to daily hyperforin doses of 24 mg for 14 days exhibited significant reductions in digoxin AUC (−25%) and C_{max} (−35%). The magnitude of these effects were strikingly similar to those reported by Johnne et al [25] and Mueller et al [27] whose subjects also ingested hyperforin doses in excess of 10 mg/day for 11-14 days.

Our findings also demonstrated that 14 days of St John's wort (24 mg/day hyperforin) reduced digoxin exposure comparable to that observed after 7 days of rifampin (600 mg/day). Rifampin, like hyperforin, is a ligand for SXR and a well-recognized inducer of P-gp [37]. Interestingly, rifampin is also a substrate for P-gp [20], as well as an inhibitor of the solute carrier organic anion transporter family, member 1B1 (SLCO1B1, also known as organic anion transporting polypeptide 1B1, OATP1B1), and its simultaneous administration with digoxin may actually increase digoxin exposure [38]. For this reason, we refrained from administering rifampin concomitantly with digoxin on the days of pharmacokinetic sampling. This practice allowed us to better gauge the clinical magnitude of P-gp induction on digoxin disposition. To our knowledge, this is the first direct comparison of St. John's wort to rifampin from an herb-drug interaction perspective.

Unlike rifampin and St. John's wort, clarithromycin markedly increased digoxin AUC, $C_{\max}$, and $T_{1/2}$. The extent of these increases were similar to that previously reported by Rengelshausen et al [18] and Gurley et al [15,17]. Accordingly, these results confirm clarithromycin's clinical utility as a positive control for P-gp inhibition. Although none of the botanical supplements tested here inhibited P-gp, future studies with other botanicals may produce statistically significant differences. By juxtaposing the effects of botanical supplementation to that of clarithromycin, or other clinically-recognized P-gp inhibitors (e.g. quinidine), a more meaningful interpretation can be obtained.

Echinacea, marketed as an "immune system booster" and alternative treatment for the common cold, ranks among the top-selling botanical supplements worldwide [39]. Due to its popularity, knowledge of its herb-drug interaction profile is much needed. In vitro studies suggest that extracts of *Echinacea* species are capable of modulating CYP activity [40-45], but no assessments of *Echinacea* on P-gp function have been reported to date. Such studies are warranted given the variety of phytochemicals present in *Echinacea*, particularly the alkamides, which are bioavailable when administered orally [46-48]. While a few clinical studies have investigated *Echinacea*'s effect on human CYP isoforms [49,50], to our knowledge this is the first report of *Echinacea*'s effect on human P-gp activity in vivo. Unlike St. John's wort, the *Echinacea* supplementation regimen used in this study produced no significant effects on digoxin disposition. This finding implies that *Echinacea* is not a potent modulator of human P-gp in vivo, and thus poses no clinically significant interaction risk with digoxin or other P-gp substrates. Previous in vivo assessments of human CYP isoforms suggest that *Echinacea* has little impact on CYP2C9, CYP2D6, CYP2E1, and CYP1A2, but may selectively modulate the activity of CYP3A4 at hepatic and intestinal sites [49,50]. When taken together, the existing clinical evidence suggests that *Echinacea*'s drug interaction potential appears to be relatively minor, however, given the inter-product variability in phytochemical content and potency among *Echinacea* supplements [51-53], these results may not extend to regimens utilizing higher dosages, longer supplementation periods, or products with improved dissolution and/or bioavailability characteristics.

In summary, botanical supplements may interact with conventional medications, but the magnitude of such interactions is not readily predictable from in vitro studies. Clinical studies offer the best means of assessing the drug interaction potential of botanical supplements. However, such studies often neglect to include appropriate benchmarks for gauging the clinical significance of any observed herb-mediated effects. Using rifampin and clarithromycin as positive controls for P-gp induction and inhibition, respectively, the effects of St. John's wort and *Echinacea* on the disposition of digoxin were evaluated. Our findings revealed that St. John's wort products containing sufficient hyperforin are equivalent to rifampin in terms of P-gp induction, whereas, *Echinacea* had no bearing on P-gp activity. These observations illustrate too the spectrum of effects that may occur during human herb-drug interaction studies and provide a means for assessing their clinical relevance. Accordingly, in vivo herb-drug

interaction studies may be better served when known modulators of drug metabolism and transport are included in their design.

Acknowledgements

This work was supported by the NIH/NIGMS under grant GM71322 and by the NIH/NCRR to the General Clinical Research Center of the University of Arkansas for Medical Sciences under grant RR14288.

Abbreviations

AUC, area under the curve; $C_{\max}$, maximum serum concentration; k_e , elimination rate constant; P-gp, P-glycoprotein; SXR, steroid xenobiotic receptor; $T_{\max}$, time of maximum serum concentration; $T_{1/2}$, elimination half-life.

REFERENCES

1. Cupp MJ. Herbal remedies: adverse effects and drug interactions. *Am. Fam. Phys* 1999;59:1239–1244.
2. Ioannides C. Pharmacokinetic interactions between herbal remedies and medicinal drugs. *Xenobiotica* 2002;32:451–478. [PubMed: 12160480]
3. Zhou S, Gao Y, Jiang W, Huang M, et al. Interactions of herbs with cytochrome P450. *Drug Metab. Rev* 2003;35:35–98. [PubMed: 12635815]
4. Zhou S, Lim LY, Chowbay B. Herbal modulation of P-glycoprotein. *Drug Metab. Rev* 2004;36:57–104. [PubMed: 15072439]
5. Mills E, Wu P, Johnston BC, Galliano K. Natural health product-drug interactions: a systematic review of clinical trials. *Ther. Drug Monit* 2005;27:549–557. [PubMed: 16175124]
6. Hu Z, Yang X, Ho PCL, Chan SY, et al. Herb-drug interactions: a literature review. *Drugs* 2005;65:1239–1282. [PubMed: 15916450]
7. John A, Roots I. Clinical drug interactions with medicinal herbs. *Evid. Based Integrative Med* 2005;2:207–228.
8. van den Bout-van den Beukel CJP, Koopmans PP, van der Ven AJAM, DeSmet PAGM, et al. Possible drug-metabolism interactions of medicinal herbs with antiretroviral agents. *Drug Metab. Rev* 2006;38:477–514. [PubMed: 16877262]
9. Ulbricht C, Basch E, Weissner W, Hackman D. An evidence-based systematic review of herb and supplement interactions by the Natural Standard Research Collaboration. *Expert Opin. Drug Saf* 2006;5:719–728. [PubMed: 16907661]
10. Lam, YWF.; Huang, S-M.; Hall, SD., editors. *Herbal Supplements—Drug Interactions: Scientific and Regulatory Perspectives*. Informa Healthcare; New York: 2006.
11. Wold RS, Lopez ST, Yau L, Butler LM, et al. Increasing trends in elderly persons' use of nonvitamin, nonmineral dietary supplements and concurrent use of medications. *J. Am. Diet. Assoc* 2005;105:54–63. [PubMed: 15635346]
12. Shahrokh LE, Lukaszuk JM, Prawitz AD. Elderly herbal supplement users less satisfied with medical care than nonusers. *J. Am. Diet. Assoc* 2005;105:1138–1140. [PubMed: 15983535]
13. Gardiner P, Graham RE, Legedza ATR, Eisenberg DM, Phillips RS. Factors associated with dietary supplement use among prescription medication users. *Arch. Intern. Med* 2006;166:1968–1974. [PubMed: 17030829]
14. Venkataramanan R, Komoroski B, Strom S. In vitro and in vivo assessment of herb drug interactions. *Life Sci* 2006;78:2105–2115. [PubMed: 16427092]
15. Gurley BJ, Swain A, Barone G, Williams DK, et al. Effect of goldenseal (*Hydrastis canadensis* and kava kava (*Piper methysticum*) supplementation on digoxin pharmacokinetics in humans. *Drug Metab. Dispos* 2007;35:240–245. [PubMed: 17079360]
16. Gurley B, Hubbard MA, Williams DK, Thaden J, et al. Assessing the clinical significance of botanical supplementation on human cytochrome P450 3A activity: comparison of a milk thistle and black cohosh product to rifampin and clarithromycin. *J. Clin Pharmacol* 2006;46:201–213. [PubMed: 16432272]

17. Gurley BJ, Barone GW, Williams DK, Carrier JC, et al. Effect of milk thistle (*Silybum marianum*) and black cohosh (*Cimicifuga racemosa*) supplementation on digoxin pharmacokinetics in humans. *Drug Metab. Dispos* 2006;34:69–74. [PubMed: 16221754]
18. Rengelshausen J, Goggelmann C, Burhenne J, Riedel K-D, et al. Contribution of increased oral bioavailability and reduced nonglomerular renal clearance of digoxin to the digoxin-clarithromycin interaction. *Br. J. Clin. Pharmacol* 2003;56:32–38. [PubMed: 12848773]
19. Greiner B, Eichelbaum M, Fritz P, Kreichgauer H-P, et al. The role of intestinal P-glycoprotein in the interaction of digoxin and rifampin. *J. Clin. Invest* 1999;104:147–153. [PubMed: 10411543]
20. Schuetz EG, Schinkel AH, Relling MV, Schuetz JD. P-glycoprotein: a major determinant of rifampin-inducible expression of cytochrome P4503A in mice and humans. *Proc. Natl. Acad. Sci* 1996;93:4001–4005. [PubMed: 8633005]
21. Liu FF, Ang CYW, Heinze TM, Rankin JD, et al. Evaluation of major active components in St. John's wort dietary supplements by high-performance liquid chromatography with photodiode array detection and electrospray mass spectrometric confirmation. *J. Chromatogr. A* 2000;888:85–92. [PubMed: 10949475]
22. Molgaard P, Johnsen S, Christensen P, Cornett C. HPLC method validated for the simultaneous analysis of cichoric acid and alkamides in *Echinacea purpurea* planta and products. *J. Agric. Food Chem* 2003;51:6922–6933. [PubMed: 14611148]
23. Anonymous. United States Pharmacopeia and National Formulary, 27th revision. 22nd ed.. United States Pharmacopeial Convention, Inc.; Rockville: 2004. Disintegration and dissolution of dietary supplements; p. 2645-2646.
24. Cohen, J. Statistical Power Analysis for the Behavioral Sciences. 2nd ed.. Lawrence Erlbaum Associates; Hillsdale: 1988.
25. Johne A, Brockmöller J, Bauer S, Maurer A, et al. Pharmacokinetic interaction of digoxin with an herbal extract from St. John's wort (*Hypericum perforatum*). *Clin. Pharmacol. Ther* 1999;66:338–345. [PubMed: 10546917]
26. Dürr D, Stieger B, Kullak-Ublick GA, Rentsch KM, et al. St. John's wort induces intestinal P-glycoprotein/MDR1 and intestinal and hepatic CYP3A4. *Clin. Pharmacol. Ther* 2000;68:598–604. [PubMed: 11180019]
27. Mueller SC, Uehleke B, Woehling H, Petzsch M, et al. Effect of St John's wort dose and preparations on the pharmacokinetics of digoxin. *Clin. Pharmacol. Ther* 2004;75:546–557. [PubMed: 15179409]
28. Wang Z, Hamman MA, Huang S-M, Lesko LJ, Hall SD. Effect of St John's wort on the pharmacokinetics of fexofenadine. *Clin. Pharmacol. Ther* 2002;71:414–420. [PubMed: 12087344]
29. Dresser GK, Schwarz UI, Wilkinson GR, Kim RB. Coordinate induction of both cytochrome P4503A and MDR1 by St. John's wort in healthy subjects. *Clin. Pharmacol. Ther* 2003;73:41–50. [PubMed: 12545142]
30. Xie R, Tan LH, Polasek EC, Hong C, et al. CYP3A and P-glycoprotein activity induction with St. John's wort in healthy volunteers from 6 ethnic populations. *J. Clin. Pharmacol* 2005;45:352–356. [PubMed: 15703370]
31. Barone GW, Gurley BJ, Ketel BL, Lightfoot ML, Abul-Ezz SR. Drug interaction between St. John's wort and cyclosporine. *Ann. Pharmacother* 2000;34:1013–1016. [PubMed: 10981246]
32. Bauer S, Störmer E, Johne A, Krüger H, et al. Alterations in cyclosporine A pharmacokinetics and metabolism during treatment with St John's wort in renal transplant patients. *Br. J. Clin. Pharmacol* 2003;55:203–211. [PubMed: 12580993]
33. Mai I, Bauer S, Perloff ES, Johne A, et al. Hyperforin content determines the magnitude of the St John's wort-cyclosporine drug interaction. *Clin. Pharmacol. Ther* 2004;76:330–340. [PubMed: 15470332]
34. Moore LB, Goodwin B, Jones SA, Wisely GB, et al. St. John's wort induces hepatic drug metabolism through activation of the pregnane X receptor. *Proc. Nat. Acad. Sci* 2000;97:7500–7502. [PubMed: 10852961]
35. Wentworth JM, Agostini M, Love J, Schwabe JW, Chatterjee VKK. St. John's wort, a herbal antidepressant, activates the steroid X receptor. *J. Endocrinol* 2000;166:R11–R16. [PubMed: 10974665]

36. Arold G, Donath F, Maurer A, Diefenbach K, et al. No relevant interaction with alprazolam, caffeine, tolbutamide, and digoxin by treatment with a low-hyperforin St John's wort extract. *Planta Med* 2005;71:331–337. [PubMed: 15856409]
37. Geick A, Eichelbaum M, Burk O. Nuclear receptor response elements mediate induction of intestinal MDR1 by rifampin. *J. Biol Chem* 2001;276:14581–14587. [PubMed: 11297522]
38. Lam JL, Shugarts SB, Okochi H, Benet LZ. Elucidating the effect of final-day dosing of rifampin in induction studies on hepatic drug disposition and metabolism. *J. Pharmacol Exp. Ther* 2006;319:864–870. [PubMed: 16905688]
39. Barnes J, Anderson LA, Gibbons S, Phillipson JD. Echinacea species (*Echinacea angustifolia* (DC.) Hell., *Echinacea pallida* (Nutt.) Nutt., *Echinacea purpurea* (L.) Moench): a review of their chemistry, pharmacology and clinical properties. *J. Pharm. Pharmacol* 2005;57:929–954. [PubMed: 16102249]
40. Budzinski JW, Foster BC, Vandenhoeck S, Arnason JT. An in vitro evaluation of human cytochrome P450 3A4 inhibition by selected commercial extracts and tinctures. *Phytomed* 2000;7:273–282.
41. Foster BC, Vandenhoeck S, Hana J, Krantis A. In vitro inhibition of human cytochrome P450-mediated metabolism of marker substrates by natural products. *Phytomed* 2003;10:334–342.
42. Strandell J, Neil A, Carlin G. An approach to the in vitro evaluation of potential for cytochrome P450 enzyme inhibition from herbals and other natural remedies. *Phytomed* 2004;11:98–104.
43. Matthias A, Gillam EMJ, Penman KG, Matovic NJ, et al. Echinacea alkamide disposition and pharmacokinetics in humans after tablet ingestion. *Chem. Biol. Interact* 2005;155:62–70. [PubMed: 15885678]
44. Yale SH, Glurich I. Analysis of the inhibitory potential of Ginkgo biloba, Echinacea purpurea, and Serenoa repens on the metabolic activity of cytochrome P450 3A4, 2D6, and 2C9. *J. Altern. Complement. Med* 2005;11:433–439. [PubMed: 15992226]
45. Hellum BH, Hu Z, Nilsen OD. The induction of CYP1A2, CYP2D6 and CYP3A4 by six trade herbal products in cultured primary human hepatocytes. *Basic Clin. Pharmacol. Toxicol* 2007;100:23–30. [PubMed: 17214607]
46. Dietz B, Heilmann J, Bauer R. Absorption of dodeca-2E,4E,8Z,10E/Z-tetraenoic acid isobutylamides after oral application of Echinacea purpurea tincture. *Planta Med* 2001;67:863–864. [PubMed: 11745026]
47. Matthias A, Addison RS, Penman KG, Dickinson RG, et al. Echinacea alkamide disposition and pharmacokinetics in humans after tablet ingestion. *Life Sci* 2005;77:2018–2029. [PubMed: 15919096]
48. Woelkart K, Koidl C, Grisold A, Gangemi JD. Bioavailability and pharmacokinetics of alkamides from the roots of Echinacea angustifolia in humans. *J. Clin. Pharmacol* 2005;45:683–689. [PubMed: 15901750]
49. Gorski JC, Huang S-M, Pinto A, Hamman MA, et al. The effect of Echinacea (*Echinacea purpurea* root) on cytochrome P450 activity in vivo. *Clin. Pharmacol. Ther* 2004;75:89–100. [PubMed: 14749695]
50. Gurley BJ, Gardner SF, Hubbard MA, Williams DK, et al. In vivo assessment of botanical supplementation on human cytochrome P450 phenotypes: Citrus aurantium, Echinacea purpurea, milk thistle, and saw palmetto. *Clin. Pharmacol. Ther* 2004;76:428–440. [PubMed: 15536458]
51. Binns SE, Livesey JF, Arnason JT, Baum BR. Phytochemical variation in Echinacea from roots and flowerheads of wild and cultivated populations. *J. Agric. Food Chem* 2002;50:3673–3687. [PubMed: 12059142]
52. Gilroy CM, Steiner JF, Byers T, Shapiro H, Georgian W. Echinacea and truth in labeling. *Arch. Intern. Med* 2003;163:699–704. [PubMed: 12639203]
53. Pellati F, Benvenuti S, Melegari M, Lasseigne T. Variability in the composition of anti-oxidant compounds in Echinacea species by HPLC. *Phytochem. Anal* 2005;16:77–85. [PubMed: 15881113]

Figure 1. Digoxin concentration-time profiles (0-6 hours) before and after each supplementation/drug phase. (A) pre- and post-clarithromycin; (B) pre- and post-*Echinacea*; (C) pre- and post-rifampin; (D) pre- and post-St. John's wort. Black squares = pre-experimental mean serum digoxin concentrations. Gray circles = post-experimental mean serum digoxin concentrations. Error bars = s.d.

Table 1Digoxin pharmacokinetic parameters before and after supplementation/drug phases (mean $\pm$ s.d.)

Supplement/Drug Phase	AUC₍₀₋₃₎ (ng·hr/mL)	AUC₍₀₋₂₄₎ (ng·hr/mL)	T_{1/2} (hours)	C_{max} (ng/mL)
Pre-Clarithromycin	2.2 $\pm$ 0.8	7.3 $\pm$ 2.0	32.5 $\pm$ 9.5	1.2 $\pm$ 0.3
Post-Clarithromycin	3.7 $\pm$ 1.3*	10.7 $\pm$ 3.4*	56.3 $\pm$ 18.6*	2.1 $\pm$ 0.9*
Pre-Rifampin	2.3 $\pm$ 0.7	8.1 $\pm$ 1.9	34.6 $\pm$ 11.5	1.3 $\pm$ 0.4
Post-Rifampin	1.6 $\pm$ 0.6*	6.1 $\pm$ 1.9*	31.3 $\pm$ 10.9	0.8 $\pm$ 0.3*
Pre-St. John's wort	2.5 $\pm$ 0.6	7.8 $\pm$ 1.6	30.1 $\pm$ 9.3	1.4 $\pm$ 0.3
Post-St. John's wort	1.8 $\pm$ 0.5*	6.0 $\pm$ 1.3*	27.7 $\pm$ 12.0	0.9 $\pm$ 0.2*
Pre-Echinacea	2.2 $\pm$ 0.5	7.3 $\pm$ 1.6	32.5 $\pm$ 8.9	1.3 $\pm$ 0.3
Post-Echinacea	2.1 $\pm$ 0.6	7.5 $\pm$ 1.8	31.5 $\pm$ 9.6	1.2 $\pm$ 0.3

AUC = area under the concentration-time curve, C_{max} = maximum serum concentration,*
= p < 0.05

Table 2

Phytochemical analysis and disintegration times for botanical dosage forms (n = 6).

Supplement (dosage form)	Compound	Content (mg/dosage unit)	Daily Dose) (mg)	Disintegration Time (minutes)
St. John's wort (coated tablet)	Hyperforin	8.0 ± 0.06	24.0	44.8 ± 1.8
	Hypericin	0.1 ± 0.001	0.3	
	Pseudohypericin	0.3 ± 0.001	0.9	
	<u>Flavonoids</u>			
	Rutin	5.6 ± 0.06	16.8	
	Hyperoside	3.6 ± 0.04	10.8	
	Isoquercetin	2.1 ± 0.03	6.3	
	Quercetin	0.8 ± 0.01	2.4	
	<u>Quercetin</u>	<u>0.8 ± 0.01</u>	<u>2.4</u>	
	Total	21.3	63.9	
Echinacea (softgel capsule)	<u>Phenolic acids</u>			6.1 ± 0.7
	Chicoric acid	3.3 ± 0.03	29.7	
	Echinacoside	1.3 ± 0.01	11.7	
	Caftaric acid	1.4 ± 0.01	12.6	
	Cynarin	0.3 ± 0.001	2.7	
	Chlorogenic acid	0.2 ± 0.001	1.8	
	<u>Isobutylamides</u>	<u>1.3 ± 0.01</u>	<u>11.7</u>	
	Total	7.8	70.2	