PREHOSPITAL CARE

The cape triage score: a new triage system South Africa. Proposal from the cape triage group

S B Gottschalk, D Wood, S DeVries, L A Wallis, S Bruijns, On behalf of the Cape Triage Group

Emerg Med J 2006;23:149–153. doi: 10.1136/emj.2005.028332

The Cape Triage Group (CTG) convened with the intention of producing a triage system for the Western Cape, and eventually South Africa. The group includes in-hospital and prehospital staff from varied backgrounds. The CTG triage protocol is termed the Cape Triage Score (CTS), and has been developed by a multi-disciplinary panel, through best available evidence and expert opinion. The CTS has been validated in several studies, and was launched across the Western Cape on 1 January 2006.

The CTG would value feedback from readers of this journal, as part of the ongoing monitoring and evaluation process.

INTRODUCTION

The health burden on emergency services in South Africa is enormous – an estimated 60 000 South Africans die of road traffic accidents and homicides each year.¹ At least 2.5 million cases of non-fatal injuries require emergency care annually – this translates to about 66 trauma presentations per 1000 population per annum,² way ahead of trauma levels in other countries. This trauma load is added to the huge burden of illness, with HIV/AIDS and Tuberculosis rife through many parts of the population.

There exists a need to properly prioritise the care of these patients, in both the prehospital and emergency unit setting. Such prioritisation is termed triage – the process of sieving or sorting patients according to need. Many triage tools exist for use in both prehospital and the emergency unit, however, such algorithms have problems preventing their adaptation in the South African setting.

In-hospital triage systems include Manchester Triage,³ the Canadian Triage Assessment Scale (CTAS),⁴ and the Australian Triage Score (ATS):⁵ each of these triage tools require extensive training to implement, making their widespread adoption in South Africa problematic. Furthermore, the time taken to triage each patient is too long for most emergency units in the South African setting, where the case-load presenting to many of our emergency units is so large that a rapid system is required.

Prehospital tools are typically generalisable to many different countries, however, they lack the sensitivity and specificity to make them safe for emergency unit use. Furthermore, some are validated only for trauma triage,^{6–8} whilst others are too detailed to be of roadside use.^{3–5}

The Cape Triage Group (CTG) was convened in Cape Town in 2004 to produce a triage system suitable for local use. The CTG falls under the auspices of the Joint Division of Emergency Medicine of the University of Cape Town (UCT) and University of Stellenbosch. The group is multidisciplinary and comprises doctors, sisters, and paramedics from the state and private sectors, ensuring expertise in all aspects of

emergency care from prehospital through to admission from the emergency unit.

The intention of the CTG was to develop a system of triage for use in the Cape.

CURRENT TRIAGE SYSTEMS IN SOUTH AFRICA

A. Prehospital

The prehospital use of triage in South Africa varies from region to region, but is generally categorised into four priorities (represented by colours). The systems in use at selected South African ambulance training colleges are presented in table 1.

Such triage typically uses instability of vital signs to differentiate high from low priority patients, following the recommendations of MacMahon.⁹ Discrepancies in triage appear when personnel of differing levels of medical experience and qualifications classify patients, as there are no clear definitions of ‘unstable’ physiology. Without objective clinical parameters, such variations in patient assessment are inevitable. To complicate matters further, classifications such as ‘stable red’ or ‘unstable yellow’ patients confuse the true severity of the patient’s clinical profile. The terms ‘stable’ and ‘unstable’ are poorly understood and fail to accurately reflect the patient’s clinical condition.

Triage must be rapid to undertake whilst being easy to use. The current system of triage fulfils both these criteria but lacks sensitivity and specificity (as well as consistency).

B. Emergency Unit

Public sector emergency units are under enormous pressure with large patient numbers, high case severity, understaffing and poor resources. GF Jooste hospital is a typical example, seeing over 4500 patients per month in the emergency unit but staffed to see around half of this number. This unit deals with the current top four epidemics in South Africa in equal proportions: HIV/AIDS, non-HIV related infectious diseases (tuberculosis), traumatic injuries, and diseases of lifestyle (hypertension, diabetes, etc.).¹⁰ Roughly 35% of these cases require immediate to urgent attention, whilst only 25% are considered non-urgent.¹¹ This leads to long waiting times and, more importantly, poor management of clinical risk. These pressures are not unique to the state sector, with some of the busier private units seeing more patients of higher severity than they are resourced to deal with. Private units (such as Milnerton Mediclinic) see over 2000 patients per month, including polytrauma and medical emergencies, with insufficient staffing to avoid the need for triage. However, no formal triage systems are in place, relying rather on the “eyeball” assessment of a nurse.

In order to maximise efficiency of the use of resources, and to minimise risk to the patient, an effective triage system with high sensitivity and specificity is required.

Table 1 Current EMS triage criteria as instructed at colleges throughout South Africa

COLLEGE	RED (P1)	YELLOW (P2)	GREEN (P3)	BLUE (DEAD)
Cape Technikon, Cape Town	Primary Survey compromised	Maintaining own Primary Survey. Injury/illness requires treatment within 60 minutes.	Injury/illness that should not compromise the Primary Survey within 60 minutes	The obviously dead
Wits Technikon, Gauteng	Primary Survey compromised	Maintaining own Primary Survey. Injury/illness requires treatment within 60 minutes.	Injury/illness that should not compromise the Primary Survey within 60 minutes	The obviously dead
DTI, Natal	Life-threatening emergencies	Non life-threatening emergencies requiring hospital treatment.	Minor injury/illness. Walking wounded	The obviously dead
Lebone Ambulance College (Pretoria)	Treatable life-threatening injuries/illness	Serious non life-threatening injuries.	Minor, easily managed injury/illness that may not require Ambulance transportation	The obviously dead
Natal Ambulance College	Life-threatening emergencies	Seriously injured patients	Moderate injuries	The obviously dead
EMS College, Cape Town	Primary Survey is compromised or an injury that will lead to permanent disability	Maintaining own Primary Survey. Injury/illness requires treatment within 60 minutes.	Injury/illness that will not compromise Primary Survey within 60 minutes	Mortal injury

METHODS

The CTG endeavoured to design and test a simple, effective triage tool. This instrument had to fulfil the requirements of being easy to use by all levels of prehospital providers, nursing staff and doctors - from the roadside through to the EU. A universally accepted and utilised objective triage system would avoid discrepancies in patient classification. Thus, for example, 'red' would mean 'red', and 'yellow' would mean 'yellow' to all, despite different levels of experience and qualifications. The CTG reviewed existing triage instruments and developed a new combined CTG triage system and scoring sheet. Each of the following factors was considered in developing the system:

- Physiological derangement
- Discriminators
 - Mechanism of injury
 - Symptom complexes
 - Discriminators
 - Senior Health-Care Personnel's Opinion
- Nomenclature

Physiological derangement

Triage is dynamic: a patient's status can change at any time and so their triage category may also change. This important ability to change priority¹²⁻¹³ is exclusive to physiological assessment. It was felt essential that the triage system contains physiological assessment. The Revised Trauma Score, based on Glasgow Coma Score (GCS), Systolic Blood Pressure and Respiratory Rate, has been successfully used to identify seriously injured trauma cases presenting to an emergency unit (systolic hypotension was found to be a particularly useful sign of serious injury).¹⁴ However, this score may not include sufficient clinical parameters to differentiate between medical cases. Previous research has focussed on developing criteria for identifying medical in-patients at risk of deterioration. Acute dyspnoea has been found to be particularly useful as a physiological marker of imminent arrest.¹⁵⁻¹⁶ The Medical Early Warning Score (MEWS) uses physiological parameters such as systolic blood

pressure, heart rate, temperature, respiratory rate and AVPU level of consciousness. MEWS has been used to successfully identify physiological deterioration of medical in-patients¹⁷ – in this study, MEWS scores of five or more were associated with increased risk of death, ICU and High Dependency Unit admission (approximately 25% combined outcome). The MEWS score is a 'simple, practical method of using routine physiological measurements to flag, from among the mass of admissions, patients who should not be ignored'.¹⁸

In 1997 the Audit Commission in the UK considered the issue of critical care outreach services to medical wards to consult on medical in-patients who were deteriorating. This led to the development of Intensive Care Outreach Services (ICORS) whereby intensive care consultation in medical wards was identified prior to the need for ICU admission. Summarising abnormal physiology into the MEWS score was found to be a particularly useful tool in this process.¹⁸ Using the MEWS as an ICORS referral tool led to decreased admission rate to ICU and shorter length of stay.¹⁹

A triage system that incorporated the MEWS score had been implemented at GF Jooste hospital in March 2004 and its substantial success initially alerted the CTG to the possibility of its use as a triage tool. The EU saw a considerable improvement in patient flow, and a significant reduction in mortality. However, the MEWS score is flawed with respect to triage in that it has a medical bias. Trauma patients (who are generally previously healthy and therefore have more physiological reserve) may have severe injuries and yet have a low MEWS score if they have stable physiology. The addition of both a mobility parameter and trauma factor were felt necessary to increase the severity score for trauma cases. This modified MEWS score was renamed the Triage Early Warning Score (TEWS). The TEWS scoring system has the following advantages:

- It requires a comprehensive assessment of the ill patient early on.
- It translates parameters that can be easily measured in both the prehospital and emergency unit setting, by even the basic trained levels of staff, to an equally easy and interpretable triage score.
- It encompasses both trauma and medical patients, in both the prehospital and emergency unit setting.

- By using this system, health care providers will be able to classify patients similarly, allowing for transparency of communication between medical staff and with more appropriate transfer of patients.

Despite the in-hospital success of the MEWS, it has not been formally validated as a triage tool. The TEWS score has been validated as part of a Masters degree,²⁰ in addition to another Masters dissertation²¹ and a PhD thesis,²² involving the prospective assessment of almost 25,000 patients in both the public and private settings. A pre-hospital study will commence in 2006.

Discriminators

a. Mechanism of Injury

Mechanism of injury scores have been shown to be highly sensitive at identifying casualties with severe trauma. However, they have also been shown to have high rates of overtriage.²³ Mechanism of injury was included but limited to high energy transfer.

b. Presentations

Triage systems often use presentations as core elements: examples of these include level of consciousness, haemorrhage and pain. We deliberately avoided long lists that would lead to long algorithms or a bulky triage tool. Although there are some specific diagnoses in the triage algorithm (e.g. burn – face/inhalation), most are presentation types such as chest pain.

c. Pain

Pain is an important part of the assessment of patients in the EU. The CTS includes an assessment of pain as part of the triage decision making process.

d. Senior health-care personnel's opinion

There is evidence that experienced health care professional can improve the triage process by adding their opinion to other parameters.²² However, this is experienced based and it would not be appropriate for more junior staff to triage by this "gut feel". In the CTG protocol, the senior health care provider involved with the patient may alter the triage coding.

Nomenclature

The CTG considered whether a colour-coded or 'Priority' (e.g. P1–P4) nomenclature was more practical. Due to the fact that colours are more readily identifiable to both medical staff and patients (considering practical points such as colour-coded stickers on folders and even coloured zones in emergency units) it was felt that a colour-based system was preferential.

Many triage systems use a 5-tiered approach (UK, Canada, Australia). Due to the problems of the 'stable red' and the 'unstable yellow' patients, a fifth category, orange, has been introduced into the South African system. The triage categories chosen by the CTG are:

- Red - resuscitation/physiologically unstable patients
- Orange - serious cases with potentially unstable physiology or potentially life/limb threatening pathology
- Yellow - 'physiologically stable' cases with reasonably serious medical or trauma problems
- Green - minor injuries/illness
- Blue - those who are clearly dead

Most of the cases referred to an emergency unit fall into the orange or yellow category (44% of all attendees at GF Jooste¹¹). The orange category essentially reduces the number of patients in the very large yellow category (38% in the

previous four colour system at GF Jooste) whilst limiting the red category (26% in previous system at GF Jooste) to resuscitation or acutely unstable cases.

RESULTS

The final CTS is a 5-colour coding system comprising the following:

- TEWS score
- Discriminators
 - Mechanism of injury
 - Presentation
 - Pain
 - Senior health care professional's discretion

The CTS is printed on posters and credit card sized keycards, which can then be used by medical personnel from the roadside through the EU (figure 1). Poster sized versions will be available in EUs to facilitate triage.

There are three versions: adult, child, and infant. The adult version is intended for patients aged over 12 years, or taller than 150 cm. The infant version is for children under three years, or less than 95 cm, and the child version is for other children (three to 12 years, 95 to 150 cm). The adult version is shown below, and the child and infant versions are available on the web.

Use of the CTS

The use of the TEWS score encourages the healthcare professional undertaking the triage to actually measure physiological parameters, a step that is worthwhile not only for the purposes of triage. This is the first step in the triage process, with the recorded physiological values being awarded a score from the TEWS sheet: these are totalled and the resulting value is read off from the first line of the triage coding chart.

It is important to emphasise that it is not necessary to "tick all the boxes" for triage of individual patients – if a patient's TEWS score make them orange then they are orange, unless there is an obvious other problem that would make them red. A current seizure, for example, will triage the patient red regardless of his TEWS score.

DISCUSSION

The CTS was rolled out across the Western Cape on 1 January 2006, following an extensive training programme. The CTS is fully supported by the provincial Department of Health. Experience at GF Jooste²¹ suggests that even inexperienced staff can learn the triage system following only a brief teaching period. It is accepted that there will be a "bedding in" period as staff become familiar with the system. There will be ongoing evaluation and assessment of the CTS, which will determine any modifications that need to be made to the system.

A request has been made to form a national Triage Group from 2006, with a view to taking the CTS nationwide.

Clearly the CTS is not intended for mass casualty situations: it is standard practice to use a simple triage method for a mass casualties situations and a more complex protocol on a day-to-day basis. Such mass casualties systems must be easy to learn and to use, fast to undertake and accurate.²⁵ Examples of such systems include the Triage Sieve²⁵ (used throughout the UK, Netherlands, Sweden, parts of India and Australia, and NATO military organisations), Careflight²⁶ (Australia) and START²⁷ (USA).

While South African EMS providers are likely to be involved in mass-casualty situations on a more frequent

TEWS	3	2	1	0	1	2	3
Mobility				Walking	With help	Stretcher/ immobile	
RR		< 9		9–14	15–20	21–29	≥ 30
HR		? 40	41–50	51–100	101–110	111–129	≥ 130
Syst BP	? 70	71–80	81–100	101–199		≥ 200	
Temp		< 35		35–38.4		≥ 38.5	
AVPU				A lert	V oice	P ain	U nresponsive

	RED	ORANGE	YELLOW	GREEN	BLUE
TEWS	> 7	6 to 7	3 to 5	0 to 2	Dead
MOI	Entrapment	Impact – high	Impact – low		
SYMPTOM					
Respiratory	Asthma – status	Asthma			
Cardiac		Chest pain			
Vascular		Haemorrhage – arterial			
AVPU	Unresponsive	Responds to <u>P</u> ain	Responds to <u>V</u> oice	<u>A</u> lert	
Neuro	Seizure – current	Seizure – post-ictal			
Psych		Psychosis/Aggression			
Ortho		Limb – threatened			
		Dislocation – major joint	Dislocation – minor joint		
		Fracture – open	Fracture – closed		
Burn	Burn – face/ inhalation	Burn > 20%	Burn – minor		
Medical	Hypoglycaemia < 2.2	Overdose/Poisoning	Abdominal pain		
GIT		Haemetemesis – fresh blood			
Obs and gynae		Pregnancy – trauma	Pregnancy – PV bleed		
ANATOMY	Trauma – airway	Trauma – head/neck torso	Trauma – limb		
		Evisceration			
PAIN		Severe	Moderate	Mild	
~~~~~Senior health-care professional's discretion~~~~~					
'trauma' = penetrating/blunt 'torso' = chest/abdo/back					

Figure 1 Cape Triage Group Triage System

basis than most ambulance services (around 10 such calls per week in the Cape Town Metropole), such events are still comparatively rare: around 0.1–0.2% of Metro Western Cape's workload (*personal communication, Dr W Smith Metro EMS, 2005*). A two-triage system approach for South Africa is still entirely appropriate, although a mass casualty system does not yet exist in the Western Cape (where the same protocol is used as is used for single patients). It is not part of the remit of the CTG to develop a mass casualty triage tool.

CONCLUSION

There is a clear need for a simple yet robust triage instrument for use in the South African medical arena. All role players, from prehospital to the emergency units, must use whatever system is implemented. The “buy in” of all key role players is therefore essential.

As with any tool, the development of this new triage instrument is a dynamic process. Whilst the science behind the CTS has been demonstrated in several studies, the CTS is

a living tool and must be monitored and evaluated. If changes are necessary, they will be made. As part of the ongoing evaluation, we wish to receive the input or as varied an audience as is possible in order to ensure that CTS evolves as a properly structured triage scheme. We value feedback from the readers of this journal (positive or otherwise) – either through the journal or direct to the authors.

CONTRIBUTION

All authors contributed equally to the article, on behalf of the Cape Triage Group. Dr L Wallis is the guarantor of this paper.

Authors' affiliations

S B Gottschalk, D Wood, S DeVries, L A Wallis, S Bruijns, Division of Emergency Medicine, University of Cape Town, Observatory, Cape Town

Funding: none

Competing interests: none declared

Correspondence to: Dr Lee A Wallis, PO Box 901, Wellington, 7654, South Africa; leewallis@bvr.co.za

Accepted for publication 25 August 2005

REFERENCES

- 1 **Matzopoulos R**, Seedat M, Cassim M. A profile of fatal injuries in South Africa: Fourth annual report of the National Injury Mortality Surveillance System (NIMSS). 2002, South African Medical Research Council.
- 2 **Matzopoulos RG**, Prinsloo MR, Bopape JL, *et al.* Estimating the South African trauma caseload as a basis for injury surveillance. South African Medical Research Council, 1999.
- 3 **Manchester Triage Group**. Emergency Triage. Manchester: BMJ Publishing Group, 1997.
- 4 **Canadian Association of Emergency Physicians**. Implementation Guidelines for the Canadian Emergency Department Triage and Acuity Scale (CTAS). 1998. Canadian Association of Emergency Physicians. <http://www.caep.ca/002.policies/002-02.ctas.htm> (accessed Nov 2004).
- 5 **Australian College of Emergency Medicine**. Guidelines for the Implementation of the Australian Triage Score. 98. Australian College of Emergency Medicine. <http://www.acem.org.au/open/documents/triageguide.htm> 6 (accessed Nov 2004).
- 6 **Champion HR**, Sacco WJ, Copes WS, *et al.* A Revision of the Trauma Score. *J Trauma* 1989;**29**:623–9.
- 7 **Koehler JJ**, Baer LJ, Malafa SA, *et al.* Pre-hospital index: a scoring system for field triage of trauma victims. *Ann Emerg Med* 1986;**15**:178–82.
- 8 **Henry MC**, Alicandro JM, Hollander JE, *et al.* Evaluation of American College of Surgeons trauma triage criteria in a suburban and rural setting. *Am J Emerg Med* 1996;**14**:124–9.
- 9 **MacMahon AG**. Sorting out triage in urban disasters. *SAMJ* 1985;**67**:555–6.
- 10 **Bradshaw D**, Groenewald P, Laubscher R, *et al.* Initial Burden of Disease estimate for South Africa 2000. South African Medical Research Council.
- 11 **GF Jooste Emergency Unit**. GF Jooste Emergency Unit Audit 2004.
- 12 **Martin TE**. Resolving the casualty evacuation conflict. *Injury* 1993;**24**:514–6.
- 13 **Vayer JS**, Ten Eyck RP, Cowan ML. New concepts in triage. *Ann Emerg Med* 1986;**15**:927–30.
- 14 **Champion HR**, Sacco WJ, Copes WS, *et al.* A revision of the Trauma Score. *J Trauma* 1989;**29**:623–9.
- 15 **Sax FL**, Charlson ME. Medical patients at high risk for catastrophic deterioration. *Crit Care Med* 1987;**15**:510–5.
- 16 **Subbe CP**, Kruger M, Rutherford P, *et al.* Validation of a modified Early Warning Score in medical admissions. *QJM* 2001;**94**:521–6.
- 17 **Subbe CP**, Davies RG, Williams E, *et al.* Effect of introducing the Modified Early Warning score on clinical outcomes, cardio-pulmonary arrests and intensive care utilisation in acute medical admissions. *Anaesthesia* 2003;**58**:797–802.
- 18 **Goldhill DR**. The critically ill: following your MEWS. *QJM* 2001;**94**:507–10.
- 19 **Pittard AJ**. Out of our reach? Assessing the impact of introducing a critical care outreach service. *Anaesthesia* 2003;**58**:882–5.
- 20 **Gottschalk S**. Evaluation of the Modified Early Warning Score as a triage tool in the Western Cape emergency healthcare services. In progress. University of Cape Town.
- 21 **Baxt WG**, Berry CC, Epperson MD, *et al.* The failure of pre-hospital trauma prediction rules to classify trauma patients accurately. *Ann Emerg Med* 1989;**18**:1–8.
- 22 **Fries GR**, McCalla G, Levitt MA, *et al.* A prospective comparison of paramedic judgement and the trauma triage rule in the pre-hospital setting. *Ann Emerg Med* 1994;**24**:885–9.
- 23 **Bruijns S**. A prospective evaluation of the feasibility of using enrolled nursing auxiliaries to triage patients in the emergency unit of an urban public hospital in South Africa. In Progress, University of Cape Town.
- 24 **Twomey, M**. Determination of a paediatric version of the Cape Triage Score. In progress. University of Cape Town.
- 25 **Advanced Life Support Group**. Major Incident Management and Medical Support: the practical approach. London: BMJ Publishing Group, 2002.
- 26 **Nocera A**, Garner A. An Australian mass casualty incident triage system for the future based upon triage mistakes of the past: the Homebush Triage Standard. *Aust N Z J Surg* 1999;**69**:603–8.
- 27 **Benson M**, Koenig KL, Schultz CH. Disaster triage: START, then SAVE—a new method of dynamic triage for victims of a catastrophic earthquake. *Pre-hospital Disaster Med* 1996;**11**:117–24.