PostScript

LETTERS

A nation divided

Corfield *et al*¹ demonstrate improvements in door-to-needle time with a relatively simple organisational change. Other examples of innovation can be found across the country.² Driving these changes in England and Wales is the National Service Framework which sets a standard of 60 minutes from calling for professional help until the initiation of thrombolysis.³ The Myocardial Infarction National Audit Project (MINAP) demonstrates the success of this approach.⁴

The only current standard which applies in Scotland is that set by Quality Improvement Scotland (QIS). This is a door-to-needle target of 30 minutes for the delivery of thrombolysis in the context of an acute ST elevation myocardial infarction.⁵ There is no standard set for call-to-needle and furthermore there is no national audit framework to monitor progress. In our own area although we achieve and exceed the current QIS target with a median door-to-needle time of 24 minutes for patients presenting to the Accident and Emergency department, if the National Framework target for call-to-needle is applied we would fail to achieve the standard by 27% (median call-to-needle time 76 minutes). We can see no reason why the prehospital phase of the treatment of acute myocardial infarction has been ignored by QIS. Without this as an audit standard, many departments simply do not collect the data, and the opportunity for local organisational change is lost.

If national targets are to affect the performance of the National Health Service, they need to be applied to the entire United Kingdom.

R J Cook, J Ronald, D K Pedley, W G Morrison
Accident and Emergency Department, Ninewells Hospital and Medical School, Dundee, Scotland

Correspondence to: R J Cook, Accident and Emergency Department, Ninewells Hospital and Medical School, Dundee, Scotland; ronald.j.cook@tuht.scot.nhs.uk

doi: 10.1136/emj.2005.024448

References

- 1 Corfield AR, Graham CA, Adams JN, *et al*. Emergency department thrombolysis improves door to needle times. *Emerg Med J* 2004;21:676-80.
- 2 Department of Health. Review of early thrombolysis. London: Department of Health, 2003, <http://www.dh.gov.uk/assetRoot/04/06/82/51/04068251.pdf>
- 3 Department of Health. National service framework for coronary heart disease. London: HMSO, 2000, 28-35. <http://www.dh.gov.uk/assetRoot/04/05/75/25/04057525.pdf>
- 4 Royal College of Physicians of London. Myocardial infarction national audit project. London: Royal College of Physicians of London, 2004, www.rcplondon.ac.uk/pubs/books/minap/index.htm (accessed 2 November 2005).
- 5 Clinical Standards Board for Scotland. Secondary prevention following ST elevation myocardial infarction. Edinburgh: Clinical Standards, 2000, 29-32. <http://www.show.scot.nhs.uk/sehd/publications/cdtt/cdtt.pdf>

NICE head injury guidelines—expensive? Yes, but what are the alternatives?

I read with some surprise Dr Leaman's article on the impact of the National Institute for Clinical Excellence (NICE) guidelines on a district general hospital.¹ I am truly confused at the level of anxiety this topic is causing in the UK. In most other developed countries, the use of skull x rays has long been abandoned in favour of the selected use of computed tomography (CT) scans in patients with head injuries. Although I appreciate the increased resource implications inherent in the implementation of these guidelines, I really do not see a viable alternative, apart from reliance on an outdated mode of investigation (skull x ray) in what should be a first world setting. In our paediatric department, we started relying on a modified version of the American Academy of Pediatrics guidelines in 2001. The use of skull x rays has fallen dramatically, although there has only been a modest rise in the use of CT scans in patients with minor head injuries.

I do agree with the Dr Leaman's point regarding out of hours scans, but we really have no firm evidence on which to base our assumption that patients requiring a scan can always wait until the morning. On the other hand, implementation of the NICE guidelines, along with careful audit will provide us with such data, at which time we can safely modify the guidelines to suit local practice. It is time that the UK falls into line with modern practice regarding the management of head injuries, even if this means an increase in the use of resources.

I Sammy

University of the West Indies, St Augustine, Trinidad and Tobago; psam@tstt.net.tt

doi: 10.1136/emj.2005.024430

Reference

- 1 Leaman AM. The NICE guidelines for the management of head injury: the view from a district hospital. *Emerg Med J* 2004;21:400.

ATLS: there are alternatives

I was heartened by the courageous but constructive critique of the Advanced Trauma Life Support (ATLS) course by Driscoll and Wardrope.¹ It promises an end to my lonely position as ATLS sceptic, which began in 1999 when I suggested that, although ATLS was "the greatest reformation in trauma care" in the late twentieth century,² it was "an American solution in a British context" (your readers may have noticed the subsequent regrettable trend in which car occupants who are one minute squabbling vigorously about responsibility for a collision are the next abruptly strait-jacketed in yellow plastic). I fretted that the Course was based on a dubious notion that trauma is a "surgical" disease, that it had become the New Dogma, and was suspiciously clubby ("I'm an ATLS instructor and you're not"). But my real concerns were that:

- The ATLS course arguably represents a major drain on scarce human and educational resources (How many days do consultants in emergency medicine spend away from their departments on these courses? How much is the NHS spending on the course fees?).
- ATLS only addresses the initial part of trauma care and is not a panacea (injury prevention and rehabilitation are at least as important).
- While evidence based medicine is something to which we all now aspire, the massive changes in the ATLS manual since its inception are a worrying illustration of the dearth of serious science underpinning what ATLS promotes.

I share Driscoll and Wardrope's disappointment at the continuing exclusion of emergency physicians from the development of ATLS. But happily, five years after I quit the NHS, the Royal College of Surgeons in Ireland has addressed my concerns with a set of clinical guidelines³ that I believe recognise the realities of trauma care in these islands.

May I invite your readers to peruse a document³ which is by no means perfect, but which reflects their daily experience, along with that of their anaesthetic, surgical, and other colleagues? The sheer number of consultants and trainees in emergency medicine who contributed to this important document may reassure those of your readers who feel a little excluded by the ATLS "elite" that their contribution to trauma care is in fact incalculable. I hope it may also offer some encouragement to those who believe that there are other options in trauma care on this side of the Atlantic, and that emergency medicine will show the way.

C Luke

Cork University Hospital, Cork, Ireland; chrisluke@eircom.net

doi: 10.1136/emj.2005.025080

References

- 1 Driscoll P, Wardrope J. ATLS: past, present, and future. *Emerg Med J* 2005;22:2-3.
- 2 Luke LC. ATLS: has the end become the means? *CPD Anaesthesia* 1999;1:94-6.
- 3 Clinical Guidelines Committee, Royal College of Surgeons in Ireland. Initial Management of the Severely Injured Patient, Dublin, November 2003. Available at: www.rcsi.ie/documents/SEVERELYINJUREDGUIDELINES.pdf (accessed 2 November 2005).

Auditory and visual prompts during cardiopulmonary resuscitation in the emergency department

We read with interest Williamson *et al*'s¹ study on the use of audio prompts in automatic external defibrillators to improve cardiopulmonary resuscitation (CPR) in untrained and trained lay subjects. We report a study from an emergency department (ED) using trained ED staff (doctors and nurses) and paramedics. Through our own

departmental video audit we have observed great variability in the rate of external cardiac compressions (ECC) (median 140 compressions per minute (cpm); range 100–180 cpm). This rapid rate of ECC is of concern since it is faster than the recommended 100 cpm.² Studies suggest that rescuers may fatigue after only one minute of CPR³ and there is an associated decline in the quality of chest compressions with time.⁴

We undertook two separate studies to test specifically (a) whether a piece of music influenced the rate of ECC when compared with nothing and a metronome and (b) whether using a specially adapted clock influenced the rate of ECC when compared with a metronome. In both studies participants were videotaped while performing CPR on an intubated resuscitation manikin for three minutes in the resuscitation bay of the ED. The rate of ECC was subsequently analysed by using video playback.

In the first study 50 participants were randomised in blocks of five to listen to a minute of silence followed by a second minute of a randomly selected piece of music, and then a final minute of a recording of a metronome set at a 100 beats per minute (bpm). Participants were not told that the metronome was set at 100 bpm. Three pieces of music had a rate of 100 bpm and one much less than 100 bpm and another much faster than 100 bpm. In the second study 43 participants were randomised (in blocks of 10) to two groups: metronome group or "resuscitation clock" group. The metronome group listened to a minute of silence followed by two minutes of a recording of a metronome set to 100 bpm. The resuscitation clock group listened to a minute of silence and were then asked to look at the resuscitation clock and use it to help time the rate of ECC. The resuscitation clock was essentially a normal clock with the numerals removed and the figures 25, 50, 75, and 100 placed at the 3, 6, 9, and 12 o'clock positions, respectively.

In the first study the rate of ECC was not statistically different between either the silence, music, or metronome groups (Freidman's test $\chi^2 = 3.6$, $p = 0.16$; range 54–156, median 116 cpm). Those participants who had received formal CPR training within three months of taking part in the study did not have a statistically different rate of ECC compared with those without recent training (Mann-Whitney U test = 139, $p = 0.14$). In the second study the metronome group achieved the target compression rate with a mean of 99.8 (95% CI 99.2 to 100.4), compared with clock group mean 106.9 (95% CI 101.6 to 112.2) and the control (silence) group mean 114.6 (95% CI 109.6 to 119.3).

We concluded that without the use of auditory prompts trained ED personnel are poor at performing ECC at 100 bpm even if recent formal CPR training has taken place. The use of a simple auditory prompt can help trained personnel perform ECC at the correct rate.

J France, S Wilson, N Whitton
Great Western Hospital, Swindon, UK

Correspondence to: J France, Great Western Hospital, Swindon, UK; jfrance@btopenworld.com

doi: 10.1136/emj.2005.025395

References

- 1 Williamson LJ, Larsen PD, Tzeng TC, et al. Effects of automatic external defibrillator audio prompts

on cardiopulmonary resuscitation performance. *Emerg Med J* 2005;**22**:140–3.

- 2 Lockey A, Nolan J. Cardiopulmonary resuscitation in adults. *BMJ* 2001;**323**:819–20.
- 3 Javier Ochoa F, Ramalle-Gomara E, Lisa V, et al. The effect of rescuer fatigue on the quality of chest compressions. *Resuscitation* 1998;**37**:149–52.
- 4 Ashton A, McCluskey A, Gwinnett CL, et al. Effect of rescuer fatigue on performance of continuous external chest compressions over 3 min. *Resuscitation* 2002;**55**:151–5.

Steroids for meningitis in the Emergency Department?

The European Dexamethasone study for adults with proven bacterial meningitis¹ was a multi-centred double blind randomised controlled trial of 300 patients, comparing 10 mg intravenous dexamethasone six hourly for four days with placebo. The study demonstrated an absolute risk reduction (ARR) in an unfavourable outcome of 26% (95% CI 8–44%) in those patients with pneumococcal meningitis who received dexamethasone. The results were less impressive in all patients with meningitis: ARR of 8% (95% CI 1–15%) for all types of bacteria and an ARR in death of 10% (95% CI 1–19%) in those receiving dexamethasone. The complications were surprisingly low.

We performed a retrospective survey of patients in our hospital that had a discharge diagnosis of bacterial meningitis. We conducted the survey over an eight year period (1996–2004), reviewing 35 case notes. We found that our patient characteristics were very similar to those in the study (see table 1).

Our survey also revealed of the 28 cases where documentation of time of antibiotics was recorded, 19 patients (68%) had antibiotics given prior to LP. In the study all patients had antibiotic therapy withheld until after LP had been performed (within 30 minutes of patient arrival) and the CSF analysis known. In our survey the mean door-antibiotic time for all patients was 216 minutes and the medical team performed all LPs.

It would appear to us that where Emergency Physicians (EP) give antibiotics to patients with suspected meningitis before an LP is performed, they are actually preventing patients with pneumococcal meningitis from getting the significant benefit that steroids would confer. Such EPs should therefore consider performing LPs themselves in patients with suspected bacterial meningitis and withholding antibiotics until CSF analysis is performed. The infrastructure to allow this to happen in a very timely manner, regardless of the grade of doctor seeing the patient, would have to be put in place.

Alternatively, EPs should consider giving 10 mg dexamethasone concurrently with antibiotics to patients for whom they have a high clinical suspicion of bacterial meningitis and only continuing to give subsequent doses to those with proven pneumococcal meningitis. This is very important when patients present with a depressed conscious level and suspected meningitis as these patients have the most to gain from steroids and are unlikely to have an LP for a significant length of time.

In conjunction with the Medical team in our hospital, a proforma has been constructed to help doctors select appropriate patients where steroids are to be given prior to LP. We are currently reviewing the final diagnoses of those patients who are given dexamethasone.

Are you implementing the evidence or ignoring it?

G L A Cumberbatch, L J Clark, B Neate

Emergency Department, Poole Hospital NHS Trust, Poole, UK

Correspondence to: Dr G L A Cumberbatch, Emergency Department, Poole Hospital, Longfleet Rd, Poole, Dorset BH15 2JB, UK; Gary.Cumberbatch@poole.nhs.uk

Accepted for publication 23 April 2005

Reference

- 1 De Gans J, Van de Beek D, for the European Dexamethasone in Adult Bacterial Meningitis Study Investigators. Dexamethasone in Adults with Bacterial Meningitis. *N Engl J Med* 2002;**347**:1549–56.

Handheld echocardiography – core skill for emergency physicians or another expensive course?

At the end of last year, I attended a course entitled Hand Held Echo for Critical Care, which was organised by Houman Ashrafi and Richard Bogle at Ealing Hospital. As a trainee in Emergency Medicine and Intensive Care I felt there was a good chance that I would pick up something useful for one or other of my chosen specialties. In fact, the course was hugely rewarding and in my mind I felt the concept of handheld echo within emergency medicine had jumped from a novelty to an essential requirement.

The success of this course (places are now booked up well in advance) is in part due to the enthusiasm of the organisers. Although they are both experienced cardiologists, they are convinced that echo can be successfully

Table 1

	Our Study	De Gans Study	Statistical test
Mean Age	50.3 SD 21.1	45 SD 19 (average of two arms of trial)	$p = 0.12$ Independent samples t-test
Male Sex Ratio	54%	56%	$p = 0.98$ Chi-squared test for assoc. with Yates Connection
Glasgow Coma Score <8	21%	16%	$p = 0.69$
Pneumococcus	40%	36%	$p = 0.79$
Meningococcus	20%	32%	$p = 0.19$
Negative CSF culture	40%	30%	$p = 0.4$
Mortality	11%	15% placebo group	$p = 0.83$