PostScript

LETTERS

Response to Thyagarajan *et al*'s audit: "An audit of corneal abrasion management following the introduction of local guidelines in an accident and emergency department."

We read with interest the audit by Thyagarajan *et al*¹ of corneal abrasion management after the introduction of local guidelines in an accident and emergency department, and believe it is worth specifically highlighting why it is so important to ask about contact lens use in the history as illustrated by the following case:

A 30-year-old contact lens wearer recently presented with a 2-day history of a painful, red left eye. Topical chloramphenicol had been started 24 h before for a corneal abrasion. The best corrected visual acuity of her left eye was counting fingers. Examination showed a severe central corneal ulcer. The patient was admitted for intensive topical ofloxacin. Culture results showed *Pseudomonas aeruginosa*, resistant to chloramphenicol. Four months later, the best corrected visual acuity of her left eye was 6/18 due to residual stromal scarring.

Contact lens use is the most important risk factor for the development of microbial keratitis,^{2,3} a potentially sight-threatening condition if correct treatment is not started promptly. In a 15-year retrospective review of antibiotic resistance in bacterial keratitis in London, >30% of the isolates were resistant to chloramphenicol,⁴ the usual prophylactic treatment in the UK for corneal abrasions. The incidence of chloramphenicol resistance increased markedly during the study period, and is attributable to an increase in the proportion of *Pseudomonas* species isolates,⁴ which are the most frequently found organisms in contact lens-related corneal ulcers.⁵

As the early signs of microbial keratitis can be subtle, we recommend that all contact lens wearers who have a red, sore eye with a corneal epithelial defect should be referred to the eye unit for urgent assessment.

A Day

Eye Unit, Chelsea and Westminster Hospital, London, UK

G Carp

The Western Eye Hospital, London, UK

Correspondence to: A Day, Eye Unit, Chelsea and Westminster Hospital, 369 Fulham Road, London SW10 9NH, UK; alex@acday.co.uk

doi: 10.1136/emj.2006.039784

Accepted for publication 29 June 2006

Competing interests: None declared.

References

- 1 Thyagarajan SK, Sharma V, Austin S, *et al*. An audit of corneal abrasion management following the introduction of local guidelines in an accident and emergency department. *Emerg Med J* 2006;23:526-9.

- 2 Schaefer F, Bruttin O, Zografos L, *et al*. Bacterial keratitis: a prospective clinical and microbiological study. *Br J Ophthalmol* 2001;85:842-7.
- 3 Bourcier T, Thomas F, Borderie V, *et al*. Bacterial keratitis: predisposing factors, clinical and microbiological review of 300 cases. *Br J Ophthalmol* 2003;87:834-8.
- 4 Tuft SJ, Matheson M. In vitro antibiotic resistance in bacterial keratitis in London. *Br J Ophthalmol* 2000;84:687-91.
- 5 Cheng KH, Leung SL, Hoekman HW, *et al*. Incidence of contact-lens-associated microbial keratitis and its related morbidity. *Lancet* 1999;354:181-5.

Intranasal diamorphine integrated care pathway for paediatric analgesia in the accident and emergency department

Intranasal diamorphine (IND) is a safe and effective analgesic in children attending A&E with clinically suspected limb fractures.¹ To introduce IND for paediatric analgesia in our unit, we developed an integrated care pathway (ICP)² and subjected its use to an audit cycle.³

To formulate the INDICP, various information sources were interrogated for evidence about IND use in paediatric analgesia (Medline database, article bibliographies, published guidelines). Only one multi-centre randomised controlled trial was identified, in which IND was found to have a faster onset of pain relief and better tolerability and acceptability than intramuscular morphine.¹ IND use was mentioned in guidelines from the British Association for Emergency Medicine (<http://www.emergencymed.org.uk/BAEM/>) and the Scottish Intercollegiate Guideline Network (www.sign.ac.uk/guidelines/fulltext/58/evidence.html). The final INDICP incorporated inclusion and exclusion criteria, tasks to be performed, and their sequence and timescales.² These included recording of weight, baseline clinical observations, pain score, verbal consent to treatment, how to deliver IND, monitoring of oxygen saturation, and further pain scores at 10 and 20 min. Appropriate staff training was undertaken before the INDICP was introduced.

An audit of the first 6 months of INDICP use was undertaken.³ This identified good safety (no patient required naloxone) and acceptability of IND use, but poor recording of verbal consent (essential for the unlicensed use of a licensed medicine), clinical observations, and pain scores. Accordingly a simple, one page, clinical proforma, which could be included in the patient notes, was developed to capture these data. Re-audit by means of a rapid-cycle sampling method³ found verbal consent, clinical observations, and pain scores were recorded in 50%, 66%, and 50% of cases, respectively.

To sustain this improvement in patient care, further cycles of staff instruction and re-audit will be required. We believe that the ICP and the clinical proforma are generally applicable for paediatric analgesia in other A&E departments.

K K Gahir, P A Ransom

A&E Department, Princess Royal Hospital, Lewes Road, Haywards Heath, West Sussex, RH16 4EX, UK

Correspondence to: K K Gahir, East Sussex Hospitals NHS Trust, Eastbourne District General Hospital, Kings Drive, Eastbourne, BN21 2UD, UK; Kalvinder.Gahir@esht.nhs.uk

INDICP and clinical proforma are available from the first author on request

doi: 10.1136/emj.2006.034637

Competing interests: None declared.

References

- 1 Kendall JM, Reeves BC, Latter VS, on behalf of the Nasal Diamorphine Trial Group. Multicentre randomised controlled trial of nasal diamorphine for analgesia in children and teenagers with clinical fractures. *BMJ* 2001;332:261-5.
- 2 Campbell H, Hatchkiss R, Bradshaw N, *et al*. Integrated care pathways. *BMJ* 1998;316:133-7.
- 3 National Institute for Clinical Excellence. *Principles for best practice in clinical audit*. Oxford: Radcliffe Medical Press, 2002.

BOOK REVIEW

ABC of sexually transmitted infections

Edited by Michael W Adler, Frances Cowan, Patrick French, Helen Mitchell, John Richens. London: Published by BMJ Publishing Group, 2004, £22.49 (paperback).

For many reasons, this is not a book to read on the train or bus on the way to work, which is a shame as it is an excellent text. I have always felt a little uncomfortable managing potential sexually transmitted diseases (STDs) in the emergency department, not only because I felt I did not know that much about them but also because of the feeling that perhaps we should be doing more in the emergency department, thereby obviating the need for referral to genitourinary medicine clinics which are under considerable strain at the moment.

Having read this concise and easily accessible text, I am now more confident in the assessment of STDs and also much happier that the place to investigate and treat is ideally in a dedicated facility. The chapters on the importance, control and prevention of STDs reinforce this and give an insight into the current scale of what is an increasingly important public health problem. The role of an STD service in this regard is beyond what the average emergency department can deliver, and these patients clearly need referral.

There are several aspects to this text which I think are particularly good. Firstly, the authors have chosen not just to recommend treatment regimens based solely on their own practice, but also to give recommendations from a wide number of clearly identifiable sources. Secondly, the overlap of other conditions that may present as a possible sexually transmitted infection is particularly valuable for those working in emergency medicine.

So, my advice would be to get a hold of this book, but if you intend to read it on the train then perhaps put it inside the ABC of something else or you may get a few odd looks from your fellow passengers.

S Carley