Clinical, molecular, and genotype–phenotype correlation studies from 25 cases of oral–facial–digital syndrome type 1: a French and Belgian collaborative study

C Thauvin-Robinet, M Cossée, V Cormier-Daire, L Van Maldergem, A Toutain, Y Alembik, E Bieth, V Layet, P Parent, A David, A Goldenberg, G Mortier, D Héron, P Sagot, A M Bouvier, F Huet, V Cusin, A Donzel, D Devys, J R Teyssier, L Faivre

J Med Genet 2006;43:54–61. doi: 10.1136/jmg.2004.027672

Oral–facial–digital syndrome type 1 (OFD1) is characterised by an X linked dominant mode of inheritance with lethality in males. Clinical features include facial dysmorphism with oral, tooth, and distal abnormalities, polycystic kidney disease, and central nervous system malformations. Large interfamilial and intrafamilial clinical variability has been widely reported, and 18 distinct mutations have been previously reported within *OFD1*. A French and Belgian collaborative study collected 25 cases from 16 families. *OFD1* was analysed using direct sequencing and phenotype–genotype correlation was performed using χ^2 test. X inactivation studies were performed on blood lymphocytes. In 11 families, 11 novel mutations, including nine frameshift, one nonsense, and one missense mutation were identified, which spanned nine different exons. A combination of our results with previously reported cases showed that the majority of mutations (65.5%) was located in exons 3, 8, 9, 13, and 16. There was phenotype–genotype correlation between (a) polycystic kidney disease and splice mutations; (b) mental retardation and mutations located in exons 3, 8, 9, 13, and 16; and (c) tooth abnormalities and mutations located in coiled coil domains. Comparing the phenotype of the families with a pathogenic mutation to families with absence of *OFD1* mutation, polycystic kidneys and short stature were significantly more frequent in the group with no *OFD1* mutation, whereas lingual hamartomas were significantly more frequent in the group with *OFD1* mutation. Finally, an X inactivation study showed non-random X inactivation in a third of the samples. Differential X inactivation between mothers and daughters in two families with high intrafamilial variability was of particular interest. Slight phenotype–genotype correlations were established, and X inactivation study showed that skewed X inactivation could be partially involved in the pathogenesis of intrafamilial clinical variability.

Oral–facial–digital syndrome type 1 (OFD1; OMIM #311200) or Papillon–Léage–Psaume syndrome belongs to the heterogeneous group of OFD syndromes. Incidence is 1/50 000 live births. Clinical features include hypertelorism, broad nasal bridge, hypoplasia of the nasal alae, small median cleft or pseudocleft of the upper lip (45%), hyperplastic buccal frenulae, cleft palate (80%), bifid or lobulated tongue (30–45%), lingual hamartomas (70%), hypodontia, especially frequent on low lateral incisors, and facial asymmetry. Abnormalities of the fingers (45%) are more frequent than toes (25%), and include brachydactyly, syndactyly, clinodactyly, and more rarely, polydactyly.^{1 2}

Polycystic kidney disease is common (50%).^{3 4} Central nervous system malformations are variable (40%), including corpus callosum agenesis, arachnoid cysts, cerebellar abnormalities, hydrocephalus, and porencephaly, and may be associated with mental retardation (40%).^{5 6} Characteristic radiological features are represented by irregular pattern of radiolucency and/or spicule-like formation in metacarpals and phalanges on hand x rays. Clinical variability is high and diagnosis may be difficult in minor cases. In addition, the phenotype overlaps with those reported in the other 11 forms of OFD syndromes.^{7–10} OFD1 is distinguished by its X linked dominant mode of inheritance with lethality in males and common association with polycystic kidney disease.^{3 4 11 12} The *OFD1* gene, also named *Cxorf5* (Xp22.2–22.3), contains 23 coding exons.^{13–16} *OFD1* is ubiquitously expressed in adult tissues, and in vitro studies have shown that it seems to escape X inactivation.¹⁶ Eighteen mutations have been reported in the literature.^{15 17–19} This French and Belgian collaborative study collected 25 observations of patients with OFD1 from 16 families. Clinical, molecular, and genotype–phenotype correlation studies are presented.

PATIENTS AND METHODS

Patients

Patients' clinical data and samples were collected from genetic clinics through a French and Belgian collaboration. Female patients with oral, facial, and digital features, associated or not with a family history compatible with X linked dominant mode of inheritance, were included. Patients with an OFD syndrome other than type 1, or with familial history ruling out dominant X linked pattern, were excluded. Family 1 was referred for prenatal diagnosis of porencephaly, and clinical details have been reported in detail previously.²⁰ The other 15 families were ascertained through the association of facial dysmorphism, oral features, and/or cerebral, renal, and digital malformations.

Methods

Mutation analysis

Blood samples and informed consents were collected from 24/25 patients. DNA was extracted from peripheral EDTA, according to previously reported methods.²¹ In the female fetus reported in family 1, DNA was also extracted from fetal blood, frozen liver, lung, fibroblasts, thymus, and paraffin block for the tongue. Mutations in *OFD1* were screened by using primer sequences covering all 23 exons.¹⁵ PCR amplifications of the samples were run through 30 cycles

Abbreviations: CNS, central nervous system; OFD1, oral–facial–digital syndrome type 1; PKD, polycystic kidney disease

Figure 1 Pedigrees of the familial cases reported in this article.

consisting of 40 seconds at 94°C (denaturation), 40 seconds at 50°C or 55°C (annealing), and 40 seconds at 72°C (extension), with a final extension step of 10 minutes. Sequencing of PCR products was performed using the ABI Prism7 BigDye Terminator Cycle Sequence kit (version 1.1; Applied Biosystems, Foster City, CA, USA) in both ranges, and analysed using an ABI Prism7 3100 Genetic Analyzer, according to the manufacturer's instructions. Identification of a mutation in a proband was followed by direct sequencing of the affected exon in each family member. When a missense mutation was found, 100 French control DNAs were sequenced for the region of interest in order to exclude a polymorphism.

Linkage studies

In familial cases in which no pathogenic mutations were found, DNA was amplified by PCR using fluorescent primers flanking microsatellite polymorphisms of the region of interest. The primers were chosen according to their location on the Marshfield map, as previously used for linkage studies.²² Primer sequences were obtained from the National Center for Biotechnology Information database. The selected markers used were DXS1224 and DXS7108, located telomeric to *OFD1* (0.59 and 16.56 cM respectively) and DXS7105 and DXS8018, located centromeric to *OFD1* (11.19 and 12.64 cM respectively). Microsatellite analysis was undertaken using the ABI 3700 Genescan/Genotyper

Figure 2 Clinical pictures of the reported patients showing oral abnormalities. Left side, patients with mutations identified in *OFD1*: (A) Individual IV-3, family 1 with buccal frenulae and lingual hamartoma. (B, C) Proband of family 4 with cleft and lobulated tongue, buccal frenulae, and surgery for cleft palate. (D) Proband of family 5 with cleft and lobulated tongue, buccal frenulae, and lingual hamartoma. (E) Proband of family 6 with cleft and lobulated tongue, buccal frenulae, lingual hamartoma, narrow palate, tooth abnormalities, and surgery for cleft palate. (F, G) Twin 1 and 2, family 8 with buccal frenulae, hypodontia, and abnormally shaped teeth. (H) Proband of family 9 with lobulated tongue, hypodontia, and abnormally shaped teeth of the inferior maxilla. Right side, patients without mutations in *OFD1*: (A) Individual IV-2, family 12 with cleft palate. (B) Individual III-2, family 12 with lingual hamartoma. (C) Individual III-3, family 13 with cleft and lobulated tongue and dental misposition. (D) Proband of family 16 with cleft and lobulated tongue. Permission from the patients was obtained for publication of these photographs.

Figure 3 Clinical pictures of the reported patients showing distal abnormalities. Left side, patients with mutations identified in *OFD1*: (A) Individual IV-3 in family 1 with postaxial polydactyly. (B) Individual III-3 in family 1 with brachydactyly predominant on the fourth and clinodactyly on the third finger. (C) Proband of family 4 with severe brachydactyly predominant on the fourth finger and clinodactyly. (D, E) Proband of family 5 at 7 months and at 11 ½ years of age, with brachydactyly on fingers III-IV and clinodactyly. (F) Proband of family 6 with severe generalised brachydactyly predominant on finger I and clinodactyly. (G) Proband of family 11 with brachydactyly. Right side, patients without mutations in *OFD1*: (A) Individual III-2, family 12 with brachydactyly, clinodactyly, and low set and deviated left thumb. (B) Individual III-3, family 13 with brachydactyly, clinodactyly, and syndactyly. (C) Individual II-2, family 13 with severe brachydactyly. (D) Proband of family 14 with severe brachydactyly predominant on finger IV. (E) Proband of family 16 with brachydactyly and syndactyly. Permission from the patients was obtained for publication of these photographs.

system, according to the manufacturer's instructions. Genotypes were recorded manually.

Genotype-phenotype correlation

The mutations found in the present study were added to those previously described in the literature. A correlation between some clinical features (hypertelorism, cleft lip and palate, buccal frenulae, lingual hamartomas, tooth abnormalities, brachydactyly, syndactyly, polydactyly, polycystic kidney disease, corpus callosum agenesis, mental retardation), mutation types (frameshift, splice mutations), and mutation locations (coiled coil domains, exons 3, 8, 9, 13, and 16) was studied using a χ^2 test. The study was not performed when the sample was insufficient to be contributive for genotype-phenotype correlation ($n < 5$). The correlation for the same clinical features between the group of patients with a pathogenic *OFD1* mutation and patients with absence of *OFD1* mutation was also studied using a χ^2 test.

X chromosome inactivation

The X chromosome inactivation pattern in lymphocytes was studied in 24 female cases. The FRAXA locus and the human androgen receptor assay were used with experimental conditions modified from the literature.^{23–26} Primer sequences and experimental conditions are available on request.

RESULTS

Clinical studies

From 16 families, 25 affected female cases were collected, average age 22 years. Eight cases were sporadic. Pedigrees of the familial cases are presented in fig 1. Clinical features of the whole series are summarised in tables 1–3. Facial

dysmorphism was a consistent feature, including buccal frenulae (76%), hypertelorism (67%), hypoplasia of the nasal alae (58%), cleft palate or lip (56%), lingual hamartomas (52%), lobulated or cleft tongue (48%), downslanting palpebral fissures (50%), and pseudocleft of the upper lip (21%). Available clinical pictures are presented in fig 2. Tooth abnormalities (44%) included hypodontia and malposition. Distal abnormalities with predominant brachydactyly (64%) were more frequent in hands (64%) than in feet (28%) (fig 3). Associated malformations included mainly corpus callosum agenesis (71%) and polycystic kidney (44%). Mild to moderate mental retardation was present in 11/23 cases (48%), but information was not available in 2/25 cases (fetus and 3 week old newborn). Additional cerebral features, including arachnoid cysts (4/14), cerebellar hypoplasia (3/14), hydrocephalus (2/14), and porencephaly (1/14) were more rarely associated with *OFD1* syndrome. Short stature was also described (4/25). Hand x rays were performed in only four cases, and showed spicule-like formations in metacarpals and/or phalanges in three of those.

Mutation analysis

Eleven novel mutations in *OFD1* were found in eight sporadic and three familial cases. Mutations located in nine different exons included eight deletions and one insertion (nine frameshift), one stop mutation, and one point mutation (table 4). In family 6, a G138S missense mutation was found, and was therefore tested in 100 controls. In families 12 and 15, the same base exchange was identified 295 bases proximal to the start codon, and was also found in an unaffected relative male, indicating a polymorphism. No mutation was found in five patients, including three familial

Table 1 Clinical features of the reported patients with *OFD1* mutation

Families	1			2			3	4	5	6	7	8	9	10	11
Case	IV 3	IV 2	III 3	II 2	IV 7	III 9						Twin 1	Twin 2		
Sex	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F
Age*	f	5	25	51	4	NA	15 days	39	11.5	11	3 weeks	11.5	11.5	36	28
Familial history	+	+	+	+	+	+	–	A	–	–	–	A	A	–	+
Short stature	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Frontal bossing	–	–	–	NA	–	–	+	–	–	+	–	–	–	–	–
Flat face	–	–	–	NA	–	–	+	–	–	–	–	–	–	–	–
Hypertelorism	+	–	–	NA	–	–	+	–	–	+	+	+	+	+	–
Epicanthus	+	–	–	NA	–	–	–	+	–	+	–	–	–	–	–
Telecanthus	–	–	–	NA	–	–	–	–	–	–	–	–	–	–	–
Down-slanting palpebral fissures	+	–	–	NA	+	–	–	+	+	+	–	–	–	+	+
Hypoplasia of the nasal alae	+	–	–	NA	+	+	+	+	+	+	–	–	–	–	+
Low set ears	+	–	–	NA	–	–	+	–	–	+	–	–	–	–	–
Pseudocleft of the upper lip	–	+	+	NA	–	+	–	–	–	+	+	–	–	–	–
Cleft lip	–	–	–	NA	+	–	+	–	–	–	–	–	–	+	+
Microretrognathia	–	–	–	NA	–	–	+	–	–	+	+	–	–	–	+
Abnormal hairs	–	–	–	NA	–	–	–	+	+	–	–	–	–	–	+
Cleft tongue	–	–	–	–	–	–	–	+	+	+	–	–	–	+	+
Lobulated tongue	–	–	–	–	–	+	+	+	+	+	–	–	–	+	+
Buccal frenulae	+	+	–	–	+	+	+	+	+	+	+	+	+	–	+
Lingual hamartomas	+	+	–	+	+	–	+	+	+	+	+	–	+	–	+
Cleft palate	–	–	–	–	–	–	+	+	–	+	+	–	+	–	–
Tooth abnormalities	–	–	–	–	–	–	–	+	+	+	+	+	+	+	–
Cerebellar hypoplasia	+	–	NA	NA	–	NA	–	–	–	–	–	NA	NA	–	NA
Corpus callosum agenesis	+	–	NA	NA	+	NA	–	+	+	+	+	NA	NA	+	NA
Hydrocephalus	–	–	NA	NA	–	NA	–	–	+	+	–	NA	NA	–	NA
Porencephaly	+	–	NA	NA	–	NA	–	–	–	–	–	NA	NA	–	NA
Arachnoid cysts	+	–	NA	NA	+	NA	–	+	+	–	–	NA	NA	–	NA
Hands															
Brachydactyly	–	–	+	–	–	–	+	+	+	+	+	+	–	+	+
Clinodactyly	–	–	+	+	–	+	–	+	+	+	+	–	–	–	–
Syndactyly	+	–	–	–	–	–	–	+	–	–	–	–	–	+	–
Polydactyly															
Preaxial	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Postaxial	+	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Feet															
Brachydactyly	–	–	+	–	–	–	+	–	+	–	–	–	–	–	+
Clinodactyly	–	–	–	–	–	–	–	–	–	+	–	–	–	–	+
Syndactyly	+	–	–	–	–	–	–	+	–	–	–	–	–	–	–
Polydactyly															
Preaxial	+	–	–	–	–	–	–	–	–	–	–	–	–	+	–
Postaxial	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Polycystic kidney disease	–	NA	–	NA	–	NA	NA	+	–	–	–	–	–	+	NA
Mental retardation	NA	–	–	–	+	–	–	+	+	+	NA	–	–	+	+

*In years unless otherwise stated. NA, not available; A, adopted; f, fetus; F, female.

cases (families 12, 13, and 15) and two sporadic cases (cases 14 and 16).

Linkage studies

In family 12, microsatellite polymorphism analysis of the Xp22.2–22.3 region showed compatible linkage with the *OFD1* region, as all affected individuals had inherited the same chromosome X.

Genotype–phenotype correlation

Renal cysts were significantly more frequently associated with splice mutations compared with other mutations ($\chi^2 = 0.012$). Mental retardation was significantly more frequently associated with mutations located in exons 3, 8, 9, 13, and 16 compared with mutations located in other exons ($\chi^2 = 0.04$). Tooth abnormalities were significantly more frequently associated with mutations located in coiled coil domains compared with mutations with other locations ($\chi^2 = 0.007$). No correlation was found between *OFD1* clinical features and frameshift compared with other mutations.

When comparing the phenotype of the 11 families with a pathogenic mutation with that of the five families with absence of mutation within *OFD1* (table 3), we showed that there was no significant difference between the presence or

absence of hypertelorism, cleft lip and palate, buccal frenulae, tooth abnormalities, digital abnormalities, corpus callosum agenesis, or mental retardation in both groups. However, short stature was only seen in the group of patients with no *OFD1* mutation, and polycystic kidney disease was significantly more frequent in the group with absence of *OFD1* mutation compared with the group with a pathogenic mutation ($\chi^2 = 0.035$). Conversely, lingual hamartomas were less common in the group with absence of mutation compared with the group with a pathogenic *OFD1* mutation ($\chi^2 = 0.041$).

X chromosome inactivation

X chromosome inactivation results in lymphocytes could not be produced in 1/24 patients because of homozygosity at both loci (patient II-2, family 1). Analyses showed non-random inactivation in 7/23 patients (30%) from six families (table 5), three of which were familial cases. Determination of the parental origin of the skewed allele was possible in two families. In family 1, the skewed X chromosome was of maternal origin in patient III-3 (mother) and of paternal origin in patient IV-3 (fetus). X chromosome inactivation studies in fetal tissues were not successful because of poor quality of the DNA. In family 15, the skewed X chromosome was of paternal origin in patient III-2.

Table 2 Clinical features of the reported patients without OFD1 mutation

Families	12	13	14	15	16
Case	III 2	IV 2	III 3	II 2	III 2
Sex	F	F	F	F	F
Age*	61	NA	29	65	20
					17 months
Familial history	+	+	+	+	+
Short stature	—	—	+	+	—
Frontal bossing	—	—	—	—	—
Flat face	—	—	—	+	—
Hypertelorism	+	+	+	+	+
Epicanthus	—	—	+	+	—
Telecanthus	—	—	+	+	—
Down-slanting palpebral fissures	+	+	+	+	—
Hypoplasia of the nasal alae	+	+	+	+	+
Low set ears	—	—	+	—	—
Pseudocleft of the upper lip	—	—	—	—	—
Cleft lip	—	—	+	+	+
Microretrognathia	—	—	—	—	—
Abnormal hairs	—	+	—	—	—
Cleft tongue	+	—	+	—	+
Lobulated tongue	—	—	+	+	+
Buccal frenula	—	—	+	+	+
Lingual hamartomas	+	—	—	+	—
Cleft palate	+	+	+	+	—
Tooth abnormalities	—	—	+	+	+
Cerebellar hypoplasia	+	—	NA	NA	NA
Corpus callosum agenesis	+	+	NA	NA	NA
Hydrocephalus	—	—	NA	NA	NA
Porencephaly	—	—	NA	NA	NA
Arachnoid cysts	—	—	NA	NA	NA
Hands					
Brachydactyly	+	—	+	+	+
Clinodactyly	+	—	+	—	—
Syndactyly	—	—	+	—	+
Polydactyly					
Preaxial	—	—	—	—	—
Postaxial	—	—	—	—	—
Feet					
Brachydactyly	—	—	+	+	—
Clinodactyly	—	—	—	+	—
Syndactyly	—	—	+	+	—
Polydactyly					
Preaxial	—	—	—	—	—
Postaxial	—	—	—	—	—
Polycystic kidney disease	+	NA	+	+	NA
Mental retardation	+	+	+	+	—

*In years unless otherwise stated. NA, not available; F, female.

DISCUSSION

We report a large French and Belgian clinical and molecular study of 25 patients from 16 families presenting with suspected OFD1 syndrome. Mutations in *OFD1* were identified in 16/25 patients.

The frequencies of the major clinical features were similar to those reported in the literature. Facial dysmorphism was a consistent feature, and included hypertelorism (67%), buccal frenulae (76%), hypoplasia of the nasal alae (58%), lingual hamartomas (52%), and cleft palate or/and lip (56%). Tooth abnormalities (44%), including hypodontia and malposition were not always seen in association with cleft palate and lip; 2/11 patients without cleft palate or/and lip presented noticeable tooth abnormalities. Considering distal abnormalities, brachydactyly involving predominantly the second, third, and fourth fingers was characteristic (64%).

Recently, the increasing use of renal ultrasound scan revealed that polycystic kidney disease (PKD) is commonly associated with OFD1 syndrome.³⁻⁴ Our study confirms these data, as PKD was present in 7/16 cases (44%). However, 8/9 patients were too young (from fetal period to 11.5 years of age) to draw definite conclusions about the absence of kidney involvement.

The central nervous system (CNS) may also be involved in as many as 40% of the cases.³ In our group, corpus callosum

agenesis (10/14 cases) was more frequent than previously reported. However, results may be biased, as cerebral magnetic resonance imaging was not systematically performed, particularly when mental retardation was absent. Moderate to mild mental retardation was frequent (48%), but could not always be correlated with CNS malformations. Spicule-like formations in metacarpals and/or phalanges were present in 3/4 cases. This previously reported sign is frequent and specific to OFD1 syndrome but hand x rays were rarely performed (4/25 cases).

The majority of clinical features overlaps with those reported in the other OFD syndromes.²⁷ Clinical classification remains imperfect. For example, OFD7 syndrome has been previously reported in only one family with renal involvement and mother to daughter transmission.²⁸ Later, the birth of a new affected female baby with corpus callosum agenesis in the third generation suggested OFD1 syndrome. Subsequent linkage studies at the *OFD1* locus showed compatible linkage despite a non-significant lod score (1.7).²² The authors suggested that OFD7 should not be a distinct clinicopathological entity. OFD8 syndrome (OMIM #311200), characterised by polydactyly and tibial dysplasia, is described as a recessive X linked heritable disease, but this type could also be allelic with OFD1.²⁹

Table 3 Clinical features of the reported patients with and without OFD1 mutation

Features	Mutations		
	Cases with	Cases without	Total
Familial history	7/13	14/22	14/22
Short stature	0/16	4/25	4/25
Frontal bossing	2/15	0/9	2/24
Flat face	1/15	1/9	2/24
Hypertelorism	8/15	8/9	16/24
Epicanthus	3/15	2/9	5/24
Telecanthus	0/15	2/9	2/24
Down-slanting palpebral fissures	7/15	5/9	12/24
Hypoplasia of the nasal alae	8/15	6/9	14/24
Low set ears	3/15	1/9	4/24
Pseudocleft of the upper lip	5/15	0/9	5/24
Cleft lip	4/15	3/9	7/24
Microretrognathia	4/15	0/9	4/24
Abnormal hairs	3/15	1/9	4/24
Cleft tongue	5/16	3/9	7/25
Lobulated tongue	7/16	4/9	11/25
Buccal frenula	13/16	6/9	19/25
Lingual hamartomas	11/16	2/9	13/25
Cleft palate	5/16	5/9	10/25
Tooth abnormalities	8/16	3/9	11/25
Cerebellar hypoplasia	2/10	1/4	3/14
Corpus callosum agenesis	8/10	2/4	10/14
Hydrocephalus	2/10	0/4	2/14
Porencephaly	1/10	0/4	1/14
Arachnoid cysts	4/10	0/4	4/14
Hands			
Brachydactyly	10/16	6/9	16/25
Clinodactyly	8/16	2/9	10/25
Syndactyly	3/16	2/9	5/25
Polydactyly			
Preaxial	1/16	0/9	1/25
Postaxial	1/16	0/9	1/25
Feet			
Brachydactyly	4/16	3/9	7/25
Clinodactyly	2/16	1/9	3/25
Syndactyly	2/16	2/9	4/25
Polydactyly			
Preaxial	2/16	0/9	2/25
Postaxial	0/16	0/9	0/25
Polycystic kidney disease	2/10	5/6	7/16
Mental retardation	6/14	5/9	11/23

Direct sequencing of *OFD1* in 16 families revealed 11 different mutations. In 5/16 families, no mutation was found. The detection level of 67% in our series is higher than previously reported (50%).²² Surprisingly, the detection rate was higher in sporadic (80%) than in familial cases (50%), although compatible with dominant X linked inheritance (families 13 and 15), with the presence of spicule-like formation metacarpals and/or phalanges in family 15. Similarly, linkage was compatible with the *OFD1* locus in family 12, associated with PKD and corpus callosum agenesis. In the two sporadic cases without identified mutation (cases 14 and 16), the presence of PKD strongly suggests OFD1 syndrome. In such cases, complementary molecular studies would be necessary to detect intronic mutations, mutations within the promoter, or large rearrangements such as deletions, as recently described in a recent report.³⁰ When comparing the phenotype of the families with or without a pathogenic *OFD1* mutation, we found that the presence of polycystic kidneys and short stature and the absence of lingual hamartomas suggested the absence of an *OFD1* mutation, although such data could not be useful at an individual level.

There have been 18 *OFD1* mutations reported in the literature; we identified a further 29 in this study, including 15 deletions, four insertions, two nonsense, five missense,

Table 4 *OFD1* mutations identified in our series

Family	Mutation	Exon	Effect
1	2349delC	17	Frameshift
2	1821delG	16	Frameshift
3	615delAGAAAT	7	Frameshift
4	1360delCTTA	13	Frameshift
5	T431A	6	L144X
6	G412A	4	G138S
7	675delC	8	Frameshift
8	1587delA	15	Frameshift
9	895insGA	9	Frameshift
10	712delT	8	Frameshift
11	919delG	9	Frameshift

and three splice mutations.^{15 17–19} No mutation was found in exons 18–23. The majority of mutations occurred in exons 3, 8, 9, 13, and 16 (19/29 cases, 66%), suggesting that these exons may represent regions for mutational hotspots. This hypothesis has previously been postulated for exons 3, 13, and 16.¹⁸

Mutational analysis of *OFD1* may be of interest for diagnosis and genetic counselling, especially in sporadic cases. Indeed, in family 2, family history is both compatible with dominant X linked and autosomal recessive mode of inheritance (OFD2 syndrome) with pseudodominance because of familial consanguinity. Identification of an out of frame deletion in exon 16 confirms OFD1 syndrome and therefore a risk of transmission to daughters of 50%. Another example is given by family 5. The mother presented with isolated bifid uvula, and identification of a nonsense mutation (L144X) only in the proband confirmed a de novo mutation and a consequent low risk of recurrence in other children, limited to the risk of gonadal mosaicism.³¹

When analysing the phenotype–genotype correlations, mental retardation and cleft lip/palate were reported more frequently in association with mutations in exons 3, 8, 9, 13, and 16 compared with mutations in other exons. PKD was

Table 5 X chromosome inactivation studies in our series

Family	Case	Mean % inactivation
1	IV 3	20/80
	III 3	17/83
	II 2	–
2	IV 7	46/54
	III 9	20/80
		42/58
3		15/85
4		45/55
5		15/85
6		47/53
7		29/71
8	J1	30/70
9		40/60
10		40/60
11		44/56
12	V 3	40/60
	IV 2	34/66
	III 2	47/53
13	III 3	NA
	II 2	26/74
		29/71
14		18/82
15	III 2	37/63
16	II 1	18/82

Significant results in **bold**.

more frequently associated with splice mutations compared with other mutations, and tooth abnormalities more frequently associated with mutations in coiled coil domains compared with other domains. Unfortunately, no correlation between CNS abnormalities and mutations within the LisH motif could be tested because of the small size of the sample.

Skewed X inactivation was found in 7/23 cases (30%) from six families, a higher rate than in the general population (10%).³² Skewed X inactivation was found in four cases from three families: in two severe cases, the paternal allele was inactivated, and in a moderate case, the maternal allele was inactivated. In family 2, the mother presented a moderate phenotype with skewed inactivation of unknown origin (parental origin could not be identified because of de novo mutation and unavailability of parental DNA samples), whereas her daughter has a severe phenotype with random inactivation. These results suggest that non-random X inactivation may in part explain intrafamilial clinical variability in these three families. In the three sporadic cases with non-random X inactivation, mental retardation and PKD were present in 2/3 cases, also suggesting that non-random X inactivation could explain in part the severe phenotype in sporadic cases. Additional studies in other tissues and in other patients would be of interest.

In conclusion, we report here the identification of 11 new mutations in *OFD1* in 16 French and Belgian families who presented with OFD1 syndrome. Slight phenotype-genotype correlations were identified, and the X inactivation study showed that skewed X inactivation could be partially involved in the pathogenesis of intrafamilial clinical variability. These results should be confirmed on larger samples.

ELECTRONIC DATABASE INFORMATION

Online Mendelian Inheritance in Man (OMIM), <http://www.nih.nlm.nih.gov/omim/>. National Center for Biotechnology Information database, <http://www.ncbi.nlm.nih.gov/>. Marshfield map, <http://www.marshmed.org/genetics/>.

ACKNOWLEDGEMENTS

This work was supported by a grant from the University Hospital of Dijon (Appel d'Offre Interne 2002) and by the Conseil Régional de Bourgogne. We thank the Centre d'Investigation Clinique et d'Epidémiologie Clinique (CIC-EC) for its help in the statistical analysis and J P Ramot for photographic assistance.

Authors' affiliations

C Thauvin-Robinet, Centre de Génétique, Hôpital d'Enfants, CHU Dijon, France

M Cossée, D Devys, Laboratoire de Diagnostic Génétique, CHRU Strasbourg, France

V Cormier-Daire, A Goldenberg, Département de Génétique, Hôpital Necker-Enfants Malades, Paris, France

L V Maldergem, Institut de Pathologie et de Génétique, Lovreval, Belgique

A Toutain, Service de Génétique, CHU Tours, France

Y Alembik, Service de Génétique Médicale, Hôpital Hautepierre, CHU Strasbourg, France

E Bieth, Laboratoire de Génétique, CHU Purpan, Toulouse, France

V Layet, Unité de Cytogénétique et Génétique médicale, CH Le Havre, France

P Parent, Département de Pédiatrie et de Génétique Médicale, CHU Brest, France

A David, Service de Génétique Médicale, CHU Nantes, France

G Mortier, Centre de Génétique Médicale, Ghent, Belgique

D Héron, Département de Génétique Cytogénétique et Embryologie, Hôpital La Pitié-Salpêtrière, Paris, France

P Sagot, Clinique Gynécologique et Obstétrique, Maternité du Bocage, CHU Dijon, France

A M Bouvier, Centre d'Investigation Clinique et d'Epidémiologie Clinique, Faculté de Médecine, Dijon, France

F Huet, Service de Pédiatrie 1, Hôpital d'Enfants, CHU Dijon, France

V Cusin, A Donzel, J R Teyssier, Laboratoire de Génétique Moléculaire, Hôpital du Bocage, CHU Dijon, France

Competing interests: none declared

Correspondence to: Dr C Thauvin-Robinet, Centre de Génétique, Hôpital d'Enfants, 10 Bd maréchal de Lattre de Tassigny, 21034 Dijon cedex, France; christel.thauvin@chu-dijon.fr

Received 30 September 2004

Revised 3 December 2004

Accepted 7 December 2004

REFERENCES

- 1 **Anneren G**, Arvidson B, Gustavson KH, Jorulf H, Carlsson G Oro-facio-digital syndromes I and II: radiological methods for diagnosis and the clinical variations. *Clin Genet* 1984;**26**:178–86.
- 2 **Salinas CF**, Pai GS, Vera CL, Milutinovic J, Hagerty R, Cooper JD, Cagna DR. Variability of expression of the orofaciadigital syndrome type I in black females: six cases. *Am J Med Genet* 1991;**38**:574–82.
- 3 **Donnai D**, Kerzin-Storarr L, Harris R. Familial orofaciadigital syndrome type I presenting as adult polycystic kidney disease. *J Med Genet* 1987;**24**:84–7.
- 4 **Scolari F**, Valzorio B, Carli O, Vizzardi V, Costantino E, Grazioli L, Bondioni MP, Savoldi S, Maiorca R. Oral-facial-digital syndrome type I: an unusual cause of hereditary cystic kidney disease. *Nephrol Dial Transplant* 1997;**12**:1247–50.
- 5 **Odent S**, Le Marec B, Toutain A, et al. Central nervous system malformations and early end-stage renal disease in oro-facio-digital syndrome type I: a review. *Am J Med Genet* 1998;**75**:389–94.
- 6 **Wood BP**, Young LW, Townes PL. Cerebral abnormalities in the oral-facial-digital syndrome. *Pediatr Radiol* 1975;**3**:130–6.
- 7 **Toriello HV**. Oral-facial-digital syndromes 1992. *Clin Dysmorphol* 1993;**2**:95–105.
- 8 **Figuera LE**, Rivas F, Cantu JM. Oral-facial-digital syndrome with fibular aplasia: a new variant. *Clin Genet* 1993;**4**:190–2.
- 9 **Gabrielli O**, Ficcacanti A, Fabrizio G, Perri P, Mercuri A, Coppa GV, Giorgi P. Child with oral, facial, digital and skeletal anomalies and psychomotor delay: a new OFDS form? *Am J Med Genet* 1994;**53**:290–3.
- 10 **Moran-Barroso V**, Valdes Flores M, Garcia-Cavazos R, Kofman-Alfaro S, Saavedra-Ontiveros D. Oral-facial-digital (OFD) syndrome with associated features: a new syndrome or genetic heterogeneity and variability? *Clin Dysmorphol* 1998;**7**:55–7.
- 11 **Goodship J**, Platt J, Smith R, Burn J. A male with type I orofaciadigital syndrome. *J Med Genet* 1991;**28**:691–4.
- 12 **Gillertot Y**, Heimann M, Fournieu C, Verellen-Dumoulin C, Van Maldergem L. Oral-facial-digital syndrome type I in a newborn male. *Am J Med Genet* 1993;**46**:335–8.
- 13 **Feather SA**, Woolf AS, Donnai D, Malcolm S, Winter RM. The oral-facial-digital syndrome type 1 (OFD 1), a cause of polycystic kidney disease and associated malformations, maps to Xp22.2-Xp22.3. *Hum Mol Genet* 1997;**6**:1163–7.
- 14 **Gedeon AK**, Oley C, Nelson J, Turner G, Mulley JC. Gene localization for oral-facial-digital syndrome type 1 (OFD1: MIM 311200) proximal to DXS85. *Am J Med Genet* 1999;**82**:352–4.
- 15 **Ferrante MI**, Giorgio G, Feather SA, Bulfone A, Wright V, Ghiani M, Selicorni A, Gammato L, Scolari F, Woolf AS, Sylvie O, Bernard L, Malcolm S, Winter R, Ballabio A, Franco B. Identification of the gene for oral-facial-digital type I syndrome. *Am J Hum Genet* 2001;**68**:569–76.
- 16 **de Conciliis L**, Marchitelli A, Wapenaar MC, Borsani G, Giglio S, Mariani M, Consalez GG, Zuffardi O, Franco B, Ballabio A, Banfi S. Characterisation of Cxorf5 (71–7A), a novel human cDNA mapping to Xp22 and encoding a protein containing coiled coil alpha helical domains. *Genomics* 1998;**51**:243–50.
- 17 **Stoll C**, Sauvage P. Long-term follow-up of a girl with oro-facio-digital syndrome type I due to a mutation in the OFD1 gene. *Ann Genet* 2002;**45**:59–62.
- 18 **Rakkolainen A**, Ala-Mello S, Kristo P, Orpana A, Jarvela I. Four novel mutations in the OFD1 (Cxorf5) gene in Finnish patients with oral-facial-digital syndrome I. *J Med Genet* 2002;**39**:292–6.
- 19 **Romio L**, Wright V, Price K, Winyard PJ, Donnai D, Porteous ME, Franco B, Giorgio G, Malcolm S, Woolf AS, Feather SA. OFD1, the gene mutated in oral-facial-digital syndrome type 1, is expressed in the metanephros and in human embryonic renal mesenchymal cells. *J Am Soc Nephrol* 2003;**14**:680–9.
- 20 **Thauvin-Robinet C**, Rousseau T, Durand C, Laurent N, Maingueneau C, Faivre L, Sagot P, Nivelon-Chevallier A. Familial orofaciadigital syndrome type I revealed by ultrasound prenatal diagnosis of porencephaly. *Prenat Diagn* 2001;**21**:466–70.
- 21 **Miller SA**, Dykes DD, Polesky HF. A simple salting out procedure for extracting DNA from nucleated cells. *Nucleic Acids Res* 1988;**16**:1215.
- 22 **Nowaczyk MJ**, Zeeman S, Kam A, Taylor SA, Carter RF, Whelan DT. Oral-facial-digital syndrome VII is oral-facial-digital syndrome I: a clarification. *Am J Med Genet* 2003;**123A**:179–82.
- 23 **Carrel L**, Willard HF. An assay for X inactivation based on differential methylation at the fragile X locus, FMR1. *Am J Med Genet* 1996;**64**:27–30.

- 24 Fu YH, Kuhl DP, Pizzuti A, Pieretti M, Sutcliffe JS, Richards S, Verkerk AJ, Holden JJ, Fenwick RG Jr, Warren ST. Variation of the CGG repeat at the fragile X site results in genetic instability: resolution of the Sherman paradox. *Cell* 1991;**67**:1047–58.
- 25 Martinez F, Tomas M, Millan JM, Fernandez A, Palau F, Prieto F. Genetic localisation of mental retardation with spastic diplegia to the pericentromeric region of the X chromosome: X inactivation in female carriers. *J Med Genet* 1998;**35**:284–7.
- 26 Allen RC, Zoghbi HY, Moseley AB, Rosenblatt HM, Belmont JW. Methylation of HpaII and HhaI sites near the polymorphic CAG repeat in the human androgen-receptor gene correlates with X chromosome inactivation. *Am J Hum Genet* 1992;**51**:1229–39.
- 27 Gorlin RJ, Cohen MM, Hennekam RCM. 2001. Oral-facial-digital syndromes. In, *Syndromes of the head and neck*, 4th ed., New York, Oxford University Press, 2001:832–43.
- 28 Whelan DT, Feldman W, Dost I. The oro-facial-digital syndrome. *Clin Genet* 1975;**8**:205–12.
- 29 Edwards M, Mulcahy D, Turner G. X-linked recessive inheritance of an orofacioidigital syndrome with partial expression in females and survival of affected males. *Clin Genet* 1988;**34**:325–32.
- 30 Morisawa T, Yagi M, Surono A, Yokoyama N, Ohmori M, Terashi H, Matsuo M. Novel double-deletion mutations of the OFD1 gene creating multiple novel transcripts. *Hum Genet* 2004;**115**:97–103.
- 31 Has C, Bruckner-Tuderman L, Muller D, Floeth M, Folkers E, Donnai D, Traupe H. The Conradi-Hunermann-Happle syndrome (CDPX2) and emopamil binding protein: novel mutations, and somatic and gonadal mosaicism. *Hum Mol Genet* 2000;**13**:1951–5.
- 32 Lyon MF. 1974. Evolution of X-chromosome inactivation in mammals. *Nature*, **250**:651–3.

Clinical Evidence—Call for contributors

Clinical Evidence is a regularly updated evidence-based journal available worldwide both as a paper version and on the internet. *Clinical Evidence* needs to recruit a number of new contributors. Contributors are healthcare professionals or epidemiologists with experience in evidence-based medicine and the ability to write in a concise and structured way.

Areas for which we are currently seeking contributors:

- Pregnancy and childbirth
- Endocrine disorders
- Palliative care
- Tropical diseases

We are also looking for contributors for existing topics. For full details on what these topics are please visit www.clinicalevidence.com/ceweb/contribute/index.jsp

However, we are always looking for others, so do not let this list discourage you.

Being a contributor involves:

- Selecting from a validated, screened search (performed by in-house Information Specialists) epidemiologically sound studies for inclusion.
 - Documenting your decisions about which studies to include on an inclusion and exclusion form, which we keep on file.
 - Writing the text to a highly structured template (about 1500–3000 words), using evidence from the final studies chosen, within 8–10 weeks of receiving the literature search.
 - Working with *Clinical Evidence* editors to ensure that the final text meets epidemiological and style standards.
 - Updating the text every 12 months using any new, sound evidence that becomes available.
- The *Clinical Evidence* in-house team will conduct the searches for contributors; your task is simply to filter out high quality studies and incorporate them in the existing text.

If you would like to become a contributor for *Clinical Evidence* or require more information about what this involves please send your contact details and a copy of your CV, clearly stating the clinical area you are interested in, to CECommissioning@bmjgroup.com.

Call for peer reviewers

Clinical Evidence also needs to recruit a number of new peer reviewers specifically with an interest in the clinical areas stated above, and also others related to general practice. Peer reviewers are healthcare professionals or epidemiologists with experience in evidence-based medicine. As a peer reviewer you would be asked for your views on the clinical relevance, validity, and accessibility of specific topics within the journal, and their usefulness to the intended audience (international generalists and healthcare professionals, possibly with limited statistical knowledge). Topics are usually 1500–3000 words in length and we would ask you to review between 2–5 topics per year. The peer review process takes place throughout the year, and out turnaround time for each review is ideally 10–14 days.

If you are interested in becoming a peer reviewer for *Clinical Evidence*, please complete the peer review questionnaire at www.clinicalevidence.com/ceweb/contribute/peerreviewer.jsp