REVIEW

Cerebral cavernous malformation: new molecular and clinical insights

N Revencu, M Vikkula

J Med Genet 2006;43:716–721. doi: 10.1136/jmg.2006.041079

Cerebral cavernous malformation (CCM) is a vascular malformation causing neurological problems, such as headaches, seizures, focal neurological deficits, and cerebral haemorrhages. CCMs can occur sporadically or as an autosomal dominant condition with variable expression and incomplete penetrance. Familial forms have been linked to three chromosomal loci, and loss of function mutations have been identified in the *KRIT1*/*CCM1*, *MGC4607*/*CCM2*, and *PDCD10*/*CCM3* genes. Recently, many new pieces of data have been added to the CCM puzzle. It has been shown that the three CCM genes are expressed in neurones rather than in blood vessels. The interaction between *CCM1* and *CCM2*, which was expected on the basis of their structure, has also been proven, suggesting a common functional pathway. Finally, in a large series of *KRIT1* mutation carriers, clinical and neuroradiological features have been characterised. These data should lead to more appropriate follow up, treatment, and genetic counselling. The recent developments will also help to elucidate the precise pathogenic mechanisms leading to CCM, contributing to a better understanding of normal and pathological angiogenesis and to the development of targeted treatment.

between the cells. No astrocytic foot and no normal nervous tissue is present within the lesion, and pericytes are rare.

Both sporadic and familial forms of CCM have been identified. In the familial form inheritance is autosomal dominant with incomplete penetrance and variable expression.^{9–10} The proportion of familial cases is estimated to be 50% in Hispanic American patients of Mexican descent,¹¹ but seems to be less in other populations.¹² Often, sporadic cases are characterised by the presence of one lesion, whereas in familial CCM, multiple lesions are present and their number is strongly correlated with the patients' age (fig 1).^{9–11} In the past, it was considered that up to 31% of familial CCM cases had only one lesion (as reviewed by Siegel¹⁰), but this was probably an overestimation because of the relative insensitivity of the radiological techniques used. The dynamic nature of the inherited lesions has also been emphasised.¹³ The diameter of lesions ranges from a few millimetres to several centimetres, and new lesions appear at a rate of 0.2 to 0.4 lesions per patient-year. The prospective haemorrhage rate is of 3.1%, and the new onset seizure rate is 2.4% per patient-year.¹⁴

Although rare, in populations other than Hispanic Americans, familial cases allow us to unravel pathogenic molecular mechanisms. Indeed, by linkage analysis, three genetic loci have been implicated, on chromosomal arms 7q (*CCM1*), 7p (*CCM2*), and 3q (*CCM3*) (table 1).^{15–16} The symptoms seem to be similar, yet the clinical penetrance was estimated at 88, 100 and 63% in the three loci, respectively.¹⁶ The first break through led to the identification of *KRIT1* (Krev Interaction Trapped1) as the *CCM1* gene.^{17–18} It encodes a protein containing known protein–protein interaction domains: four ankyrin repeats, a C-terminal FERM domain and one NPxY motif. Although without known function, *KRIT1* was thought to be an intracellular effector owing to its structure and its initial identification by yeast two-hybrid screen as a *KREV1*/*RAP1a* interactor.¹⁹ Characterisation of *KRIT1* expression during early angiogenesis and in human adult organs, and demonstration of a role in arterial morphogenesis and identity in mice lacking *Ccm1*, have started to unravel *KRIT1* function.^{20–23} In addition, the identification of the *CCM2*^{24–25} and *CCM3* genes²⁶ has paved the way toward understanding the molecular pathways involved in CCM formation.

Cerebral cavernous malformation (CCM; OMIM 116860) is a vascular malformation characterised by closely clustered enlarged capillary-like channels with a single layer of endothelium without intervening brain parenchyma.¹ CCMs are mostly located in the brain, but are also observed in the spinal cord, retina, and as hyperkeratotic cutaneous capillary-venous malformations on the skin.^{2–3} The prevalence is about 0.5%, based on cerebral magnetic resonance imaging (MRI) and necropsy studies of large cohorts of patients.^{4–5} However, the clinical prevalence is much lower, as only 20–30% of individuals are symptomatic,⁴ usually between the third and the fifth decade of life. The main symptoms are headaches, seizures, focal neurological deficits, and cerebral haemorrhages.

Ultrastructurally, these lesions consist of endothelium lined vascular sinusoids embedded in a dense collagenous matrix. They are characterised by the absence or abnormality of blood–brain barrier components.^{6–8} The tight junctions between the endothelial cells are poorly formed or absent, with gaps observed

See end of article for authors' affiliations

Correspondence to: Professor Miikka Vikkula, Laboratory of Human Molecular Genetics, Christian de Duve Institute of Cellular Pathology, Université catholique de Louvain, Avenue Hippocrate 74, BP 75.39, B-1200 Brussels, Belgium; vikkula@bchm.ucl.ac.be

Received 20 January 2006
Revised version received 8 March 2006
Accepted for publication 9 March 2006
Published Online First 29 March 2006

Abbreviations: CCM, cerebral cavernous malformation

Figure 1 (A) Axial T2 weighted magnetic resonance imaging sequence; solitary lesion (white arrow) in the right occipital lobe in a 3 year old boy. (B) Axial T2 weighted gradient echo sequence; multiple supratentorial cerebral cavernous malformations (black arrows) in a young adult.

ADVANCEMENTS REGARDING KRIT1

Clinical developments

As expected on the basis of linkage,¹⁶ a mutation in *KRIT1* was found in approximately 40% of familial CCM cases. So far about 100 distinct mutations have been reported; however, until recently the associated phenotype has been poorly described. Denier *et al* carried out a detailed clinical, neuroradiological, and molecular analysis of 64 *KRIT1* families.²⁷ Nearly half the patients had their first neurological symptoms before 25 years of age, which is earlier than in sporadic CCM. In their cohort, the maximum clinical penetrance was about 62% on the basis of 202 *KRIT1* mutation carriers (table1). This percentage is smaller than the clinical penetrance previously associated with the CCM1 locus (88%),¹⁶ and highlights the importance of large clinical studies to allow precise genetic counselling. This discrepancy could be explained by the size of the cohort (64 v 8 families) but also by the age limit of the *KRIT1* mutation carriers included (10 v 20 years), older patients being more susceptible to developing symptoms than younger ones. Denier *et al* also showed that the radiological penetrance is incomplete, as at least five asymptomatic mutation carriers—aged from 27 to 48 years—did not show any cerebral lesion on highly sensitive gradient echo MRI sequences. In addition, at least four patients had only one lesion on gradient echo MRI sequences, showing that in some patients the familial nature of the lesion on the basis of multiplicity could be overlooked.

New insights into the expression and function of KRIT1 protein

During recent years, many studies have attempted to identify the function of KRIT1 and the mechanism through which it causes CCM. Despite the vascular nature of CCMs, KRIT1 mRNA and protein have been detected in astrocytes, neurones, and various epithelial cells in adults.^{20–22} Importantly, KRIT1 protein, but not mRNA, was also detected in vascular endothelial cells during early angiogenesis, as well as in capillaries and arterioles of several human adult organs.^{20–22} Krit1 was also shown to co-localise with microtubules in bovine endothelial cells. However, the Krit1 polyclonal antibody used, detected a much smaller protein (58–60 kDa) than predicted (78–84 kDa).²⁸ Another antibody has since identified the predicted size protein on western blot, but the primary location of Krit1 awaits elucidation.²⁹

Murine embryos lacking the *Ccm1* gene suggested an essential role of Krit1 in arterial morphogenesis and identity.²³ The homozygous mutant embryos died in mid-gestation and the vascular defect was associated with downregulation of artery specific markers: *Efnb2*, *Dll4*, and *Notch4*. This suggests that the starting point of the development of CCM could be defective arterialisation.³⁰ This is surprising, as CCMs are slow flow lesions. However, it could be that Krit1 is important for arterial differentiation, and if deficient leads by default to venous identity, with abnormal morphology. If so, Krit1 could also be involved in the pathogenic pathways leading to arteriovenous malformations.

Krit1 does not have exactly the same effect in mouse as in man. Analysis of 20 *Ccm1*^{+/-} mice did not show any cavernous malformation.³⁰ In contrast to man, an additional deficit may be needed, as shown by *Ccm1/Trp53* double mutant mice. Cerebral vascular malformations, presenting features reminiscent of cavernous malformation but also of venous malformation and capillary telangiectasia, were observed in 55% of the *Ccm1*^{+/-}*Trp53*^{-/-} mice. As neither *Trp53*^{+/-} nor *Trp53*^{-/-} mice present vascular lesions, the *p53* tumour suppressor gene is not enough to cause them, but suggests that cell cycle regulators can act as modifiers in the pathogenesis of cerebrovascular malformations.

The initially reported interaction between KRIT1 and RAP1a has not been confirmed. Instead, an interaction with integrin cytoplasmic domain associated protein-1α (ICAP1α) was identified (fig 2).^{31–32} The latter is known to participate in integrin β1 mediated cell adhesion and migration. The interactions of β1 integrin and KRIT1 with ICAP1α occur similarly, through a NPxY motif/PTB domain, which suggests that integrin signalling plays a role in CCM pathogenesis. ICAP1α and KRIT1 possess a functional nuclear localisation sequence and both of them have the capacity to shuttle between the cytoplasm and the nucleus (fig 2).^{29–33} Moreover,

Table 1 Genes involved in familial cerebral cavernous malformations (http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=OMIM)			
	CCM1	CCM2	CCM3
Locus	7q11–q22	7p15–13	3q25.2–q27
Gene	<i>KRIT1</i>	<i>MGC4607</i> (malcavernin)	<i>PDCD10</i>
OMIM	604214	603284	603285
Mutation type	Loss of function	Loss of function?	Loss of function?
Second hit/trans-heterozygosity	1 Patient	?	?
	?	?	?
Molecular function	Modulator of ICAP1α, malcavernin interaction, association with microtubules?	KRIT1 interaction, scaffold for MEKK3	?
Cellular/tissue function	Cell adhesion/migration, arterial morphogenesis/identity	Osmoregulation ?	Apoptosis?
Clinical penetrance	62–88%	?	?
References	12–15, 20, 24, 26, 28, 29, 40	13, 21, 22, 26	13, 23, 26

Figure 2 Schematic presentation of molecular pathways involving the CCM proteins. ● NPXY motif; ○ phosphotyrosine binding (PTB) domain; dashed lines, hypothetical interactions. ICAP1 α interacts with the β 1 integrin cytoplasmic domain and controls cell spreading/cell proliferation on fibronectin. It was suggested that KRIT1 could be a modulator of this pathway. KRIT1/CCM2/ICAP1 α can form a ternary complex. ICAP1 α and KRIT1 can go to the nucleus. It is not known if they go separately or together. Hyperosmotic environment stimulates in mammalian cells the Rac1-OSM(CCM2)-MEKK3-MKK3-p38MAPK pathway leading to osmoregulation. KRIT1/CCM2/MEKK3 form a ternary complex.

ICAP1 α is able to sequester KRIT1 in the nucleus.²⁹ It seems that KRIT1 acts as an intracellular signalling molecule through extracellular/adhesion signals, which may be important for the activation of differentiation programmes that determine arterial identity.

IDENTIFICATION OF THE CCM2 GENE

Two teams independently identified *MGC4607* (malcavernin) to be the *CCM2* gene (table 1).^{24, 25} They used two different approaches: sequencing of positional candidate genes and loss of heterozygosity mapping, respectively. After exclusion of mutations in the *KRIT1* gene, Liquori and co-workers identified eight different mutations in nine of 27 patients, and Denier *et al* in 10 of 30 patients. This corresponds to the expected frequency of about 20%, based on previous linkage data.¹⁶ All but one mutation provoked either a premature termination codon or a deletion of the first exon, suggesting loss of function. In one case, a missense mutation, L198R, was reported. Verlaan *et al* did not identify any mutation in *MGC4607* in a cohort of 31 sporadic patients lacking *KRIT1* mutation: 21 with a single and 10 with multiple malformations.³⁴

The *CCM2* gene has several orthologues in various vertebrates, but no paralogue. Northern blot analysis showed ubiquitous expression in human adult tissues (MTC1 panel, Clontech), similar to *KRIT1*. In situ hybridisation studies have shown *Ccm2* mRNA expression in neurones^{35, 36} and astrocytes,³⁷ similar to *Ccm1* expression at embryonic and adult stages. *Ccm2* expression was also transiently observed in meningeal and parenchymal cerebral vessels.

Interestingly, the *CCM2* protein (malcavernin) contains a phosphotyrosine binding (PTB) domain, similar to ICAP1 α , suggesting an interaction between *KRIT1* and malcavernin, and a common functional pathway. Indeed, Zawistowski *et al* showed that *KRIT1* and malcavernin interact and that malcavernin is capable of sequestering *KRIT1* in the cytoplasm (fig 2).²⁹ The only *CCM2* missense mutation

reported so far—L198R, located in the PTB domain—was able to inhibit this interaction. This suggests that loss of the *KRIT1*–malcavernin interaction could contribute to CCM pathogenesis, possibly by its regulation of *KRIT1* shuttling to and from the nucleus. Compared with ICAP1 α , the NPXY motif was not critical for the *KRIT1*–malcavernin interaction. Moreover, *KRIT1*, malcavernin, and ICAP1 α were found in a ternary complex, suggesting that other interaction sites exist.

The murine orthologue of malcavernin, OSM (osmosensing scaffold for MekK3), was shown to modulate the MekK3 dependent p38Mapk activation induced by hyperosmotic shock (fig 2).³⁸ Knockdown of endogenous *CCM2* in HEK293 cells resulted in marked inhibition of sorbitol mediated p38 activation, which was confirmed in murine *Ccm2*^{+/−} embryonic fibroblasts.^{29, 38} As *KRIT1* was identified by immunoprecipitation in a ternary complex with malcavernin and MEKK3, this suggests a possible function of the CCM1/2 complex in p38MAPK activation.²⁹ Moreover, upon sorbitol treatment the CCM1/2 complex was shifted from the cytoplasm to the cell periphery. It is well known that PTB domain containing proteins have the capacity to bind phospholipids in addition to peptides, and preliminary experiments suggest that OSM would be able to do this as well.³⁹ It could be that this pathway also lies downstream of integrin mediated activation. Interestingly, mice lacking MekK3 or p38 α Map kinase have significant defects in placental angiogenesis and in blood vessel development, in particular in the head region.^{40, 41}

IDENTIFICATION OF THE CCM3 GENE

The latest breakthrough in unravelling CCM pathogenesis was the identification of *PDCD10* as the *CCM3* gene (table 1).²⁶ Bergametti and coworkers again used loss of heterozygosity mapping, proving the efficiency of this method in loss of function based disorders. They found seven distinct mutations including one large deletion in eight of 20 families studied. As point/small mutations in the *KRIT1* and the

MGC4607 genes had already been excluded, a larger proportion of mutations in the *PDCD10* gene was expected, according to previous linkage data.¹⁶ Similarly, two other groups identified *PDCD10* mutations in a minority of CCM families lacking *KRIT1* and *MGC4607* mutations: two of 15 and three out of 29, respectively.^{42–43} Moreover, in the study undertaken by Liquori and co-workers, in one large CCM3 locus linked family, the *PDCD10* gene was excluded by a recombination event and these investigators suggested the existence of a fourth CCM gene mapping near or within chromosome 3q26.3–3q27.2.

As in the case of the *MGC4607* gene, no paralogue was identified for *PDCD10* in the human genome. Instead, database searches revealed several highly conserved orthologues in vertebrates and invertebrates, highlighting an evolutionary conservation. Like *KRIT1* and malcavernin, *PDCD10* is ubiquitously expressed on the basis of northern blot analysis. As with *Ccm1* and *Ccm2*, in situ hybridisation showed *Ccm3* mRNA expression in neuronal cells at embryonic and adult stages.³⁵ Similarly, its expression coincided with that of *Ccm2* in meningeal and parenchymal cerebral vessels. Otherwise little is known about this gene. Curiously, its expression was found to be upregulated in the human myeloid cell line TF-1, upon induction of apoptosis.⁴⁴ As the CCM 1, 2, and 3 phenotypes seem to be very similar if not the same, and as the mRNA expression patterns overlap, *PDCD10* most probably has a function closely linked to that of CCM1 and 2.

FUTURE

The identification of the three CCM genes represents an important step towards the elucidation of the molecular basis of CCM. In addition, large prospective studies of genotype–phenotype association can now be undertaken. This would allow evaluation of clinical penetrance, age at onset, frequency and severity of symptoms, evolution, and response to treatment, all pivotal factors for designing accurate patient care guidelines. Other interesting questions could be addressed. What is the percentage of familial CCM and what is the frequency of de novo germline mutations? Is the *KRIT1* associated hyperkeratotic cutaneous capillary-venous malformation also present in patients with mutations in the *MGC4607* or the *PDCD10* genes, suggesting an important function in cutaneous angiogenesis for all three proteins? Is there a significant co-occurrence of CCMs with developmental venous anomalies as suggested,⁴⁵ and if so, what is the basis for this? Do all CCM families have a mutation in one of the three known genes, or do others exist? Twelve families in the screening carried out by Bergametti *et al* did not harbour a mutation in any of the three CCM genes.²⁶ Large deletions and non-sensitivity of the techniques used could explain this, but it also opens the door for the existence of a fourth gene, as recently suggested by Liquori *et al*.⁴³ Another aspect for deeper investigation is the aetiology of sporadic CCM: are they caused by genetic or environmental factors or both? If genetic in origin, are the causative genes the same as in the familial forms? Sporadic patients with multiples lesions seem to harbour *KRIT1* mutations in approximately the same proportion as familial cases.^{46–47} These mutations are either de novo or inherited from an asymptomatic parent. If this is also true for malcavernin and *PDCD10*, multiple CCMs can be postulated to have a genetic aetiology. Thus sporadic cases with multiple lesions need to be considered as familial cases, which is of major importance for patient care and genetic counselling. In contrast, sporadic cases with only one malformation may indeed differ in aetiology, with no increase in risk for progeny. Based on these data, we suggest a clinico-genetic risk evaluation scheme (fig 3).

Figure 3 Scheme for evaluation of genetic risk.

The pathogenic mechanism leading from the heterozygous germline mutation to CCM formation is poorly understood. Two major hypotheses have been put forward: haploinsufficiency and paradominant inheritance. The latter has been favoured and could explain several CCM features: the localised nature and the number of lesions (usually one in sporadic cases versus multiple in familial cases) and the age at the first symptom (earlier in familial cases). Moreover, this mechanism has been shown to be true for a patient with another inherited multifocal vascular malformation, glomuvenous malformation.⁴⁸ Gault and coworkers also identified two truncating biallelic mutations in *KRIT1* in one CCM patient with multiple lesions, although two previous screens had failed to detect somatic mutations in 20 and 72 patients, respectively (including familial and sporadic ones).^{49–51} While the techniques used may not have been sensitive enough, other mechanisms are likely to contribute to CCM pathogenesis.⁵² One explanation is trans-heterozygosity, in which case a patient with a germline mutation in the *CCM1* gene would have a somatic mutation in *CCM2* or *CCM3* gene and so forth. This could explain intrafamilial clinical variability. This can now be tested in CCM lesions, as well as in the *Ccm1*^{+/–} *Trp53*^{–/–} murine model, but may need laser capture dissection to enrich the subpopulation of cells with second hit.

As patients with a familial CCM history present similar neurological symptoms, one could assume that the three CCM genes are involved in a common functional pathway. Indeed, it has been shown that *KRIT1* interacts with malcavernin, and there is evidence that loss of this interaction contributes to the CCM pathogenesis.²⁹ In this scenario, the *CCM3* gene product, suspected to be involved in apoptosis, should be a member of this complex or play a role in a *KRIT1*/malcavernin pathway. As *KRIT1* and malcavernin are suggested to be involved in $\beta 1$ integrin signalling through ICAP1, with possible downstream signalling via p38MAPK, and as *PDCD10* was identified as a gene involved in apoptosis, it may be that the CCM pathway functions in cell adhesion governed survival. If the CCM genes have such a role in vascular endothelial cells or neural cells or both, the enlarged endothelial lined cerebral vascular channels could result from inhibited apoptosis. This would be similar to what has been proposed for cutaneous venous malformations, which are caused by TIE-2 point mutations that lead to increased Akt activity.^{53–54}

As CCM1, CCM2, and CCM3 seem to be expressed in neurones rather than in blood vessels, the vascular phenotype should result from a defect in signalling between these two juxtaposed structures. Interestingly, it has been shown that vascular and neuronal development are closely linked with several proteins—for example, neuropilin and VEGF, having a functional role in both (reviewed by Carmeliet⁵⁵). Moreover, some studies have shown the importance of neuronal invasion of primary capillary plexus for its proper remodeling and maturation (reviewed by Eichmann *et al*⁵⁶). Supplementary in vitro and in vivo studies on the three CCM proteins are clearly needed: co-immunoprecipitation and co-localisation studies implicating CCM3, and the generation of conditional homozygous, heterozygous, and compound heterozygous mutant mice. This should yield further fundamental insights, especially as to whether the primary defect is vascular or neuronal. Despite the lack of detailed mechanistic understanding of CCM formation, the important discoveries that have been made enable precise molecular diagnosis in patients with a family history or with multiple lesions, testing that should now be taken into clinical practice to allow more appropriate follow up, treatment, and genetic counselling.

ACKNOWLEDGEMENTS

We are grateful to Professors Ph Clapuyt and G Cosnard for supplying the MRI images. NR is supported by Cliniques universitaires St Luc and Université catholique de Louvain, Brussels, Belgium. MV, a “Maître de Recherches du FNRS”, is supported by grants from the Interuniversity Attraction Poles initiated by the Belgian Science Policy, network 5/25, the FNRS (Fonds national de la recherche scientifique), the ARC (Actions de Recherche Concertées-Communauté française de Belgique), the European Commission, programme project LSHG-CT-503573, and the National Institutes of Health, programme project P01 AR048564.

Authors' affiliations

N Revencu, M Vikkula, Laboratory of Human Molecular Genetics, Christian de Duve Institute of Cellular Pathology, Université catholique de Louvain, Brussels, Belgium

N Revencu, Centre for Human Genetics, Cliniques universitaires St Luc, Brussels, Belgium

Conflicts of interest: none declared

REFERENCES

- Russell DS, Rubenstein JLR. *Pathology of tumors of the nervous system*, 5th edition. Baltimore: Williams and Wilkins, 1989.
- Labauge P, Enjolras O, Bonerandi JJ, Laberge S, Dandurand M, Joujoux JM, Tournier-Lasserre E. An association between autosomal dominant cerebral cavernomas and a distinctive hyperkeratotic cutaneous vascular malformation in 4 families. *Ann Neurol* 1999;**45**:250–4.
- Eerola I, Plate KH, Spiegel R, Boon LM, Mulliken JB, Vikkula M. KRIT1 is mutated in hyperkeratotic cutaneous capillary-venous malformation associated with cerebral capillary malformation. *Hum Mol Genet* 2000;**9**:1351–5.
- Robinson JR, Awad IA, Little JR. Natural history of the cavernous angioma. *J Neurosurg* 1991;**75**:709–14.
- Otten P, Pizzolato GP, Rilliet B, Berney J. [131 cases of cavernous angioma (cavernomas) of the CNS, discovered by retrospective analysis of 24,535 autopsies]. *Neurochirurgie*. 1989;**35**: 82–3, 128–31.
- Wong JH, Awad IA, Kim JH. Ultrastructural pathological features of cerebrovascular malformations: a preliminary report. *Neurosurgery* 2000;**46**:1454–9.
- Clatterbuck RE, Eberhart CG, Crain BJ, Rigamonti D. Ultrastructural and immunocytochemical evidence that an incompetent blood-brain barrier is related to the pathophysiology of cavernous malformations. *J Neurol Neurosurg Psychiatry* 2001;**71**:188–92.
- Tu J, Stoodley MA, Morgan MK, Storer KP. Ultrastructural characteristics of hemorrhagic, nonhemorrhagic, and recurrent cavernous malformations. *J Neurosurg* 2005;**103**:903–9.
- Labauge P, Laberge S, Brunereau L, Levy C, Tournier-Lasserre E. Hereditary cerebral cavernous angiomas: clinical and genetic features in 57 French families. *Societe Francaise de Neurochirurgie. Lancet* 1998;**352**:1892–7.
- Siegel AM. Familial cavernous angioma: an unknown, known disease. *Acta Neurol Scand* 1998;**98**:369–71.
- Rigamonti D, Hadley MN, Drayer BP, Johnson PC, Hoenig-Rigamonti K, Knight JT, *et al*. Cerebral cavernous malformations. Incidence and familial occurrence. *N Engl J Med* 1988;**319**:343–7.
- Cave-Riant F, Denier C, Labauge P, Cecillon M, Maciazek J, Joutel A, Laberge-Le Couteux S, Tournier-Lasserre E. Spectrum and expression analysis of KRIT1 mutations in 121 consecutive and unrelated patients with Cerebral Cavernous Malformations. *Eur J Hum Genet* 2002;**10**:733–40.
- Zabramski JM, Wascher TM, Spetzler RF, Johnson B, Golfinos J, Drayer BP, Brown B, Rigamonti D, Browen G. The natural history of familial cavernous malformations: results of an ongoing study. *J Neurosurg* 1994;**80**:422–32.
- Moriarty JL, Wetzel M, Clatterbuck RE, Javedan S, Sheppard JM, Hoenig-Rigamonti K, Crone NE, Breiter SN, Lee RR, Rigamonti D. The natural history of cavernous malformations: a prospective study of 68 patients. *Neurosurgery* 1999;**44**:1166–71.
- Dubovsky J, Zabramski JM, Kurth J, Spetzler RF, Rich SS, Orr HT, Weber JL. A gene responsible for cavernous malformations of the brain maps to chromosome 7q. *Hum Mol Genet* 1995;**4**:453–8.
- Craig HD, Gunel M, Cepeda O, Johnson EW, Ptacek L, Steinberg GK, Ogilvy CS, Berg MJ, Crawford SC, Scott RM, Steichen-Gersdorf E, Sabroe R, Kennedy CT, Mettler G, Beis MJ, Fryer A, Awad IA, Lifton RP. Multilocus linkage identifies two new loci for a mendelian form of stroke, cerebral cavernous malformation, at 7p15–13 and 3q25.2–27. *Hum Mol Genet* 1998;**7**:1851–8.
- Laberge-le Couteux S, Jung HH, Labauge P, Houtteville JP, Lescoat C, Cecillon M, Marechal E, Joutel A, Bach JF, Tournier-Lasserre E. Truncating mutations in CCM1, encoding KRIT1, cause hereditary cavernous angiomas. *Nat Genet* 1999;**23**:189–93.
- Sahoo T, Johnson EW, Thomas JW, Kuehl PM, Jones TL, Dokken CG, Touchman JW, Gallione CJ, Lee-Lin SQ, Kosofsky B, Kurth JH, Louis DN, Mettler G, Morrison L, Gil-Nagel A, Rich SS, Zabramski JM, Boguski MS, Green ED, Marchuk DA. Mutations in the gene encoding KRIT1, a Krev-1/rap1a binding protein, cause cerebral cavernous malformations (CCM1). *Hum Mol Genet* 1999;**8**:2325–33.
- Serebriiskii I, Estojak J, Sonoda G, Testa JR, Golem EA. Association of Krev-1/rap1a with Krt11, a novel ankyrin repeat-containing protein encoded by a gene mapping to 7q21–22. *Oncogene* 1997;**15**:1043–9.
- Denier C, Gasc JM, Chapon F, Domenga V, Lescoat C, Joutel A, Tournier-Lasserre E. Krt11/cerebral cavernous malformation 1 mRNA is preferentially expressed in neurons and epithelial cells in embryo and adult. *Mech Dev* 2002;**117**:363–7.
- Guzeloglu-Kayisli O, Kayisli UA, Amankulor NM, Voorhees JR, Gokce O, DiLuna ML, Laurans MS, Luleci G, Gunel M. Krev1 interaction trapped-1/cerebral cavernous malformation-1 protein expression during early angiogenesis. *J Neurosurg* 2004;**100**:481–7.
- Guzeloglu-Kayisli O, Amankulor NM, Voorhees J, Luleci G, Lifton RP, Gunel M. KRIT1/cerebral cavernous malformation 1 protein localizes to vascular endothelium, astrocytes, and pyramidal cells of the adult human cerebral cortex. *Neurosurgery* 2004;**54**:943–9.
- Whitehead KJ, Plummer NW, Adams JA, Marchuk DA, Li DY. Ccm1 is required for arterial morphogenesis: implications for the etiology of human cavernous malformations. *Development* 2004;**131**:1437–48.
- Liquori CL, Berg MJ, Siegel AM, Huang E, Zawistowski JS, Stoffer T, Verlaan D, Balogun F, Hughes L, Leedom TP, Plummer NW, Cannella M, Maglione V, Squitieri F, Johnson EW, Rouleau GA, Ptacek L, Marchuk DA. Mutations in a gene encoding a novel protein containing a phosphotyrosine-binding domain cause type 2 cerebral cavernous malformations. *Am J Hum Genet* 2003;**73**:1459–64.
- Denier C, Goutagny S, Labauge P, Krivosic V, Arnoult M, Cousin A, Benabid AL, Comoy J, Frerebeau P, Gilbert B, Houtteville JP, Jan M, Lapierre F, Loiseau H, Menei P, Mercier P, Moreau JJ, Nivelon-Chevallier A, Parker F, Redondo AM, Scarabin JM, Tremoulet M, Zerah M, Maciazek J, Tournier-Lasserre E; Societe Francaise de Neurochirurgie. Mutations within the MGC4607 gene cause cerebral cavernous malformations. *Am J Hum Genet* 2004;**74**:326–37.
- Bergametti F, Denier C, Labauge P, Arnoult M, Boetto S, Clanet M, Coubes P, Echenne B, Ibrahim R, Irthum B, Jacquet G, Lonjon M, Moreau JJ, Neau JP, Parker F, Tremoulet M, Tournier-Lasserre E; Societe Francaise de Neurochirurgie. Mutations within the Programmed Cell Death 10 Gene Cause Cerebral Cavernous Malformations. *Am J Hum Genet* 2004;**76**:42–51.
- Denier C, Labauge P, Brunereau L, Cave-Riant F, Marchelli F, Arnoult M, Cecillon M, Maciazek J, Joutel A, Tournier-Lasserre E; Societe Francaise de Neurochirurgie; Societe de Neurochirurgie de Langue Francaise. Clinical features of cerebral cavernous malformations patients with KRIT1 mutations. *Ann Neurol* 2004;**55**:213–20.
- Gunel M, Laurans MS, Shin D, DiLuna ML, Voorhees J, Choate K, Nelson-Williams C, Lifton RP. KRIT1, a gene mutated in cerebral cavernous malformation, encodes a microtubule-associated protein. *Proc Natl Acad Sci USA* 2002;**99**:10677–82.
- Zawistowski JS, Stalheim L, Uhlik MT, Abell AN, Ancrile BB, Johnson GL, Marchuk DA. CCM1 and CCM2 protein interactions in cell signaling: implications for cerebral cavernous malformations pathogenesis. *Hum Mol Genet* 2005;**14**:2521–31.
- Plummer NW, Gallione CJ, Srinivasan S, Zawistowski JS, Louis DN, Marchuk DA. Loss of p53 sensitizes mice with a mutation in Ccm1 (KRIT1) to development of cerebral vascular malformations. *Am J Pathol* 2004;**165**:1509–18.
- Zhang J, Clatterbuck RE, Rigamonti D, Chang DD, Dietz HC. Interaction between krit1 and icap1alpha infers perturbation of integrin beta1-mediated angiogenesis in the pathogenesis of cerebral cavernous malformation. *Hum Mol Genet* 2001;**10**:2953–60.

- 32 **Zawistowski JS**, Serebriiskii IG, Lee MF, Golemis EA, Marchuk DA. KRIT1 association with the integrin-binding protein ICAP-1: a new direction in the elucidation of cerebral cavernous malformations (CCM1) pathogenesis. *Hum Mol Genet* 2002;**11**:389–96.
- 33 **Fournier HN**, Dupe-Manet S, Bouvard D, Luton F, Degani S, Block MR, Retta SF, Albiges-Rizo C. Nuclear translocation of integrin cytoplasmic domain-associated protein 1 stimulates cellular proliferation. *Mol Biol Cell* 2005;**16**:1859–71.
- 34 **Verlaan DJ**, Laurent SB, Rouleau GA, Siegel AM. No CCM2 mutations in a cohort of 31 sporadic cases. *Neurology* 2004;**63**:1979.
- 35 **Petit N**, Blecon A, Denier C, Tournier-Lasserre E. Patterns of expression of the three cerebral cavernous malformation (CCM) genes during embryonic and postnatal brain development. *Gene Expr Patterns*, 2006 Jan 30 [Epub ahead of print].
- 36 **Plummer NW**, Squire TL, Srinivasan S, Huang E, Zawistowski JS, Matsunami H, Hale LP, Marchuk DA. Neuronal expression of the Ccm2 gene in a new mouse model of cerebral cavernous malformations. *Mamm Genome* 2006;**17**:119–28.
- 37 **Seker A**, Pricola KL, Guclu B, Ozturk AK, Louvi A, Gunel M. CCM2 expression parallels that of CCM1. *Stroke* 2006;**37**:518–23.
- 38 **Uhlir MT**, Abell AN, Johnson NL, Sun W, Cuevas BD, Lobel-Rice KE, Horne EA, Dell'Acqua ML, Johnson GL. Rac-MEKK3-MKK3 scaffolding for p38 MAPK activation during hyperosmotic shock. *Nat Cell Biol* 2003;**5**:1104–10.
- 39 **Uhlir MT**, Temple B, Benchari S, Kimple AJ, Siderovski DP, Johnson GL. Structural and evolutionary division of phosphotyrosine binding (PTB) domains. *J Mol Biol* 2005;**345**:1–20.
- 40 **Adams RH**, Porras A, Alonso G, Jones M, Vintersten K, Panelli S, Valladares A, Perez L, Klein R, Nebreda AR. Essential role of p38alpha MAP kinase in placental but not embryonic cardiovascular development. *Mol Cell* 2000;**6**:109–16.
- 41 **Yang J**, Boerm M, McCarty M, Bucana C, Fidler IJ, Zhuang Y, Su B. Mek3 is essential for early embryonic cardiovascular development. *Nat Genet* 2000;**24**:309–13.
- 42 **Verlaan DJ**, Roussel J, Laurent SB, Elger CE, Siegel AM, Rouleau GA. CCM3 mutations are uncommon in cerebral cavernous malformations. *Neurology* 2005;**65**:1982–3.
- 43 **Liquori CL**, Berg MJ, Squitieri F, Ottenbacher M, Sorlie M, Leedom TP, Cannella M, Maglione V, Ptacek L, Johnson EW, Marchuk DA. Low frequency of PDCD10 mutations in a panel of CCM3 probands: potential for a fourth CCM locus. *Hum Mutat* 2006;**27**:118.
- 44 **Wang YG LH**, Zhang YM, Ma DL. cDNA cloning and expression of an apoptosis-related gene, human TFAR-15 gene. *Science in China series C – Life Sciences* 1999;**42**:323–9.
- 45 **Abe T**, Singer RJ, Marks MP, Norbash AM, Crowley RS, Steinberg GK. Coexistence of occult vascular malformations and developmental venous anomalies in the central nervous system: MR evaluation. *Am J Neuroradiol* 1998;**19**:51–7.
- 46 **Laurans MS**, DiLuna ML, Shin D, Niazi F, Voorhees JR, Nelson-Williams C, Johnson EW, Siegel AM, Steinberg GK, Berg MJ, Scott RM, Tedeschi G, Enevoldson TP, Anson J, Rouleau GA, Ogilvy C, Awad IA, Lifton RP, Gunel M. Mutational analysis of 206 families with cavernous malformations. *J Neurosurg* 2003;**99**:38–43.
- 47 **Verlaan DJ**, Laurent SB, Sure U, Bertalanffy H, Andermann E, Andermann F, Rouleau GA, Siegel AM. CCM1 mutation screen of sporadic cases with cerebral cavernous malformations. *Neurology* 2004;**62**:1213–15.
- 48 **Brouillard P**, Boon LM, Mulliken JB, Enjolras O, Ghassibe M, Warman ML, Tan OT, Olsen BR, Viskula M. Mutations in a novel factor, glomulin, are responsible for glomuvenous malformations (“glomangiomas”). *Am J Hum Genet* 2002;**70**:866–74.
- 49 **Gault J**, Shenkar R, Recksiek P, Awad IA. Biallelic somatic and germ line CCM1 truncating mutations in a cerebral cavernous malformation lesion. *Stroke* 2005;**36**:872–4.
- 50 **Marini V**, Ferrera L, Pigatto F, Origone P, Garre C, Dorcaratto A, Viale G, Alberti F, Mareni C. Search for loss of heterozygosity and mutation analysis of KRIT1 gene in CCM patients. *Am J Med Genet A* 2004;**130**:98–101.
- 51 **Reich P**, Winkler J, Straube A, Steiger HJ, Peraud A. Molecular genetic investigations in the CCM1 gene in sporadic cerebral cavernomas. *Neurology* 2003;**60**:1135–8.
- 52 **Gallagher AR**, Hidaka S, Gretz N, Witzgall R. Molecular basis of autosomal-dominant polycystic kidney disease. *Cell Mol Life Sci* 2002;**59**:682–93.
- 53 **Viskula M**, Boon LM, Carraway KL, Calvert JT, Diamanti AJ, Goumnerov B, Pasyk KA, Marchuk DA, Warman ML, Cantley LC, Mulliken JB, Olsen BR. Vascular dysmorphogenesis caused by an activating mutation in the receptor tyrosine kinase TIE2. *Cell* 1996;**87**:1181–90.
- 54 **Morris PN**, Dunmore BJ, Tadros A, Marchuk DA, Darland DC, D'Amore PA, Brindle NP. Functional analysis of a mutant form of the receptor tyrosine kinase Tie2 causing venous malformations. *J Mol Med* 2005;**83**:58–63.
- 55 **Carmeliet P**. Blood vessels and nerves: common signals, pathways and diseases. *Nat Rev Genet* 2003;**4**:710–20.
- 56 **Eichmann A**, Makinen T, Alitalo K. Neural guidance molecules regulate vascular remodeling and vessel navigation. *Genes Dev* 2005;**19**:1013–21.