CCN2 YAPs at cancer

Andrew Leask

Received: 17 August 2008 / Accepted: 19 August 2008 / Published online: 3 September 2008
© The International CCN Society 2008

Abstract The YAP transcription coactivator has been implicated as an oncogene and is amplified in human cancers. Previously, it has been shown that CCN2 (connective tissue growth factor, CTGF) is a target of the tumor promoting YAP and its transcription factor target TEAD. A recent report in *Genes and Development* by Zhao and colleagues (*Genes Dev* 22:1962–1971 2008) has extended these initial observations to show that CCN2 plays an important role in the growth-promoting function of YAP. These data confirm the role of CCN2 as a key oncogenic mediator. This report briefly summarizes these findings.

Keywords YAP · ETS · TEAD · CCN2 · Gene regulation · Cancer

A large body of evidence has indicated that the expression of the CCN family of proteins is dysregulated in cancers and promotes angiogenesis and metastasis by an integrin-mediated pathway (Babic et al. 1998; Rachfal et al. 2004; Jiang et al. 2004; Xie et al. 2004; Shimo et al. 2006; Pickles and Leask 2007). CCN2 expression appears to correlate with high tumor grade and metastasis (Deng et al. 2007). Collectively, these results suggest that CCN2 expression may be both a marker and mediator of metastasis and that studying the basis of why CCN2 is overexpressed in cancers may enhance our understanding of the molecular basis of cancers as well as the role that CCN2 may play in these pathologies (Blom et al. 2002). Indeed, it was recently

shown that CCN2 was expressed downstream of ras in pancreatic cancer cells (Pickles and Leask 2007).

In NIH 3T3 fibroblasts, The CCN2 promoter is activated by the oncogenic transcriptional coactivator YAP and regulated by the YAP target TEAD (Leask et al. 2003). In an intriguing recent report, Zhao and colleagues (2008) extend this observation by first confirming that YAP and members of the TEAD family interact (Vassilev et al. 2001). Among the confirmed YAP-inducible genes in NIH 3T3 fibroblasts and MCF10A breast cancer cells was CCN2, and this response depended on TEAD transcription factors (Zhao et al. 2008). Chromatin immunoprecipitation experiments verified that TEAD and YAP bound the endogenous CCN2 promoter. Zhao and colleagues (2008) YAP activated the CCN2 promoter via previously identified TEAD/YAP response elements (Leask et al. 2003). Zhao and colleagues (2008) used siRNA to knockdown CCN2 expression in the YAP-overexpressing MCF10A cells, and showed that reduction in CCN2 expression decreased the acini growth and reversed the rough surface morphology in 3D culture. However, CCN2 knockdown did not affect the apparent epithelial-mesenchymal transition (EMT) in monolayer culture. These results indicate that in CCN2 may play an important role in the growth-promoting function but may not be required for the EMT-inducing activity of YAP.

It is interesting to note that the TEAD/YAP response elements of the CCN2 promoter are also functional binding elements for the Ets family (ETS) of transcription factors (van Beek et al. 2006) which also are implicated in the pathology of cancer (Liebermann and Zerbini 2006). ETS transcription factors are characterized by an evolutionally conserved Ets domain and play important roles in cell development, cell differentiation, cell proliferation, apoptosis and tissue remodeling. Most of them are downstream nuclear

A. Leask (✉)
CIHR Group in Skeletal Development and Remodeling,
Division of Oral Biology and Department of Physiology and
Pharmacology, Schulich School of Medicine and Dentistry,
Dental Sciences Building, University of Western Ontario,
London, ON N6A 5C1, Canada
e-mail: Andrew.leask@schulich.uwo.ca

targets of Ras-MAP kinase signaling, and the deregulation of ETS genes results in the malignant transformation of cells (Oikawa 2004). Possible functional interactions between the TEAD/YAP and ETS transcription factors have not been speculated on before. However, it is interesting to note that Ets1 is required, but TEAD1 blocks, the TGF β -induction of CCN2 in fibroblasts (Leask et al. 2003; van Beek et al. 2006).

Collectively the above results suggest that although the TEAD/YAP and ETS factors may have similar activities in some contexts, alteration in TEAD/YAP:ETS ratios may have profound effects on other areas, such as the resistance of cancer cells to TGF β . But, perhaps more important from the context of the CCN family, the results from Zhao and colleagues (2008) strongly support the notion that CCN2 plays a key role in oncogenesis and that blocking the action of CCN2, or other members of the CCN family, may be suitable for precisely targeted drug therapy.

References

- Babic AM, Kireeva ML, Kolesnikova TV, Lau LF (1998) CYR61, a product of a growth factor-inducible immediate early gene, promotes angiogenesis and tumor growth. *Proc Natl Acad Sci USA* 95:6355–6360, doi:[10.1073/pnas.95.11.6355](https://doi.org/10.1073/pnas.95.11.6355)
- Blom IE, Goldschmeding R, Leask A (2002) Gene regulation of connective tissue growth factor: new targets for antifibrotic therapy? *Matrix Biol* 21:473–482, doi:[10.1016/S0945-053X\(02\)00055-0](https://doi.org/10.1016/S0945-053X(02)00055-0)
- Deng YZ, Chen PP, Wang Y, Yin D, Koeffler HP, Li B et al (2007) Connective tissue growth factor is overexpressed in esophageal squamous cell carcinoma and promotes tumorigenicity through beta catenin-T-cell factor/Lef signaling. *J Biol Chem* 282:36571–36581, doi:[10.1074/jbc.M704141200](https://doi.org/10.1074/jbc.M704141200)
- Jiang WG, Watkins G, Fodstad O, Douglas-Jones A, Mokbel K, Mansel RE (2004) Differential expression of the CCN family members Cyr61, CTGF and Nov in human breast cancer. *Endocr Relat Cancer* 11:781–791, doi:[10.1677/erc.1.00825](https://doi.org/10.1677/erc.1.00825)
- Leask A, Holmes A, Black CM, Abraham DJ (2003) Connective tissue growth factor gene regulation. Requirements for its induction by transforming growth factor-beta 2 in fibroblasts. *J Biol Chem* 278:13008–13015, doi:[10.1074/jbc.M210366200](https://doi.org/10.1074/jbc.M210366200)
- Libermann TA, Zerbini LF (2006) Targeting transcription factors for cancer gene therapy. *Curr Gene Ther* 6:17–33, doi:[10.2174/156652306775515501](https://doi.org/10.2174/156652306775515501)
- Oikawa T (2004) ETS transcription factors: possible targets for cancer therapy. *Cancer Sci* 95:626–633, doi:[10.1111/j.1349-7006.2004.tb03320.x](https://doi.org/10.1111/j.1349-7006.2004.tb03320.x)
- Pickles M, Leask A (2007) Analysis of CCN2 promoter activity in PANC-1 cells: regulation by ras/MEK/ERK. *J Cell Commun Signal* 1:85–90, doi:[10.1007/s12079-007-0008-9](https://doi.org/10.1007/s12079-007-0008-9)
- Rachfal AW, Luquette MH, Brigstock DR (2004) Expression of connective tissue growth factor (CCN2) in desmoplastic small round cell tumour. *J Clin Pathol* 57:422–425, doi:[10.1136/jcp.2003.012344](https://doi.org/10.1136/jcp.2003.012344)
- Shimo T, Kubota S, Yoshioka N, Ibaragi S, Isowa S, Eguchi T et al (2006) Pathogenic role of connective tissue growth factor (CTGF/CCN2) in osteolytic metastasis of breast cancer. *J Bone Miner Res* 21:1045–1059, doi:[10.1359/jbmr.060416](https://doi.org/10.1359/jbmr.060416)
- van Beek JP, Kennedy L, Rockel JS, Bernier SM, Leask A (2006) The induction of CCN2 by TGF beta1 involves Ets-1. *Arthritis Res Ther* 8:R36
- Vassilev A, Kaneko KJ, Shu H, Zhao Y, DePamphilis ML (2001) TEAD/TEF transcription factors utilize the activation domain of YAP65, a Src/Yes-associated protein localized in the cytoplasm. *Genes Dev* 15:1229–1241, doi:[10.1101/gad.888601](https://doi.org/10.1101/gad.888601)
- Xie D, Yin D, Wang HJ, Liu GT, Elashoff R, Black K et al (2004) Levels of expression of CYR61 and CTGF are prognostic for tumor progression and survival of individuals with gliomas. *Clin Cancer Res* 10:2072–2081, doi:[10.1158/1078-0432.CCR-0659-03](https://doi.org/10.1158/1078-0432.CCR-0659-03)
- Zhao B, Ye X, Yu J, Li L, Li W, Li S et al (2008) TEAD mediates YAP-dependent gene induction and growth control. *Genes Dev* 22:1962–1971, doi:[10.1101/gad.1664408](https://doi.org/10.1101/gad.1664408)