Published in final edited form as:

Arch Microbiol. 2008 November ; 190(5): 547–557. doi:10.1007/s00203-008-0405-3.

An *N*-acyl homolog of mycothiol is produced in marine actinomycetes

Gerald L. Newton and Robert C. Fahey

Department of Chemistry and Biochemistry, University of California, San Diego, La Jolla, CA 92093, USA e-mail: rcfahay@ucsd.edu fax: 858-5344864

Paul R. Jensen and William Fenical

Center for Marine Biotechnology and Biomedicine, Scripps Institution of Oceanography, University of California, San Diego, La Jolla, CA 92093, USA

John B. MacMillan

Department of Biochemistry, UT Southwestern Medical Center, 5323 Harry Hines Blvd, Dallas TX 75390-9038, USA

Abstract

Marine actinomycetes have generated much recent interest as a potentially valuable source of novel antibiotics. Like terrestrial actinomycetes the marine actinomycetes are shown here to produce mycothiol as their protective thiol. However, a novel thiol, U25, was produced by MAR2 strain CNQ703 upon progression into stationary phase when secondary metabolite production occurs and became the dominant thiol. MSH and U25 were maintained in a reduced state during early stationary phase but become significantly oxidized after 10 days in culture. Isolation and structural analysis of the monobromobimane derivative identified U25 as a homolog of mycothiol in which the acetyl group attached to the nitrogen of cysteine is replaced by a propionyl residue. This *N*-propionyl-desacetyl-mycothiols was present in 13 of the 17 strains of marine actinomycetes examined, including 5 strains of *Salinispora* and representatives of the MAR2, MAR3, MAR4 and MAR6 groups. Mycothiol and its precursor, the pseudodisaccharide 1-*O*-(2-amino-2-deoxy- α -D-glucopyranosyl)-D-*myo*-inositol, were found in all strains. High levels of mycothiol *S*-conjugate amidase activity, a key enzyme in mycothiol-dependent detoxification, was found in most strains. The results demonstrate that major thiol/disulfide changes accompany secondary metabolite production and suggest that mycothiol-dependent detoxification is important at this developmental stage.

Keywords

mycothiol; *N*-propionyl-desacetyl-mycothiols; marine actinomycetes; *Salinispora arenicola*; Mca

Introduction

There has been a growing interest in recent years in the study of marine actinomycetes (Bull et al. 2005; Fenical and Jensen 2006; Bull and Stach 2007). A major motivation underlying such studies is the promise that they represent a source of new antibiotics, and a number of potentially important compounds have been isolated from marine-derived actinomycetes (Feling et al. 2003; Fiedler et al. 2005; Jensen et al. 2005b). Phylogenetic analyses based on sequences of 16S rRNA have provided evidence that marine actinomycetes are distinct from

their terrestrial counterparts (Mincer et al. 2002). Although characterization of secondary metabolites produced by marine actinomycetes has received considerable attention, little is known about the levels of common metabolic intermediates in these new species. In the present study we set out to determine the low-molecular-weight thiol distribution in marine actinomycetes.

The diverse biochemistry associated with major thiols produced by microorganisms and by various marine organisms is the subject of a review in which mycothiol emerges as a thiol of major significance in the gram positive actinomycetes (Hand and Honek 2005). The main protective thiol found in gram negative bacteria and most eukaryotes is glutathione but glutathione is not found in most Gram positive bacteria (Fahey et al. 1978; Fahey and Sundquist 1991). Mycothiol (MSH, Fig. 1A) is the protective thiol produced by most terrestrial actinomycetes and it has thus far not been found outside of these bacteria (Newton et al. 1996; Newton and Fahey 2002). Mycothiol biosynthesis is initiated by the glycosyltransferase MshA (Newton et al. 2003) whose substrates have been identified as UDP-GlcNAc and 1L-inositol-1-P (Newton et al. 2006). An as yet unidentified phosphatase, MshA2, a deacetylase MshB (Newton et al. 2000a), an ATP-dependent cysteine:GlcN-Ins ligase MshC (Sareen et al. 2002), and the mycothiol synthase MshD (Koledin et al. 2002) complete the biosynthesis of mycothiol as summarized in Fig. 1A.

MSH forms *S*-conjugates (MSR) with a wide range of toxins, Fig. 1B (Newton et al. 2000b), including the antibiotic rifamycin S (Steffek et al. 2003). The enzyme mycothiol *S*-conjugate amidase (Mca) cleaves such *S*-conjugates to produce a mercapturic acid, AcCySR, which can be excreted from the cell, and the pseudo-disaccharide GlcN-Ins (Fig. 1B), which can be recycled for MSH biosynthesis (Newton et al. 2000b). This detoxification process may be important for protection of antibiotic producing bacteria against their own antibiotics (Newton and Fahey 2002). Mycobacterial Mca has lower activity with MSH itself and under limiting nutrient conditions the AcCys produced is rapidly deacetylated to liberate Cys (Bzymek et al. 2007). Both Cys and GlcN-Ins can be further degraded to generate energy and important metabolic intermediates. Mycobacteria deficient in Mca or the enzymes of MSH biosynthesis exhibit enhanced sensitivity to certain antibiotics (Newton et al. 1999; Rawat et al. 2002; Buchmeier et al. 2003) and production of MSH is required for growth of *Mycobacterium tuberculosis* (Sareen et al. 2003; Buchmeier and Fahey 2006).

Because marine actinomycetes are an important source of novel antibiotics and since mycothiol in conjunction with Mca has been shown to play a role in antibiotic detoxification in terrestrial actinomycetes, it was of interest to examine the mycothiol and Mca content in marine actinomycetes. In the present study we show that mycothiol is generally found in marine actinomycetes and the level of Mca activity is quite high. However, in stationary phase when secondary metabolites are produced the thiol composition was found to change dramatically and a new thiol, shown to be a homolog of mycothiol, predominates over mycothiol.

Materials and methods

Bacterial strains and culture

The organisms studied are listed in Table 1. These strains were isolated and phylogenetically characterized as previously described (Mincer et al. 2002b; Jensen et al. 2005a). Each strain was cultured in a 1 liter of TCG medium (3 g tryptone (Fisher), 5 g casitone (Difco), 4 g glucose, 1 liter natural seawater) for 2-10 days while shaking at 215 rpm at 25-27°C.

Assays performed

For thiol (RSH) analysis, cells were extracted in warm acetonitrile containing monobromobimane (mBBBr). The acetonitrile denatures and precipitates most proteins. The thiol-labeling agent, mBBBr, rapidly alkylates thiols and produces fluorescent derivatives for analysis by HPLC. Separate cell extractions and labeling of amines with AccQ-Fluor allowed determination of the GlcN-Ins content. A third extract was prepared and used to assay for Mca activity with MSmB as substrate. For determination of disulfide forms (RSS) an extract prepared in the presence of *N*-ethylmaleimide to block thiols and then subsequently reduced with dithiothreitol and labeled with monobromobimane was analyzed by HPLC to determine RSS levels and allow calculation of the thiol redox status (RSH/RSS).

Assay of cellular thiols

Cells were collected by centrifugation at $4000 \times g$ for 30 min at 4°C. The cells were split into 6 equal pellets (pellets A-F), the wet weight was determined using preweighed 15 ml polypropylene centrifuge tubes and the tubes stored at -70°C prior to extraction. Two pellets, typically between 0.5 and 2 g wet weight, were used for duplicate thiol analysis (pellets A and B). Pellets A and B were extracted in an equal volume of warm acetonitrile containing 4 mM monobromobimane (mBBBr; Molecular Probes) and 20 μ l 1 M HEPES, pH 8.0, per ml of extraction. The cells suspensions were incubated at 60°C, vortexed every 5 min for 15 min, and then cooled on ice. To quench the reaction and slow the formation of mBBBr hydrolysis products, 5 μ l of 5 M methanesulfonic acid was added per ml of total extract. The thiol control sample (pellet C) was treated similarly, except that the extraction buffer contained 10 mM *N*-ethylmaleimide (NEM) in place of mBBBr. The sample was incubated at 60°C for 10 min after which mBBBr was added to a final concentration of 2 mM and the sample incubated for an additional 15 min at 60°C. The control was cooled on ice and 5 μ l of 5M methanesulfonic acid was added per ml of extract. The extracts were clarified by centrifugation for 30 min at $10,000 \times g$. The pellets were dried to constant weight in a vacuum oven and weighed to provide the residual dry weight used to calculate results. The supernatant was removed to a separate tube and was usually two phases due to the saline content of the cells. The supernatant was reduced to half of the starting volume in a Savant Speed Vac, producing a single phase.

The samples were assayed using three analytical HPLC protocols developed for analysis of a wide variety of thiols. Coelution of a thiol-bimane derivative with a standard on two or more HPLC protocols was considered confirmation of the identity for that thiol. HPLC methods 1 and 2 were the primary methods used and the retention times for various biological thiols have been reported for these protocols (Fahey and Newton 1987). The retention times reported for unknown thiols (U-retention time) were those recorded using method 1M, a shortened version of method 1 as described by Koledin, et al. (Koledin et al. 2002). Quantitation was based on known standards when the identity of the thiol was confirmed and the fluorescent factor for the bimane derivative of cysteine (CySmB) was used for quantitation of unknown thiols. The control sample, in which thiols were blocked by NEM prior to reaction with mBBBr, was utilized to verify that peaks resulting from nonthiol fluorescent components of the extract did not interfere with the analysis.

Assay of GlcN-Ins

Duplicate pellets labeled D and E were extracted in an equal volume of 60°C acetonitrile containing 20 μ l per ml 1 M HEPES pH 7.5. The cells were vortexed every 5 min and incubated at 60°C for 15 min. The extract was cooled on ice and centrifuged. The supernatant was reduced to half the original volume, as above, to produce one phase. A sample of this extract was reacted with AccQ-Fluor (Waters) amine labeling reagent and the extract was assayed for GlcN-Ins according to Buchmeier et al (Buchmeier et al. 2003). The retention time for AccQ-GlcN-Ins

was typically 22-24 min using the chromatographic protocol (method 4) described by Anderberg et al (Anderberg et al. 1998).

Mycothioliol S-conjugate amidase (Mca) activity

The final cell pellet (pellet F) from each culture was extracted in neutral aqueous buffer for estimation of the Mca activity. Each pellet was suspended in 5 ml per g wet weight of 50 mM NaCl, 50 mM HEPES pH 7.5 containing 3 mM 2-mercaptoethanol and 35 μ M each of *N*- α -*p*-tosyl-L-phenylalanylchloromethyl ketone and *N*- α -*p*-tosyl-L-lysinechloromethyl ketone to inhibit proteolytic degradation. The cells were disrupted by sonicating 5 times for 30 sec using a 50% pulsed sonication at maximum power (Branson Sonifier). The cells were iced between sonications and the extract was maintained between 5 and 15°C at all times. The cell free supernatant was collected by centrifugation at $12,000 \times g$ for 30 min at 4°C and the protein content was assayed by the Bradford dye binding assay (Biorad). For the amidase reaction, extraction buffer containing MSmB derived from *M. smegmatis* (Newton et al. 2000b) was preheated at 37°C. The reaction was started by the addition of 12.5 μ l of extract to produce a final volume of 250 μ l containing 0.1 mM MSmB. At 0, 3, 6, and 12 min a 50 μ l sample was removed and mixed with 200 μ l of 40 mM methanesulfonic acid to rapidly terminate the reaction. The sample was centrifuged for 3 min at $13,000 \times g$ and was analyzed by HPLC without dilution for the formation of AcCySmB as described previously (Newton et al. 2000b). The amount of extract was adjusted as necessary to produce accurate initial rate estimates and the activity is reported as specific activity, nmol/min/mg protein.

Time course analysis of strain CNQ703

A culture of CNQ703 was inoculated in 2 liters TCG medium with shaking (250 rpm) at 28°C. Samples (200 ml) were removed at 2, 3, 6, 8, and 10 days. The cells were pelleted by centrifugation into 4 pellets of equal size. Three of the pellets were extracted in 1 ml of 4 mM mBBr in acetonitrile plus 20 μ l of 1M HEPES pH 8.0 and processed as described above for thiol samples. The fourth pellet was extracted in 1 ml 10 mM NEM plus 20 μ l of 1M HEPES pH 8.0 and incubated at 60°C for 15 min. The sample was cooled on ice and the cell debris removed by centrifugation for 5 minutes at $13,000 \times g$. The supernatant was reduced in a speed vac to generate a 0.5 ml one phase supernatant. An aliquot (100 μ l) of the NEM supernatant was reacted with 2 mM mBBr for 15 min at 60°C to serve as a control for the thiol samples. The balance of the NEM sample was reacted with 10 mM dithiothreitol (DTT) for 20 min at 23°C to reduce disulfides and then with 20 mM mBBr for an additional 15 min to derivatize the DTT released thiols. A redox ratio was calculated as the mean of the triplicate thiol analyses divided by the DTT reduced soluble disulfides.

Isolation and structure elucidation of U25

U25 was isolated as the bimane derivative from a 20 g pellet derived from 5 liters of a 3-day CNQ703 culture. The cells were collected by centrifugation at $4000 \times g$ for 30 min at 4°C and extracted in 200 ml of 50% acetonitrile (20 mM HEPES pH 8.0) containing 2 mM mBBr at 60°C. After centrifugation the supernatant was reduced to 25 ml using a rotary evaporator and diluted to 100 ml with water. The bimane derivatives were concentrated by solid phase extraction on two Sep Pak C-18 (Waters, 5 g each) columns. The columns were washed with aqueous 0.1% TFA, and eluted with 0.1% aqueous TFA containing 10%, 25%, and 50% methanol. U25mB eluted in the 10-25% methanol fraction and this was concentrated on a Savant Speed Vac. U25mB was purified on a Vydac preparative HPLC column (#218TP1022) using aqueous 0.1% TFA and methanol as solvents and a linear 0-40% methanol gradient. U25mB (10 mg) eluted in about 20% methanol and was further purified in a second pass on the same column. For structural studies residual water and TFA were removed in a Savant Speed Vac.

NMR studies of U25-mB

The structure of U25-mB was determined by a combination of 1D and 2D NMR experiments. All spectra were obtained on a Varian INOVA spectrometer with ^1H NMR spectra being recorded at 500 MHz and ^{13}C NMR spectra recorded at 125 MHz, using residual solvent as an internal reference (CD_3OD , δ_{H} 3.30 ppm, δ_{C} 49.0 ppm). Carbon multiplicities were determined using a multiplicity edited HSQC experiment, while ^1H - ^1H and ^1H - ^{13}C correlations were established by gradient homonuclear correlated spectroscopy (gCOSY) and gradient homonuclear multiple bond coherence (gHMBC) NMR experiments.

Results

Initial findings

We set out to sample a broad spectrum of marine actinomycetes representative of the main MAR groupings (Table 1). Mycothiol was detected in all of the strains initially studied but we soon found that an additional thiol that could not be assigned as a known biological thiol (Fahey and Newton 1987; Newton and Fahey 1995) was also present. It was designated U25 based upon the retention time (25 min) under HPLC analysis using HPLC method 1M (Koledin et al. 2002), one of three protocols used to analyze thiols. In some cases the U25 content exceeded the MSH content. One MAR2 strain, CNQ703, appeared to produce U25 in large amounts and was selected for more detailed study.

Thiol composition and redox status of MAR2 Strain CNQ703

We determined the thiol content (Fig. 2A) and the thiol redox ratio (Fig. 2B) from day 2 to day 10 of the growth of CNQ703. The culture could not be monitored by optical density owing to the production of intensely absorbing secondary metabolites beginning on day 3, but the yield of residual dry weight per liter of culture remained constant at 4.1 ± 1.0 g per liter from day 2 to day 10 indicating that the culture was in stationary phase. During this period the MSH content fell 50-fold (Fig. 2A). However, a novel thiol (U25) was also produced whose content increased almost 40-fold from day 2 to day 3 and U25 was the dominant thiol from day 3 to day 10. A second unidentified thiol, U22, remained at a low level throughout. The penultimate intermediate in MSH biosynthesis, Cys-GlcN-Ins (Fig. 1), was also observed on day 3 (0.037 ± 0.007 $\mu\text{mol per g}$) but was not observed above background (≤ 0.001 $\mu\text{mol per g}$) at any other time.

Thiol components present in disulfide forms (RSS) were measured and this permitted calculation of the thiol redox ratios (RSH/RSS) that are shown in Fig. 2B. The only terrestrial actinomycetes for which redox ratios have been determined are *M. smegmatis* and *M. tuberculosis*. In *M. smegmatis* the MSH/MSS ratio was found to fall from ~ 1000 in exponential phase to ~ 200 in stationary phase (Newton et al. 2005) whereas in *M. tuberculosis* it drops from ~ 180 in exponential phase to ~ 50 in stationary phase (Buchmeier et al. 2006). The values from Fig. 2B (MSH/MSS = 30-50) at days 2-6 are comparable to the stationary phase values for *M. tuberculosis*. The sharp fall on days 8 and 10 reflects both a fall in MSH and an increase in MSS. The redox state of U25 is comparable to that of cysteine (Fig. 2) and is several fold more oxidized than that of MSH.

Structure of U25

The bimane derivative of unknown thiol U25 was purified from mBBR labeled cell extracts of CNQ703 by solid phase extraction and HPLC. Electrospray positive ion mass spectroscopy produced a molecular ion peak at m/z 713.2 (MNa^+) and a weaker one at m/z 691.1 (MH^+), corresponding to molecules of 14 mass units greater than that of MSH. A strong fragmentation ion at m/z 511.1 reflects the loss of a 180 Da fragment from the MH^+ peak at m/z 691.1. This

can be assigned to loss of a $C_6H_{12}O_6$ inositol unit as was seen in the mass spectrum of the bimane derivative of mycothiol, MSmB (Newton et al. 1995).

Further structural analysis of the bimane derivative of U25 (U25mB) was achieved by NMR studies. The 500 MHz proton spectrum (Fig. 3) was very similar to that of MSmB (Newton et al. 1995) but contained additional signals ascribable to a CH_3CH_2CO residue (δ 1.14 ppm (t, $J = 7.5$ Hz); δ 2.3 ppm (q, $J = 7.5$ Hz)) and it lacked the singlet resonance at 2.09 ppm for the acetyl moiety of AcCys in MSmB. The *myo*-inositol and glucosamine residues were established by a combination of gCOSY NMR correlations and coupling constants analysis (Table 2). The structure and the numbering of the carbons for *N*-propionyl-desacetyl-mycothiol is shown in Fig. 4. The amide linkage between the anomeric carbon of glucosamine (C1) and carbonyl carbon C1'' of cysteine was determined based on a HMBC NMR correlation between H1 (δ 5.11 ppm (d, $J = 4.0$ Hz)) and C1'' (δ 172.5 ppm), while the *N*-propionyl residue was established to be at C2'' based on a HMBC NMR correlation between H2'' (δ 4.56 ppm (dd, $J = 8.5, 5.0$ Hz)) and C4'' (δ 177.4 ppm). These studies establish that U25 is a close relative of MSH with the *N*-acetyl group replaced by an *N*-propionyl residue, i.e., 1-O-[2-[(2R)-2-(propionylamino)-3-mercapto-1-oxopropyl]amino]-2-deoxy- α -D-glucopyranosyl]-D-*myo*-inositol (propionyl-Cys-GlcN-Ins). We propose the common name *N*-propionyl-desacetyl-mycothiol (PdAMSH) for this homolog of mycothiol.

Survey of marine actinomycetes

Diverse marine-derived actinomycetes belonging to the families Micromonosporaceae, Streptomycetaceae, and Thermomonosporaceae were selected to test the generality of production of MSH, PdAMSH and related compounds production (Table 1). These strains include four representatives of the newly described marine genus *Salinispora* (Maldonado et al. 2005), as well as examples of the clearly defined clade within the Streptomycetaceae that has tentatively been called MAR2. An earlier study of the terrestrial *Streptomyces clavuligerus* had established that a thiol, later identified as mycothiol (Newton et al. 1995), is produced at substantial and roughly stable levels during exponential growth and into stationary phase when secondary metabolites are produced (Newton et al. 1993). For the present survey the various marine actinomycetes strains were grown and harvested 3-10 days after inoculation, following entry into stationary phase when secondary metabolite production became evident. This provided ample material from a one-liter culture to prepare samples for determination of thiols and GlcN-Ins in duplicate and to assay Mca activity.

In the *Salinispora* strains (MAR1), MSH was the major thiol with most values for MSH content between 2 and 4 μ mol per g residual dry weight (Table 3). One strain, CNH725, gave a relatively low value (0.8 μ mol per g residual dry weight) value for MSH content. The level of PdAMSH was 2-20% that of MSH in these samples. Significant levels (0.3-1.5 μ mol per g) of an additional unknown, U18, were found in all strains except CNR107 and lower levels of U27 (0.14-0.38 μ mol per g) were present in all strains except CNH725. The level of Mca activity was quite high in all *Salinispora* strains. Hydrogen sulfide was found at significant levels in all *Salinispora* strains (Table 3). This likely derives from iron-sulfur proteins, the labile sulfide being released and reacted with mBBR under our extraction conditions. The values for H_2S are typical of those found in a wide range of bacteria (Newton et al. 1996).

The three "*Marinispora*" (Kwon et al. 2006) (MAR2) strains examined all had significant levels of GlcN-Ins, MSH and PdAMSH (Table 3). PdAMSH was the dominant thiol in strains CNQ695 and CNQ703 but MSH was 6-fold higher in strain CNR252. All three strains had a high level of Mca activity.

Five strains formed a coherent clade, designated MAR3, within the Streptomycetaceae. All were found to produce GlcN-Ins and MSH at measurable levels that were well reproduced in

replicate samples of a given culture (Table 3). Significant HPLC peaks assignable to PdAMSH were found in three of the five strains but the level in strains CNQ687 and CNQ857 was undetectable ($<0.1 \mu\text{mol per g}$). Two strains of the MAR4 group gave widely different results, one strain (CNR530) producing the highest level of MSH of any strain tested and the other (CNR525) producing the lowest MSH content but both had similar amounts of GlcN-Ins (Table 3). GlcN-Ins is a dedicated intermediate in mycothiol biosynthesis and has no known function except MSH production. Strain CNR530 having the highest MSH content also had significant, but much lower PdAMSH whereas PdAMSH was undetectable in strain CNQ525.

One strain (CNR363) from the marine Thermomonosporaceae MAR5 group and a second (CNR431) from the MAR6 group differed 5-fold in the GlcN-Ins content and nearly 40-fold in MSH content but in opposite directions (Table 3). No PdAMSH ($<0.002 \mu\text{mol per g dry weight}$) was detected in strain CNR363 whereas strain CNR431 contained $0.43 \mu\text{mol per g dry weight}$. Substantial Mca activity was found in both strains.

Discussion

The most intriguing feature of the present results is the replacement of MSH by PdAMSH during secondary metabolite production. It appears that this phenomenon is not restricted to MAR2 strain CNQ703. A survey of representative marine actinomycetes revealed that PdAMSH was detected in all five MAR1 strains, in all three MAR2 strains, in three of five MAR3 strains, in one of two MAR4 strains, in the one MAR6 strain but not in the one MAR5 strain. Whether all of the PdAMSH producing strains exhibit profiles similar to CNQ703 (Fig. 2) was not determined but it is noted that PdAMSH did exceed MSH in the data for another MAR2 strain, CNQ695. Moreover, we have recently found that PdAMSH is also produced in mycobacteria under some conditions (G. L. Newton, N. Buchmeier and R. C. Fahey, unpublished data). Thus, synthesis of PdAMSH is not limited to the marine actinomycetes and may prove to be a more general phenomenon among the mycothiol producing Actinobacteria.

PdAMSH is the first derivative of MSH to be found in wild type Actinobacteria. However analogs of MSH in which acyl groups other than acetyl or propionyl are linked to Cys-GlcN-Ins have been reported in *mshD* null mutants of *M. smegmatis* and *M. tuberculosis* (Newton et al. 2005; Buchmeier et al. 2006). In these mutants Cys-GlcN-Ins accumulates to such high levels ($0.5 - 1.5 \mu\text{mol per g dry weight}$) that transacylation chemical reaction with succinyl-CoA leads to the generation of succinyl-Cys-GlcN-Ins, a process that occurs some 20-fold faster than chemical production of MSH from acetyl-CoA or PdAMSH from propionyl-CoA (Newton et al. 2005). Formyl-Cys-GlcN-Ins is also produced from Cys-GlcN-Ins in these mutants, presumably by an enzymatic process. Chemical transacylation of Cys-GlcN-Ins by propionyl-CoA cannot be the source of PdAMSH in the marine actinomycetes since the level of Cys-GlcN-Ins in these organisms is very low or undetectable ($<0.005 \mu\text{mol per g dry weight}$). Production of PdAMSH must involve an enzymatic process. *M. tuberculosis* MshD is known to utilize propionyl-CoA at 20-25% the rate of acetyl-CoA (Vetting et al. 2006). If the marine actinomycete MshD has similar substrate flexibility then it seems likely that it is responsible for the enzymatic production of PdAMSH. However, replacement of MSH by PdAMSH would require that the level of propionyl-CoA substantially exceed the level of acetyl-CoA. Marine actinomycetes are known to produce a number of polyketide secondary metabolites (Kwon et al. 2006; Jensen et al. 2007; Udvary et al. 2007) and propionyl-CoA is used as an acyl starter unit and to generate methylmalonyl-CoA extender units in polyketide synthesis (Walsh 2003). Thus, the production of propionyl-CoA during generation of secondary metabolites is expected and PdAMSH synthesis did coincide with the appearance of the black-green secondary metabolite on day 3 (Fig. 2).

What function does production of PdAMSH serve? One possibility is that it functions as a detoxification pathway. Degradation of odd chain and branched fatty acids (Boshoff and Barry 2005) (Koburger et al. 2005) and of cholesterol (Pandey and Sasseti 2008) leads to the production of propionyl-CoA, as does growth in the presence of propionate (Brock and Buckel 2004). Accumulation of propionyl-CoA during degradative metabolism is potentially toxic owing to accumulation of intermediates produced by its metabolism via the methylcitrate cycle (Brock 2005). Diversion of propionyl-CoA into PdAMSH production could be a possible way of avoiding such toxicity.

It is also possible that production of PdAMSH serves a regulatory or protective function involving protein thiol groups. The redox status of PdAMSH is significantly more oxidized than is that of MSH and becomes more oxidized in the latter stages of secondary metabolite production (Fig. 2). The disulfide of PdAMSH may be an inefficient substrate relative to that of MSH for mycothiol disulfide reductase (Patel and Blanchard 2001). In glutathione producing organisms such changes in thiol redox status lead to the glutathionylation of proteins (Prot-SSG formation) which is thought to be involved in regulation of protein function or protection against overoxidation of sensitive thiol groups (Ghezzi 2005; Hurd et al. 2005; Masip et al. 2006; Dalle-Donne et al. 2007). Analogous thiolation of proteins may be important when PdAMSH becomes the dominant thiol and could serve similar functions. Elaboration of the function of PdAMSH is being undertaken in studies of mycobacteria where availability of mutants defective in mycothiol biosynthesis and enzymes of mycothiol metabolism are available.

The values for MSH content determined here can be compared with those for terrestrial actinomycetes: 0.8 - 2.3 μmol per g residual dry weight for *Streptomyces clavuligerus*, increasing during the transition from exponential to stationary phase (Newton et al. 1993), and ~ 12 μmol per g residual dry weight for *Mycobacterium smegmatis* mc²155, remaining constant during transition from exponential to stationary phase (Newton et al. 2005). While most of the values in Table 3 are comparable, the values for strains CNR530 and CNR431 are the highest we have measured. Since these bacteria are readily harvested from liquid growth medium by filtration, such high-producing MSH strains may represent a suitable resource for practical production of MSH. Mycothiol is a difficult molecule to generate by chemical synthesis (Jardine et al. 2002; Nicholas et al. 2002; Lee and Rosazza 2004) and is more efficiently obtained by isolation from cells (Unson et al. 1998; Steenkamp and Vogt 2004).

The levels of Mca activity found are very high in most marine actinomycetes, especially in strain CNR003 which produced the highest activity, 133 nmol/min/mg protein. The only terrestrial actinomycete for which Mca activity has been measured is *M. smegmatis* where the activity of a crude extract was found to be ~ 1 nmol/min/mg (Newton et al. 2000b) or 30-130-fold lower than that observed in *Salinospora* strains. A role for MSH and Mca in the detoxification of antibiotics has been identified in *M. smegmatis* (Steffek et al. 2003; Rawat et al. 2004) and in *S. coelicolor* (Park and Roe 2008). Thus, it appears likely that MSH and the high Mca activity are associated with secondary metabolite detoxification in the marine actinomycetes.

In summary, several unusual aspects of mycothiol metabolism have been identified in studies of marine actinomycetes. A novel variant of mycothiol, *N*-propionyl-desacetyl-mycythiol, was shown to replace mycothiol when nutrient depletion occurs and secondary metabolites are formed. Exceptionally high levels of mycothiol *S*-conjugate amidase activity were found in nearly all marine actinomycetes examined. Two strains, CNR530 (MAR4 group) and CNR431 (MAR6 group) produced record high levels of mycothiol and may have utility as a source of this thiol.

Acknowledgements

This work was supported by NIH grant AI49174 from the National Institute of Allergy and Infectious Diseases and grant MCB-0235705 from the National Science Foundation to RCF. Additional support was provided from the NIH, National Cancer Institute, under grant CA-44848 (to WF), and in part by the National Sea Grant College Program of the U.S. Department of Commerce's National Oceanic and Atmospheric Administration under NOAA Grant # NA040AR4170038, project # R/MP-96, through the California Sea Grant College Program; and in part by the California State Resources Agency to WF. The views expressed herein do not necessarily reflect the views of any of those organizations. We thank Nancy Buchmeier for critical review of the manuscript.

References

- Anderberg SJ, Newton GL, Fahey RC. Mycothiol biosynthesis and metabolism: Cellular levels of potential intermediates in the biosynthesis and degradation of mycothiol. *J Biol Chem* 1998;273:30391–30397. [PubMed: 9804803]
- Boshoff HI, Barry CE 3rd. Tuberculosis - metabolism and respiration in the absence of growth. *Nat Rev Microbiol* 2005;3:70–80. [PubMed: 15608701]
- Brock M. Generation and phenotypic characterization of *Aspergillus nidulans* methylisocitrate lyase deletion mutants: methylisocitrate inhibits growth and conidiation. *Appl Environ Microbiol* 2005;71:5465–5475. [PubMed: 16151139]
- Brock M, Buckel W. On the mechanism of action of the antifungal agent propionate. *Eur J Biochem* 2004;271:3227–3241. [PubMed: 15265042]
- Buchmeier N, Fahey RC. The *mshA* gene encoding the glycosyltransferase of mycothiol biosynthesis is essential in *Mycobacterium tuberculosis* Erdman. *FEMS Microbiol Lett* 2006;264:74–79. [PubMed: 17020551]
- Buchmeier NA, Newton GL, Fahey RC. A mycothiol synthase mutant of *Mycobacterium tuberculosis* has an altered thiol-disulfide content and limited tolerance to stress. *J Bacteriol* 2006;188:6245–6252. [PubMed: 16923891]
- Buchmeier NA, Newton GL, Koledin T, Fahey RC. Association of mycothiol with protection of *Mycobacterium tuberculosis* from toxic oxidants and antibiotics. *Mol Microbiol* 2003;47:1723–1732. [PubMed: 12622824]
- Bull AT, Stach JE. Marine actinobacteria: new opportunities for natural product search and discovery. *Trends Microbiol* 2007;15:491–499. [PubMed: 17997312]
- Bull AT, Stach JE, Ward AC, Goodfellow M. Marine actinobacteria: perspectives, challenges, future directions. *Antonie Van Leeuwenhoek* 2005;87:65–79.
- Bzymek KP, Newton GL, Ta P, Fahey RC. Mycothiol import by *Mycobacterium smegmatis* and function as a resource for metabolic precursors and energy production. *J Bacteriol* 2007;189:6796–6805. [PubMed: 17644601]
- Dalle-Donne I, Rossi R, Giustarini D, Colombo R, Milzani A. S-glutathionylation in protein redox regulation. *Free Radic Biol Med* 2007;43:883–898. [PubMed: 17697933]
- Fahey RC, Brown WC, Adams WB, Worsham MB. Occurrence of glutathione in bacteria. *J Bacteriol* 1978;133:1126–1129. [PubMed: 417060]
- Fahey RC, Newton GL. Determination of low-molecular-weight thiols using monobromobimane fluorescent labeling and high-performance liquid chromatography. *Methods Enzymol* 1987;143:85–96. [PubMed: 3657565]
- Fahey RC, Sundquist AR. Evolution of glutathione metabolism. *Adv Enzymol Relat Areas Mol Biol* 1991;64:1–53. [PubMed: 1675828]
- Feling RH, Buchanan GO, Mincer TJ, Kauffman CA, Jensen PR, Fenical W. Salinosporamide A: a highly cytotoxic proteasome inhibitor from a novel microbial source, a marine bacterium of the new genus *Salinospora*. *Angew Chem Int Ed Engl* 2003;42:355–357. [PubMed: 12548698]
- Fenical W, Jensen PR. Developing a new resource for drug discovery: marine actinomycete bacteria. *Nat Chem Biol* 2006;2:666–673. [PubMed: 17108984]
- Fiedler HP, Bruntner C, Bull AT, Ward AC, Goodfellow M, Potterat O, Puder C, Mihm G. Marine actinomycetes as a source of novel secondary metabolites. *Antonie Van Leeuwenhoek* 2005;87:37–42. [PubMed: 15726289]

- Ghezzi P. Regulation of protein function by glutathionylation. *Free Radic Res* 2005;39:573–580. [PubMed: 16036334]
- Hand CE, Honek JF. Biological chemistry of naturally occurring thiols of microbial and marine origin. *J Nat Prod* 2005;68:293–308. [PubMed: 15730267]
- Hurd TR, Costa NJ, Dahm CC, Beer SM, Brown SE, Filipovska A, Murphy MP. Glutathionylation of mitochondrial proteins. *Antioxid Redox Signal* 2005;7:999–1010. [PubMed: 15998254]
- Jardine MA, Spies HS, Nkambule CM, Gammon DW, Steenkamp DJ. Synthesis of mycothiol, 1D-1-O-(2-[N-acetyl-L-cysteinyl]amino-2-deoxy- α -D-glucopyranosyl)-*myo*-inositol, principal low molecular mass thiol in the actinomycetes. *Bioorg Med Chem* 2002;10:875–881. [PubMed: 11836093]
- Jensen PR, Gontang E, Mafnas C, Mincer TJ, Fenical W. Culturable marine actinomycete diversity from tropical Pacific Ocean sediments. *Environ Microbiol* 2005a;7:1039–1048. [PubMed: 15946301]
- Jensen PR, Mincer TJ, Williams PG, Fenical W. Marine actinomycete diversity and natural product discovery. *Antonie Van Leeuwenhoek* 2005b;87:43–48. [PubMed: 15726290]
- Jensen PR, Williams PG, Oh DC, Zeigler L, Fenical W. Species-specific secondary metabolite production in marine actinomycetes of the genus *Salinispora*. *Appl Environ Microbiol* 2007;73:1146–1152. [PubMed: 17158611]
- Koburger T, Weibezahn J, Bernhardt J, Homuth G, Hecker M. Genome-wide mRNA profiling in glucose starved *Bacillus subtilis* cells. *Mol Genet Genomics* 2005;274:1–12. [PubMed: 15809868]
- Koledin T, Newton GL, Fahey RC. Identification of the mycothiol synthase gene (*mshD*) encoding the acetyltransferase producing mycothiol in actinomycetes. *Arch Microbiol* 2002;178:331–337. [PubMed: 12375100]
- Kwon HC, Kauffman CA, Jensen PR, Fenical W. Marinomycins A-D, antitumor-antibiotics of a new structure class from a marine actinomycete of the recently discovered genus “marinispora”. *J Am Chem Soc* 2006;128:1622–1632. [PubMed: 16448135]
- Lee S, Rosazza JP. First total synthesis of mycothiol and mycothiol disulfide. *Org Lett* 2004;6:365–368. [PubMed: 14748594]
- Maldonado LA, Fenical W, Jensen PR, Kauffman CA, Mincer TJ, Ward AC, Bull AT, Goodfellow M. *Salinispora arenicola* gen. nov., sp. nov. and *Salinispora tropica* sp. nov., obligate marine actinomycetes belonging to the family Micromonosporaceae. *Int J Syst Evol Microbiol* 2005;55:1759–1766. [PubMed: 16166663]
- Masip L, Veeravalli K, Georgiou G. The many faces of glutathione in bacteria. *Antioxid Redox Signal* 2006;8:753–762. [PubMed: 16771667]
- Mincer TJ, Jensen PR, Kauffman CA, Fenical W. Widespread and persistent populations of a major new marine actinomycete taxon in ocean sediments. *Appl Environ Microbiol* 2002;68:5005–5011. [PubMed: 12324350]
- Newton GL, Arnold K, Price MS, Sherrill C, delCardayré SB, Aharonowitz Y, Cohen G, Davies J, Fahey RC, Davis C. Distribution of thiols in microorganisms: Mycothiol is a major thiol in most actinomycetes. *J Bacteriol* 1996;178:1990–1995. [PubMed: 8606174]
- Newton GL, Av-Gay Y, Fahey RC. *N*-Acetyl-1-D-*myo*-inosityl-2-amino-2-deoxy- α -D-glucopyranoside deacetylase (MshB) is a key enzyme in mycothiol biosynthesis. *J Bacteriol* 2000a;182:6958–6963. [PubMed: 11092856]
- Newton GL, Av-Gay Y, Fahey RC. A novel mycothiol-dependent detoxification pathway in mycobacteria involving mycothiol *S*-conjugate amidase. *Biochemistry* 2000b;39:10739–10746. [PubMed: 10978158]
- Newton GL, Bewley CA, Dwyer TJ, Horn R, Aharonowitz Y, Cohen G, Davies J, Faulkner DJ, Fahey RC. The structure of U17 isolated from *Streptomyces clavuligerus* and its properties as an antioxidant thiol. *Eur J Biochem* 1995;230:821–825. [PubMed: 7607257]
- Newton GL, Fahey RC. Determination of biothiols by bromobimane labeling and high-performance liquid chromatography. *Methods Enzymol* 1995;251:148–166. [PubMed: 7651194]
- Newton GL, Fahey RC. Mycothiol biochemistry. *Arch Microbiol* 2002;178:388–394. [PubMed: 12420157]
- Newton GL, Fahey RC, Cohen G, Aharonowitz Y. Low molecular weight thiols in streptomycetes and their potential role as antioxidants. *J Bacteriol* 1993;175:2734–2742. [PubMed: 8478335]

- Newton GL, Koledin T, Gorovitz B, Rawat M, Fahey RC, Av-Gay Y. The glycosyltransferase gene encoding the enzyme catalyzing the first step of mycothiol biosynthesis (*mshA*). *J Bacteriol* 2003;185:3476–3479. [PubMed: 12754249]
- Newton GL, Ta P, Bzymek K, Fahey RC. Biochemistry of the initial steps of mycothiol biosynthesis. *J Biol Chem* 2006;281 in press
- Newton GL, Ta P, Fahey RC. A mycothiol synthase mutant of *Mycobacterium smegmatis* produces novel thiols and has an altered thiol redox status. *J Bacteriol* 2005;187:7309–7316. [PubMed: 16237013]
- Newton GL, Unson MD, Anderberg SJ, Aguilera JA, Oh NN, delCardayré SB, Davies J, Av-Gay Y, Fahey RC. Characterization of a *Mycobacterium smegmatis* mutant defective in 1-D-*myo*-inosityl-2-amino-2-deoxy- α -D-glucopyranoside and mycothiol biosynthesis. *Biochem Biophys Res Commun* 1999;255:239–244. [PubMed: 10049692]
- Nicholas GM, Kovac P, Bewley CA. Total synthesis and proof of structure of mycothiol bimane. *J Amer Chem Soc* 2002;124:3492–3493. [PubMed: 11929223]
- Pandey AK, Sassetti CM. Mycobacterial persistence requires the utilization of host cholesterol. *Proc Natl Acad Sci U S A* 2008;105:4376–4380. [PubMed: 18334639]
- Park J-H, Roe J-H. Mycothiol regulates and is regulated by a thiol-specific anti-sigma factor RsrA and sigmaR in *Streptomyces coelicolor*. *Mol Microbiol*. 2008 in press
- Patel MP, Blanchard JS. *Mycobacterium tuberculosis* mycothione reductase: pH dependence of the kinetic parameters and kinetic isotope effects. *Biochemistry* 2001;40:3119–3126.
- Rawat M, Newton GL, Ko M, Martinez GJ, Fahey RC, Av-Gay Y. Mycothiol-deficient *Mycobacterium smegmatis* mutants are hypersensitive to alkylating agents, free radicals and antibiotics. *Antimicrob Agents Chemother* 2002;46:3348–3355. [PubMed: 12384335]
- Rawat M, Uppal M, Newton G, Steffek M, Fahey RC, Av-Gay Y. Targeted mutagenesis of the *Mycobacterium smegmatis* *mca* gene, encoding a mycothiol-dependent detoxification protein. *J Bacteriol* 2004;186:6050–6058. [PubMed: 15342574]
- Sareen D, Newton GL, Fahey RC, Buchmeier NA. Mycothiol is essential for growth of *Mycobacterium tuberculosis* Erdman. *J Bacteriol* 2003;185:6736–6740. [PubMed: 14594852]
- Sareen D, Steffek M, Newton GL, Fahey RC. ATP-dependent L-cysteine:1D-*myo*-inosityl 2-amino-2-deoxy- α -D-glucopyranoside ligase, mycothiol biosynthesis enzyme MshC, is related to class I cysteinyl-tRNA synthetases. *Biochemistry* 2002;41:6885–6890. [PubMed: 12033919]
- Steenkamp DJ, Vogt RN. Preparation and utilization of a reagent for the isolation and purification of low-molecular-mass thiols. *Anal Biochem* 2004;325:21–27. [PubMed: 14715280]
- Steffek M, Newton GL, Av-Gay Y, Fahey RC. Characterization of *Mycobacterium tuberculosis* mycothiol S-conjugate amidase. *Biochemistry* 2003;42:12067–12076. [PubMed: 14556638]
- Uduary DW, Zeigler L, Asolkar RN, Singan V, Lapidus A, Fenical W, Jensen PR, Moore BS. Genome sequencing reveals complex secondary metabolome in the marine actinomycete *Salinispora tropica*. *Proc Natl Acad Sci U S A* 2007;104:10376–10381. [PubMed: 17563368]
- Unson MD, Newton GL, Davis C, Fahey RC. An immunoassay for the detection and quantitative determination of mycothiol. *J Immunol Meth* 1998;214:29–39.
- Vetting MW, Yu M, Rendle PM, Blanchard JS. The Substrate-induced Conformational Change of *Mycobacterium tuberculosis* Mycothiol Synthase. *J Biol Chem* 2006;281:2795–2802. [PubMed: 16326705]
- Walsh, C. Antibiotics: Actions, Origins, Resistance. ASM Press; Washington, D.C.: 2003.

Fig. 1.
(A) Structure and biosynthesis of mycothiol (MSH). **(B)** Cleavage of mycothiol *S*-conjugate (MSR) by the amidase Mca, MSH resynthesis from GlcN-Ins, and degradation of MSH by Mca as a source of energy and metabolic intermediates.

Fig. 2. Variation in thiol content (**A**) and thiol redox ratio (**B**) with time for MAR 2 strain CNQ703: ○, MSH; □, U25 (*N*-propionyl-desacetyl-mycothiols; PdAMSH); ◇, CoA; Δ, Cys; ▽, U22. Arrow indicates the onset of secondary metabolite production. Error bars represent the range of values, where larger than the symbol, from duplicate determinations.

Fig. 3. 500 MHz proton NMR spectrum of U25-mB; HDO and CHD₂OD, solvent impurities; C, contaminant.

N-propionyl-desacetyl-mycothioliol

Fig. 4. Structure of *N*-propionyl-desacetyl-mycothioliol (PdAMSH) with numbering of carbons as listed in Table 2.

Table 1

Origin and reference data of strains studied.

Strain #	Phylotype (Group ^a)	Origin	Depth (meters)	16S RNA Accession No.
CNR107	<i>Salinispora arenicola</i> (MAR1)	Guam ^b	40	AY4646534
CNR003	<i>Salinispora pacifica</i> (MAR1)	Guam ^b	184	DQ224161
CNR643	<i>Salinispora arenicola</i> (MAR1)	Bahamas ^c	1	AY040619
CNH725	<i>Salinispora arenicola</i> (MAR1)	Red Sea ^c	20	AY040621
CNQ976	<i>Salinispora arenicola</i> (MAR1)	Guam ^b	72	AY464533
CNR252	Streptomycetaceae (MAR2)	Guam ^b	3	AY464536
CMQ695	Streptomycetaceae (MAR2)	Guam ^b	500	AY464537
CNQ703	Streptomycetaceae (MAR2)	Guam ^b	75	AY464538
CNQ857	Streptomycetaceae (MAR3)	Guam ^b	10	AY464545
CNQ719	Streptomycetaceae (MAR3)	Guam ^b	<1	AY464544
CNQ698	Streptomycetaceae (MAR3)	Guam ^b	3	AY464543
CNQ766	Streptomycetaceae (MAR3)	Guam ^b	500	AY464546
CNQ687	Streptomycetaceae (MAR3)	Guam ^b	<1	AY464542
CNQ530	Streptomycetaceae (MAR4)	Guam ^b	45	AY464541
CNQ525	Streptomycetaceae (MAR4)	San Diego	160	EF177816
CNR363	Thermomonosporaceae (MAR5)	Guam ^b	3	AY464547
CNR431	Thermomonosporaceae (MAR6)	Guam ^b	500	AY464548

^aReference (Jensen et al. 2005b).^b(Fenical and Jensen 2006).^cReference (Mincer et al. 2002)

Table 2NMR Data for U25-mB recorded in CD₃OD

Carbon #	¹ H δ (mult., #, J (Hz))	¹³ C δ	HMBC correlations
D-glucosamine			
1	5.11 (d, 1H, J = 4.0)	100.2	H2, H1'
2	3.90 (dd, 1H, J = 10.5, 4.0)	55.7	H1, H3
3	3.73 (dd, 1H, J = 10.5, 9.5)	73.0	H2, H4
4	3.32 (t, 1H, J = 9.5)	72.4	H3, H5, H6
5	3.82 (m, 1H)	74.4	H4, H6
6	3.92 (m, 1H)	62.8	H4, H5
	3.65 (dd, 1H, J = 11, 2.5Hz)		
D-myo-inositol			
1'	3.47 (dd, 1H, J = 9.5, 3.0)	80.5	H1, H2', H6'
2'	4.12 (t, 1H, J = 3.0)	73.4	H1', H3'
3'	3.34 (dd, 1H, J = 9.5, 3.0)	73.2	H2', H6'
4'	3.59 (t, 1H, J = 9.5)	74.0	H3', H5'
5'	3.16 (t, 3H, J = 9.5)	76.5	H4', H6'
6'	3.78 (t, 1H, J = 9.5)	74.1	H1', H5'
N-propionyl cysteine			
1''	-	172.5	H2, H2'', H3''
2''	4.56 (dd, 1H, J = 8.5, 5.0)	54.2	H3''
3''	2.89 (dd, 1H, J = 13.5, 8.5)	35.1	H2'', H1'''
	3.14 (dd, 1H, J = 13.5, 5.0)		
4''	-	177.4	H5'', H6''
5''	2.30 (q, 2H, J = 7.5)	29.9	H6''
6''	1.14 (t, 3H, J = 7.5)	10.2	H5''
Bimane			
1'''	3.92 (s, 2H)	26.2	
2'''		114.9	H1''', H8'''
3'''		148.9	H1''', H8'''
4'''		162.3	H8'''
5'''		162.8	H9'''
6'''		149.7	H9''', H10'''
7'''		112.9	H9''', H10'''
8'''	1.89 (s, 3H)	7.0	
9'''	1.80 (s, 3H)	6.5	
10'''	2.47 (s, 3H)	11.8	

Table 3
GlcN-Ins, thiol, and Mca assays of marine actinomycetes.

Organism	Cellular Content, $\mu\text{moles per g residual dry weight}^d$					PdAMSH (U25)	Mca Activity ^b
	GlcN-Ins	Cys	MSH	CoA			
<i>Salinispora</i> MAR1 ^c	0.3-1.5	0.4-1.8	0.8-4	0.1-0.4	0.03-0.8		30-133
" <i>Marinispora</i> " MAR2 ^d	0.1-0.5	0.6-2.9	2.3-3.8	<0.02-0.8	0.6-9		27-71
Marine streptomycetaceae							
MAR3 ^e	0.1-0.9	0.1-2.1	0.5-12	<0.03-2.4	<0.1-0.4		3-38
MAR4 ^f	0.35, 0.40	0.6, 3.6	0.2, 33	0.5, 10	<0.1, 0.44		29, 43
Marine thermomonosporaceae							
MAR5 CNR363 ^g	0.6	0.3	0.7	0.14	<0.02		20
MAR6 CNR431 ^h	0.11	1.1	27	1.0	0.43		1.2

^a Range of values from duplicate analyses of each strain; GSH and Cys-GlcN-Ins <0.005 in all extracts.

^b Assayed at 37°C in 0.1 mM MSMB, 50 mM NaCl, 1 mM 2-mercaptoethanol, 50 mM HEPES pH 7.5, nmol/min/mg protein.

^c Strains CNR107, CNR003, CNH643, CNH725 and CNQ976; also contained SSO₃⁻² (0.3-1.3) and H₂S (1.5-2.5).

^d Strains CNR252, CNQ695 and CNQ703; also contained SSO₃⁻² (1.1-1.8) and H₂S (2.2-6.6).

^e Strains CNQ687, CNQ698, CNQ719, CNQ766, CNQ857; also contained SSO₃⁻² (0.5-2.5) and H₂S (0.7-6).

^f Strains CNR530 and CNQ525.

^g Also contained SSO₃⁻² (1.0) and H₂S (1.1).

^h Also contained SSO₃⁻² (1.8) and H₂S (7).