Glycan microarray of Globo H and related structures for quantitative analysis of breast cancer

Cheng-Chi Wang^{*†}, Yen-Lin Huang^{*†}, Chien-Tai Ren^{*}, Chin-Wei Lin^{*}, Jung-Tung Hung^{*}, Jyh-Cherng Yu[‡], Alice L. Yu^{*§}, Chung-Yi Wu^{*§}, and Chi-Huey Wong^{*§}

^{*}The Genomics Research Center, Academia Sinica, Taipei, Taiwan; [†]Institute of Biochemical Science, National Taiwan University, Taipei, Taiwan; and [‡]Tri-Service General Hospital, Taipei, Taiwan

Contributed by Chi-Huey Wong, May 23, 2008 (sent for review January 1, 2008)

Cancer-associated carbohydrate antigens are often found on the surface of cancer cells. Understanding their roles in cancer progression will lead to the development of new therapeutics and high-sensitivity diagnostics for cancers. Globo H is a member of this family, which is highly expressed on breast cancer cells. Here, we report the development of a glycan microarray of Globo H and its analogs for measurement of the dissociation constants on surface ($K_{D,surf}$) with three different monoclonal antibodies (VK-9, Mbr1, and anti-SSEA-3), to deduce their binding specificity. The glycan microarray was also used to detect the amount of antibodies present in the plasma of breast cancer patients and normal blood donors. It was shown that the amount of antibodies against Globo H from breast cancer patients were significantly higher than normal blood donors, providing a new tool for possible breast cancer diagnosis. Compared with the traditional ELISA method, this array method required only atto-mole amounts of materials and is more effective and more sensitive (5 orders of magnitude). The glycan microarray thus provides a new platform for use to monitor the immune response to carbohydrate epitopes after vaccine therapy or during the course of cancer progression.

array | diagnosis

Globo H is a hexasaccharide, which is a member of a family of antigenic carbohydrates that are highly expressed on a various types of cancers, especially cancers of breast, prostate and lung (1–4). It is expressed on the cancer cell surface as a glycolipid and possibly as a glycoprotein (5, 6). Furthermore, it has been established that the serum of breast cancer patients contains high levels of antibodies against the Globo H epitope, and this epitope is also targeted by the monoclonal antibodies Mbr1 (5, 7–8) and VK9 (9) in immunohistochemistry studies. Although certain normal tissues also react with Mbr1, including normal breast, pancreas, small bowel, and prostate tissue (2, 6, 10), the antigen in these tissues is predominantly localized at the secretory borders where access to the immune system is restricted. These two findings provide the rationale for evaluating human antibody against Globo H as a diagnosis for breast cancer.

Glycan microarray is a very powerful tool for glycobiological study (11–36). It may effectively mimic the presentation of glycans on the cell membrane to exhibit multivalent interactions with receptors with high affinity and specificity (37). In addition, only a very small amount of material is required for arraying. Thus, the microarray may provide a more appropriate and more sensitive system for studying protein–carbohydrate interactions than the traditional solution-phase ELISA analysis. Recently, our group has used Globo H and its truncated analog microarrays to profile the binding specificity of its monoclonal and polyclonal antibodies in a qualitative manner (38). In that preliminary study, we found that the polyclonal antibodies from a breast cancer patient can bind both Globo H and its truncated analog without fucose (Gb5), whereas Mbr1 is specific for Globo H (38). The binding specificity studies of monoclonal antibodies and human polyclonal antibodies associated with Globo H have prompted us to further investigate the following questions:

Fig. 1. Chemical structure of Globo H and abbreviations of Globo H analogs.

- Are Globo H and Gb5 differentially expressed in disease stages?
- Can the glycan array be used for the quantitative and high-sensitivity detection of anti-Globo H antibody present in the serum as an indication of cancer?
- Can the array be used to measure the activity of fucosyltransferase in breast cancer cells at different stages, to screen inhibitors of the enzyme and to monitor the immune response to the Globo H-based vaccine?

Results and Discussion

Specificities of Antibodies. The Globo H and its analogs (Fig. 1) were prepared according to the one-pot programmable protocol (38) and were covalently attached onto NHS-coated glass slides

Author contributions: A.L.Y., C.-Y.W., and C.-H.W. designed research; C.-C.W., Y.-L.H., C.-T.R., C.-W.L., J.-T.H., and J.-C.Y. performed research; C.-C.W., Y.-L.H., J.-T.H., and A.L.Y. analyzed data; and C.-C.W., A.L.Y., C.-Y.W., and C.-H.W. wrote the paper.

The authors declare no conflict of interest.

[§]To whom correspondence may be addressed. E-mail: ayu@gate.sinica.edu.tw, cyiwu@gate.sinica.edu.tw, or ch Wong@gate.sinica.edu.tw.

This article contains supporting information online at www.pnas.org/cgi/content/full/0804923105/DCSupplemental.

© 2008 by The National Academy of Sciences of the USA

Fig. 4. Glycan array is a mimic of cell surface expressing glycans for study of multivalent interaction.

saturation at 5 μM printing concentration. To determine the dissociation constant on surface, we plotted VK9 concentrations against the fluorescence intensity at different concentrations of printed sugar. At the printing concentrations from 100 μM to 5 μM , the $K_{D,\text{surf}}$ values determined were narrowly distributed from the individual curves (Fig. 3 and Table 1). However, if the printing concentration was <5 μM , the fluorescence intensity of each spot was too weak to be detected with a significant signal-to-noise (S/N) ratio (data not shown). In addition, the intensities of those spots at

lower printing concentrations did not converge to the binding curves generated by the Langmuir isotherm. This result illustrates that at the printing concentrations <5 μM , the sugar density on surface is too low to exhibit a multivalent interaction.

Using the same method, the $K_{D,\text{surf}}$ values were measured for the truncated Globo H analogs interacting with antibodies [see the supporting information (SI), Fig. 2, and Table 2]. The relative binding specificity of both VK9 and Mbr1 for the sugar epitopes was Globo H > tetrasacchride Bb4. In addition, we observed that the relative binding affinity of antibodies for Globo H was Mbr1 > VK9 but that for Bb4 was in an opposite order. This study indicates that the glycan microarray is a good mimic of cell surface expressing multiple glycans and is thus useful for the study of multivalent interaction (Fig. 4).

Detection of Serum Antibody Against Globo H. The binding intensities in a range of Globo H and tetrasaccharide Bb4 concentration (5–100 μM) for the two antibodies (VK-9 and Mbr1), as measured by the Cy3-labeled secondary antibody were relatively high and similar, whereas the binding to other truncated sequences is very weak. This result provides a rationale for development of a glycan microarray to detect the presence of anti-Globo H antibody in the serum of patients. Because we only loaded 0.7 nL to each spot, the minimum amount of sugar required for detection was 10^{-15} mol per spot if a 50 μM solution is used (Table 2 and Fig. 3). For the purpose of detection only, the concentration can further be reduced to 10^{-18} mol per spot, as shown in the study in ref. 42. This is indeed a highly

Fig. 5. Ratios of IgG levels against Globo H analogs in sera from breast cancer patients and normal blood donors. The relative fluorescence ratios were obtained from fluorescence intensity of Globo H or Globo H analogs divided by fluorescence intensity of Gb5. The mean of Globo H/Gb5 IgG ratios was significantly higher in sera of breast cancer patients.

Fig. 6. Ratios of IgM levels against Globo H analogs in sera from breast cancer patients and normal blood donors. The relative fluorescence ratios were obtained from fluorescence intensity of Globo H or Globo H analogs/fluorescence intensity of Gb5. The mean of Globo H/Gb5 IgM ratios was significantly higher in sera of breast cancer patients.

Fig. 7. The comparison of glycan microarray and ELISA for monitoring anti-Globo H response. Sera from immunized mice were serially diluted and analyzed for anti-Globo H specific IgG antibody on glycan microarray and ELISA plates. The fold level of signal over background was calculated as (mean of fluorescence of postserum – preserum)/background intensity in glycan microarray or (OD value of postserum – preserum)/background in ELISA. The antibody levels were detectable up to 1:1,920 dilution method (3.2 ± 1.0 -fold of background signal), but only 1:240 dilution in ELISA (1.08 ± 0.48 -fold of background OD). Values shown were mean $\pm$ SD of four mice.

the analysis of fucosyltransferase activity and identification of fucosyltransferase inhibitors (21) as possible therapeutics.

Materials and Methods

General. NHS-coated glass slides were purchased from Nexterion H slide (SCHOTT North America). Primary antibodies used were mouse anti-Globo H monoclonal antibodies Mbr1 (IgM; Alexis Biochemicals), VK-9 [IgG; a gift from Philip Livingston (Memorial Sloan-Kettering Cancer Center, New York)], and A488 anti-mouse/human SSEA-3 (stage-specific embryonic antigen-3) (eBioscience). The secondary antibodies used were Cy3-conjugated Goat anti-mouse IgG, goat anti-mouse IgM, goat anti-human IgG, and goat anti-human IgM (Jackson ImmunoResearch). Human plasma samples from healthy individuals and cancer patients were obtained from the Genomics Research Center, Academia Sinica; and the Tri-Service General Hospital, Taipei, Taiwan, respectively. Samples were fully encoded to protect patient confidentiality and were used under a protocol approved by the Institutional Review Board of Human Subjects Research Ethics Committee of Academia Sinica, Taipei, Taiwan. Globo H and its truncated analogs were synthesized as in ref. 38.

Glycan Microarray Fabrication. Microarrays were printed (BioDot; Cartesian Technologies) by robotic pin (SMP3; TeleChem International) deposition of ~ 0.7 nL of various concentrations of amine-containing glycans in printing buffer [300 mM phosphate buffer (pH 8.5), containing 0.005% Tween-20] from a 96-well microtiter plate onto NHS-coated glass slides. The slides for Globo H analogs were spotted with solutions of Globo H analogs 1–8 with concentrations 50 μ M every two rows for one glycan from bottom to top with 15 replicates horizontally placed in each subarray, and each slide was designed for 16 grids for further incubation experiments. Printed slides were allowed to react in an atmosphere of 80% humidity for an hour followed by desiccation overnight. These slides were designed for 16 grids in one slide, and stored at room temperature in a desiccator until use. Before the binding assay, these slides were blocked with ethanolamine [50 mM ethanolamine in 50 mM borate buffer (pH 9.2)] and then washed with water and PBS buffer (pH 7.4) twice.

Antibody Binding Assay. Mbr1, a mouse IgM anti-Globo H monoclonal antibody, VK-9, a mouse IgG anti-Globo H monoclonal antibody, or A488 anti-mouse/human SSEA-3 were prepared in 0.05% Tween 20/PBS buffer (pH 7.4) and added to cover the grid with application of a coverlip. After incubation in a humidifying chamber with shaking for 1 h, the slides were washed three times each with 0.05% Tween 20/PBS buffer (pH 7.4), PBS buffer (pH 7.4), and water. Next, Cy3-conjugated goat anti-mouse IgM (for Mbr1) or IgG (for VK-9) antibody was added to the slide and incubated in a humidifying chamber with shaking under cover lip for 1 h. The slide was washed three times each with 0.05% Tween20/PBS buffer (pH 7.4), PBS buffer (pH 7.4), and H₂O and dried. The slide was scanned at 532 nm (for Cy3-conjugated secondary antibody) and 488 nm (for A488 anti-SSEA-3 antigen antibody) with a microarray fluorescence chip reader (ArrayWorx microarray reader).

Table 5. The comparison of sensitivity of glycan microarray and ELISA

Dilution	Increase fold of signal	
	Glycan microarray	ELISA
30	374.7 $\pm$ 87.83	4.01 $\pm$ 1.58
120	188.4 $\pm$ 78.93	1.92 $\pm$ 0.75
240	102.2 $\pm$ 44.21	1.08 $\pm$ 0.48
480	44.86 $\pm$ 17.05	0.20 $\pm$ 0.10
960	12.13 $\pm$ 4.08	0.30 $\pm$ 0.14
1,920	3.203 $\pm$ 1.048	ND

Glycan microarray showed much higher sensitivity than ELISA. The increase fold of signal defined as (signal intensity of post-immune – pre-immune)/background intensity. Values are mean $\pm$ SEM. ND, not detectable.

Microarray Analysis of Cancer Patient Plasma. The plasma samples from breast cancer patients and healthy individuals were diluted 1:20 with 0.05% Tween 20/3% BSA/PBS buffer (pH 7.4), and applied to the grids on the Globo H analogs microarrays and then incubated in a humidifying chamber with shaking for 1 h. Then the slides were washed three times each with 0.05% Tween 20/PBS buffer (pH 7.4), PBS buffer (pH 7.4), and water. Next, Cy3-conjugated goat anti-human IgM or IgG antibody was added to the slide as described above and incubated in a humidifying chamber incubation with shaking under a coverlip for 1 h. The slide was washed three times each with 0.05% Tween20/PBS buffer (pH 7.4), PBS buffer (pH 7.4), and H₂O and dried. The slide was scanned at 532 nm (for Cy3-conjugated secondary antibody) with a microarray fluorescence chip reader (ArrayWorx microarray reader).

Immunization. Groups of three mice (6-week-old female C57BL/6 mice; BioLASCO) were immunized s.c. with Globo H-KLH (Optimer) once weekly for 3 weeks. Each vaccination contained 0.6 μ g of Globo H with 2 μ g of α -galactosylceramide as an adjuvant. Control mice were injected with PBS. The mouse sera were obtained before 1st immunization and 10 days after the third immunization. The serological responses were analyzed by glycan microarray with the same method as described for human plasma except that Cy3-conjugated goat anti-mouse IgG antibody was used as a secondary antibody. 0.2 μ g of Globo H-ceramide in 100 μ L of carbonate bicarbonate buffer (pH 10) were coated in 96-well plate (NUNC) at 4°C for overnight, washed with PBS, and blocked with 3% BSA for 30 min at room temperature. Serial dilutions of mice sera were added into each well and incubated for 1 h at room temperature, followed by washing with Dulbecco's phosphate buffered saline (DPBST) and 0.05% Tween 20. Goat anti-mouse IgG-AP (1:200, Southern Biotech.) was added and incubated for 45 min at room temperature. The plates were washed with PBST five times and then incubated with an alkaline phosphates substrate, p-nitrophenyl phosphate (Sigma), for 8 min at 37°C. After incubation, the reaction was stopped by adding 3 M NaOH, and the plates were read at 405 nm on the ELISA reader (SpectraMax; Molecular Devices).

Data Analysis. The software GenePix Pro (Axon Instruments) was used for the fluorescence analysis of the extracted data. The local background was subtracted from the signal at each antibody spot. The spots with obvious defects, no detectable signal, or a net fluorescence of <100 were removed from the analysis. The “medians of ratios” from replicate spots were averaged in the same array.

To determine the $K_{D,surf}$ value, the equilibrium binding data were analyzed by fitting the data to the appropriate equation (1), assuming that ligands bound to one or two independent sites, using the commercial nonlinear regression program GradPad PRISM (GraphPad). To obtain the relative binding intensities of Globo H analogs in human plasma, we used the binding intensity to Bg5 as 100%, and normalized the relative binding intensities of Globo H analogs in each plasma sample. For example, the ratio of GH/Gb5 in sera was calculated by dividing mean of Globo H replicates by mean of Gb5 replicates. Finally, the statistical analysis of IgG or IgM levels in breast cancer patients and normal individuals was performed with an unpaired t test, using the program GraphPad Prism (GraphPad).

ACKNOWLEDGMENTS. We thank Ms. Pei-Lan Hsu, a registered nurse in the management of clinical samples. This work was supported by Academia Sinica.

1. Kannagi R, et al. (1983) New globo series glycosphingolipids in human tetraocarcinoma reactive with the monoclonal-antibody directed to a developmentally regulated antigen, stage-specific embryonic antigen-3. *J Biol Chem* 258:8934–8942.

2. Zhang SL, et al. (1997) Selection of tumor antigens as targets for immune attack using immunohistochemistry. 1. Focus on gangliosides. *Int J Cancer* 73: 42–49.

3. Hakomori S, Zhang YM (1997) Glycosphingolipid antigens and cancer therapy. *Chem Biol* 4:97–104.
4. Dube DH, Bertozzi CR (2005) Glycans in cancer and inflammation. Potential for therapeutics and diagnostics. *Nat Rev Drug Discov* 4:477–488.
5. Menard S, Tagliabue E, Canevari S, Fossati G, Colnaghi MI (1983) Generation of monoclonal-antibodies reacting with normal and cancer-cells of human-breast. *Cancer Res* 43:1295–1300.
6. Livingston PO (1995) Augmenting the immunogenicity of carbohydrate tumor antigens. *Cancer Biol* 6:357–366.
7. Canevari S, Fossati G, Balsari A, Sonnino S, Colnaghi MI (1983) Immunochemical analysis of the determinant recognized by a monoclonal-antibody (Mbr1) which specifically binds to human mammary epithelial-cells. *Cancer Res* 43:1301–1305, 1983.
8. Bremer EG, et al. (1984) Characterization of a glycosphingolipid antigen defined by the monoclonal-antibody Mbr1 expressed in normal and neoplastic epithelial-cells of human mammary-gland. *J Biol Chem* 259:14773–14777.
9. Kudryashov V, et al. (1998) Characterization of a mouse monoclonal IgG3 antibody to the tumor-associated globo H structure produced by immunization with a synthetic glycoconjugate. *Glycoconjugate J* 15:243–249.
10. Zhang S, et al. (1998) Expression of potential target antigens for immunotherapy on primary and metastatic prostate cancers. *Clin Cancer Res* 4:295–302.
11. Park S, Shin I (2002) Fabrication of carbohydrate chips for studying protein-carbohydrate interactions. *Angew Chem Int Ed* 41:3180–3182.
12. Houseman BT, Mrksich M (2002) Carbohydrate arrays for the evaluation of protein binding and enzymatic modification. *Chem Biol* 9:443–454.
13. Fazio F, Bryan MC, Blixt O, Paulson JC, Wong CH (2002) Synthesis of sugar arrays in microtiter plate. *J Am Chem Soc* 124:14397–14402.
14. Bryan MC, et al. (2002) Saccharide display on microtiter plates. *Chem Biol* 9:713–720.
15. Fukui S, Feizi T, Galustian C, Lawson AM, Chai W (2002) Oligosaccharide microarrays for high-throughput detection and specificity assignments of carbohydrate-protein interactions. *Nat Biotechnol* 20:1011–1017.
16. Wang DN, Liu SY, Trummer BJ, Deng C, Wang AL (2002) Carbohydrate microarrays for the recognition of cross-reactive molecular markers of microbes and host cells. *Nat Biotechnol* 20:275–281.
17. Mellet CO, Garcia Fernandez JM (2002) Carbohydrate microarrays. *ChemBiochem* 3:819–822.
18. Feizi T, Fazio F, Chai W, Wong CH (2003) Carbohydrate microarrays-a new set of technologies at the frontiers of glycomics. *Curr Opin Struct Biol* 13:637–645.
19. Disney MD, Magnet S, Blanchard JS, Seeberger PH (2004) Aminoglycoside microarrays to study antibiotic resistance. *Angew Chem Int Ed* 43:1591–1594.
20. Ratner DM, et al. (2004) Probing protein-carbohydrate interactions with microarrays of synthetic oligosaccharides. *ChemBioChem* 5:379–382.
21. Bryan MC, et al. (2004) Covalent display of oligosaccharide arrays in microtiter plates. *J Am Chem Soc* 126:8640–8641.
22. Disney MD, Seeberger PH (2004) Aminoglycoside microarrays to explore interactions of antibiotics with RNAs and proteins. *Chem Eur J* 10:3308–3314.
23. Adams EW, et al. (2004) Oligosaccharide and glycoprotein Microarrays as tools in HIV glycobiology: Glycan-dependent gp120/protein interactions. *Chem Biol* 11:875–881.
24. Shin I, Park S, Lee MR (2005) Carbohydrate microarrays: An advanced technology for functional studies of glycans. *Chem Eur J* 11:2894–2901.
25. Bochner BS, et al. (2005) Glycan array screening reveals a candidate ligand for Siglect-8. *J Biol Chem* 280:4307–4312.
26. Calarese DA, et al. (2005) Dissection of the carbohydrate specificity of the broadly neutralizing anti-HIV-1 antibody 2G12. *Proc Natl Acad Sci USA* 102:13372–13377.
27. Manimala JC, Roach TA, Li Z, Gildersleeve JC (2006) High-Throughput carbohydrate microarray analysis of 24 lectins. *Angew Chem Int Ed* 45:3507–3610.
28. Dotan N, Altstock RT, Schwarz M, Dukler A (2006) Anti-glycan antibodies as biomarkers for diagnosis and prognosis. *Lupus* 15:442–450.
29. Lee JC, Wu CY, Apon JV, Siuzdak G, Wong CH (2006) Reactivity-based one-pot synthesis of the tumor-associated antigen N3 minor octasaccharide for the development of a photo cleavable DIOS-MS sugar array. *Angew Chem Int Ed* 45:2753–2757.
30. Lawrie CH, et al. (2006) Cancer-associated carbohydrate identification in Hodgkin's lymphoma by carbohydrate array profiling. *Int J Cancer* 118:3161–3166.
31. Paulson JC, Blixt O, Collins BE (2006) *Nat Chem Biol* 2:238–248.
32. Stevens J, et al. (2006) Structure and receptor specificity of the hemagglutinin from an H5N1 influenza virus. *Science* 312:404–410.
33. Stevens J, Blixt O, Paulson JC, Wilson IA (2006) Glycan microarray technologies: Tools to survey host specificity of influenza viruses. *Nat Rev Microbiol* 4:857–864.
34. Stevens J, et al. (2006) Glycan microarray analysis of the hemagglutinins from modern and pandemic influenza viruses reveals different receptor specificities. *J Mol Biol* 355:1143–1155.
35. Manimala JC, Roach TA, Li ZT, Gildersleeve JC (2006) High-throughput carbohydrate microarray profiling of 27 antibodies demonstrates widespread specificity problems. *Glycobiology* 17:17C–23C, 2007.
36. Ratner DM, Seeberger PH (2007) Carbohydrate microarrays as tools in HIV glycobiology. *Curr Pharm Des* 13:173–183.
37. Mammen M, Choi SK, Whitesides GM (1998) Polyvalent interactions in biological systems: Implications for design and use of multivalent ligands and inhibitors. *Angew Chem Int Ed* 37:2755–2794.
38. Hung CY, et al. (2006) Carbohydrate microarray for profiling the antibodies interacting with Globo H tumor antigen. *Proc Natl Acad Sci USA* 103:15–20.
39. Blixt O, et al. (2004) Printed covalent glycan array for ligand profiling of diverse glycan binding proteins. *Proc Natl Acad Sci USA* 101:17033–17038.
40. Jones RB, Gordus A, Krall JA, MacBeath G (2006) A quantitative protein interaction network for the ErbB receptors using protein microarrays. *Nature* 439:168–174.
41. Gordus A, MacBeath G (2006) Circumventing the problems caused by protein diversity in microarrays: Implications for protein interaction networks. *J Am Chem Soc* (2006) 28:13668–13669.
42. Liang PH, Wang SK, Wong CH (2007) Quantitative analysis of carbohydrate-protein interactions using glycan microarrays: Determination of surface and solution dissociation constants. *J Am Chem Soc* 129:11177–11184.
43. Boyle P (1997) Global burden of cancer. *Lancet* 349:S23–S26.
44. Galanina OE, Mecklenburg M, Nifantiev NE, Pazynina GV, Bovin NV (2003) GlycoChip: Multiaarray for the study of carbohydrate-binding proteins. *Lab Chip* 3:260–265.
45. Chang WW, et al. (2008) Expression of Globo H and SSEA3 in breast cancer stem cells and the involvement of fucosyl transferase 1 & 2 in Globo H synthesis. *Proc Natl Acad Sci USA* 105:11667–11672.