Published in final edited form as:

Carbohydr Res. 2008 October 13; 343(15): 2701–2705. doi:10.1016/j.carres.2008.06.006.

The structure of the carbohydrate backbone of the LPS from *Shewanella* spp. MR-4

Evgeny Vinogradov^{*,1}, Joanna Kubler-Kielb², and Anton Korenevsky³

¹Institute for Biological Sciences, National Research Council, Ottawa ON, K1A 0R6 Canada

²National Institute of Child Health and Human Development, National Institutes of Health, 9000 Rockville Pike, Bethesda, MD, 20892, USA

³Department of Microbiology, College of Biological Science, University of Guelph, Guelph ON, N1G 2W1 Canada

Abstract

The rough type lipopolysaccharide isolated from *Shewanella* spp. strain MR-4 was analyzed using NMR, mass spectroscopy and chemical methods. Two structural variants have been found, both contained 8-amino-3,8-dideoxy-D-manno-octulosonic acid and lacked L-glycero-D-manno-heptose. A minor variant of the LPS contained phosphoramidate substituent.

Shewanella spp. strain MR-4, a Gram-negative bacterium isolated from the Black Sea, is an important environmental microorganism participating in complex biogeochemical processes involving metal reduction.^{1,2} Surface components of this organism play an important role in its adhesion to mineral surfaces.³ *Shewanella* spp. MR-4 produces a large quantity of capsular polysaccharide (CPS), which gives the cells "smooth" appearance. However its LPS has no polymeric O-chain, belonging to the "rough-type". The CPS⁴ contains a monosaccharide similar to anthrose that makes it cross-reactive with the BclA, a glycoprotein located on the surface of *B. anthracis* spores.⁵ Here we describe the results of structural analysis of the *Shewanella* spp. MR-4 LPS.

Monosaccharide analysis of the *Shewanella* MR-4 LPS showed the presence of two major constituents: D-galactose and D-glycero-D-manno-heptose, and three minor constituents: glucosamine (GlcN), mannose (Man) and glucose (Glc). The GlcN originated from the backbone of lipid A, Man and Glc appeared in the GC analysis of alditol acetates as a result of the reduction of fructose, which was identified in the LPS-derived oligosaccharides by NMR. The absolute configuration of Gal, DDHep, and GlcN were determined as D using GLC of acetylated 2-butyl glycosides. Fatty acid analysis of the LPS revealed the presence of 14:0(3-OH), 14:0, and 12:0 fatty acids. O-Deacylated LPS contained 14:0(3-OH) acids, being attached as amides to the GlcN of lipid A.

Alkaline deacylation of the LPS from *Shewanella* spp. MR-4 gave a complex mixture of oligosaccharides that were separated by HPAEC. Only the three major products were analyzed. The structure of the oligosaccharide **1** as presented on Scheme 1 was determined by 2D NMR

*Corresponding author: Evgeny Vinogradov, Institute for Biological Sciences, National Research Council, 100 Sussex Dr., Ottawa, ON, K1A 0R6, Canada, Tel 1-613-990 0832, Fax 1-613-952 9092, email evgenii.vinogradov@nrc.ca.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

spectroscopy (Table 1). Assignment of the NMR spectra showed the presence of four monosaccharide residues: α -GlcN1P, β -GlcN4P, α -8-aminoKdo4P, and α -DDHep3P. The identity of the monosaccharides was established on the basis of proton and carbon chemical shifts and H,H vicinal coupling constants. The presence of the aminogroup at position 8 of the Kdo followed from the high field chemical shift of C-8 (44.3 ppm). All monosaccharides were phosphorylated, and the phosphate group position was determined using ^1H - ^{31}P NMR correlation spectroscopy. The sequence of monomeric components was determined using NOE (correlations B1A6w, E1C5s, E1C7s) and HMBC (H-C correlations B1A6, C2B6, E1C5) data. ESI mass spectra of oligosaccharide **1** (negative mode) contained a peak of a double charged ion corresponding to a molecular mass of 1072 Da (calculated for $\text{C}_{27}\text{H}_{53}\text{N}_3\text{O}_{33}\text{P}_4$ 1071.6 Da). The second major oligosaccharide had the same structure but without phosphate at O-4 of the 8-amino-Kdo. This was confirmed by NMR and MS analysis (data not shown).

Oligosaccharide **2**, eluted near the solvent front in HPAEC, contained the residues of β -fructofuranose and α -galactopyranose (Scheme 1), and many variants of the degradation products of the reducing-end monosaccharide (6-substituted galactose). The structure of the oligosaccharide **2**, except for the reducing end monosaccharide, was determined by NMR (Table 1). The β -fructofuranose ring size and anomeric configuration were identified by ^{13}C chemical shift comparison with the known values.⁶ Its position was determined from HMBC correlation between C-2 of Fru and H-6 of Gal H. This oligosaccharide was obtained by hydrolysis of the phosphodiester bond between Gal F and Hep E and the subsequent destruction of the Gal residues F and partially G.

Acetic acid hydrolysis of the LPS gave a mixture of mono- and oligosaccharides. They were separated by gel filtration and anion exchange chromatography. The major products were galactotriose and free galacturonic acid. Products originating from Hep-(1–5)-8-aminoKdo disaccharide were not analyzed because they gave many forms with variable Kdo degradation products at the end and different phosphorylation.

NMR analysis of the O-deacylated LPS (LPS-Hy) showed the presence of two structural forms (variants **1** and **2**). Variant **1** contained all components that are required to produce compounds **1–3** upon KOH or AcOH treatment. Thus, alkaline treatment cleaves the phosphodiester bond between Gal F and phosphate group and destroys Gal F completely and Gal G partially, resulting in oligosaccharide **2**. The rest of the molecule remains as the oligosaccharide **1**. The connection between Gal G and DDHep E via a phosphodiester bond can be traced in the ^1H - ^{31}P HMQC correlation (Fig. 1). The Kdo in the LPS-Hy was partially substituted by α -GalAP (residue Y) at O-4, which is visible in the ^1H - ^{31}P HMQC correlation spectrum. Cleavage of α -GalAP in KOH gave an analog of the oligosaccharide **1** without the phosphate group at O-4. The LPS which had phosphate group at C-4 of 8-amino-Kdo without galacturonic acid retained this phosphate and gave the oligosaccharide **1** after KOH treatment. ESI MS data of the LPS-Hy (Fig. 2) showed structural variants with HexA

(Hex₄HexA₁Hep₁HexN₂Kdo₁C14OH₂P₄, observed molecular mass of 2350.6 Da, calculated 2350 Da) and without HexA (Hex₄Hep₁HexN₂Kdo₁C14OH₂P₄, observed molecular mass of 2174.5 Da, calculated 2173.9 Da), as well as a peak at of a 2297.6 molecular mass corresponding to an additional PEtN substituent to the structure without GalA, and a minor peak of –28 Da probably due to the replacement of 14:0(3-OH) acyl groups by 12:0(3-OH) acyl groups.

The NMR spectra of the LPS-Hy contained the signals of another DDHep residue E', linked to the 8-amino-Kdo (LPS-Hy variant **2**). DDHep E' was phosphorylated at positions 2 and 3 (Fig. 1), and most interestingly the ^{31}P signal of the substituent at O-3 appeared at 16.5 ppm, which indicates the presence of a phosphoramidate group. A similar substituent has been identified recently in *Xanthomonas campestris* LPS.⁷ Unfortunately no derivatives originating

from variant **2** was identified in the LPS hydrolysates; this variant also showed no signal in the mass spectrum and thus its structure as presented here should be considered tentative.

The lipid A-core part of LPS from several *Shewanella* strains were elucidated previously^{8–12} Most of them contained 8-amino-Kdo residue in place of Kdo. The absolute stereochemistry of this monosaccharide has been confirmed recently.¹³ The structure described in the current publication gives one more example of 8-amino-Kdo containing LPS, which seems to be a characteristic feature of *Shewanella*.

1. Experimental

1.1 Bacterial strains and lipopolysaccharide isolation

Shewanella spp. strain MR-4 was obtained from K. Nealson, Jet Propulsion Laboratory, Pasadena, CA and was grown aerobically in trypticase soy broth to an optical density (OD₆₀₀) of 1.0. Cells were harvested by centrifugation and LPS was extracted by the Darveau and Hancock method using EDTA and SDS on cells broken in a French press⁴ followed by DNase, RNase and protease treatment. The extracted LPS was precipitated in ice-cold 95% (v/v) ethanol containing 0.375 M MgCl₂.

1.2. NMR spectroscopy and general methods

¹H and ¹³C NMR spectra were recorded using a Varian Inova 500 spectrometer in D₂O solutions at 25 °C with an acetone standard (2.23 ppm for ¹H and 31.5 ppm for ¹³C) using standard COSY, TOCSY (mixing time 120 ms), NOESY (mixing time 200 ms), HSQC, HMBC (100 ms long-range transfer delay), ¹H-³¹P HMQC, HMQC-TOCSY (optimized for 11 Hz coupling constant) pulse sequences. Spectra of the LPS-Hy were recorded at 60 °C in 5% fully deuterated SDS. 24% NH₄OH (10 μL) were added to the 0.6 mL D₂O sample. Spectra were assigned using the Pronto program.¹⁵ ESI mass spectra were obtained using a Micromass Quattro spectrometer in 50% MeCN with 0.2% HCOOH at a flow rate of 15 μL/min with direct injection. GLC, GLC-MS, methylation and monosaccharide analyses and preparation of the oligosaccharides were performed as described.^{8–10}

Abbreviations

LPS, lipopolysaccharide; DDHep, *D*-glycerol-*D*-manno-heptose; Kdo, 3-deoxy-*D*-manno-oct-2-ulonic acid; 8-aminoKdo, 8-amino-3,8-dideoxy-*D*-manno-oct-2-ulonic acid; *P*, phosphate; PEtN, phosphoethanolamine; HPAEC, high performance anion exchange chromatography.

Acknowledgements

The research was partially supported by a US-DOE Natural and Accelerated Bioremediation Research grant and by the Intramural Research Program of the NIH, NICHD. Authors thank J. Stupak (NRC, Canada) for the ESIMS analysis.

References

1. Nealson KH, Myers CR, Wimpee BB. Deep-Sea Res 1991;38:S907–S920.
2. Marsili E, Baron DB, Shikhare ID, Coursolle D, Gralnick JA, Bond DR. Proc. Natl. Acad. Sci U. S. A 2008;105:3968–3973. [PubMed: 18316736]
3. Korenevsky AA, Vinogradov E, Gorby Y, Beveridge TJ. Appl. Environ. Microbiol 2002;68:4653–4657. [PubMed: 12200327]
4. Vinogradov E, Nossova L, Korenevsky A, Beveridge TJ. Carbohydr. Res 2005;340:1750–1753. [PubMed: 15936738]
5. Kubler-Kielb J, Vinogradov E, Hu H, Leppla SH, Robbins JB, Schneerson R. Proc. Natl. Acad. Sci. U. S. A. 2008doi:

6. Bock K, Pedersen C. *Adv. Carb. Chem. Biochem* 1983;41:27–66.
7. Silipo A, Molinaro A, Sturiale L, Dow JM, Erbs G, Lanzetta R, Newman MA, Parrilli M. *J. Biol. Chem* 2005;280:33660–33668. [PubMed: 16048996]
8. Vinogradov E, Korenevsky A, Beveridge TJ. *Carbohydr. Res* 2002;337:1285–1289. [PubMed: 12151208]
9. Vinogradov E, Korenevsky A, Beveridge TJ. *Carbohydr Res* 2003;338:1991–1997. [PubMed: 14499575]
10. Vinogradov E, Korenevsky A, Beveridge TJ. *Carbohydr. Res* 2004;339:737–740. [PubMed: 15013415]
11. Silipo A, Leone S, Molinaro A, Sturiale L, Garozzo D, Nazarenko EL, Gorshkova RP, Ivanova EP, Lanzetta R, Parrilli M. *Eur. J. Org. Chem* 2005;2005:2281–2291.
12. Moule AL, Galbraith L, Cox AD, Wilkinson SG. *Carbohydr. Res* 2004;339:1185–1188. [PubMed: 15063209]
13. Leone S, Molinaro A, De Castro C, Baier A, Nazarenko EL, Lanzetta R, Parrilli M. *J. Nat. Prod* 2007;70:1624–1627. [PubMed: 17914882]

Figure 1.
 ^1H - ^{31}P HMQC spectrum of *Shewanella* spp. MR-4 LPS-Hy.

Figure 2.
Reconstructed ESI mass spectrum of the *Shewanella* spp. MR-4 LPS-Hy.

KOH products:

Reduced AcOH product:

LPS-Hy variant 1:

LPS-Hy variant 2:

Scheme 1.

Structures of the isolated oligosaccharides derived from the LPS from *Shewanella* spp. MR-4. PN is phosphoramidate. ac = 14:0(3-OH) acyl group. GalA Y is present in about 50% on each of the structural variants of LPS-Hy. Residue Xxx (G*) is a product of alkaline degradation of the Gal G.

Table 1
¹H and ¹³C NMR data for the isolated oligosaccharides.

Unit, compound	Atom	1	2 (3ax)	3 (3eq)	4	5	6(a)	7(a) (6b)	8a (7b)	8b
α -GlcN1P A, 1a	¹ H	5.74	3.44	3.93	3.52	4.12	3.91	4.26		
	¹³ C	93.2	55.4	70.9	71.1	73.8	70.4			
	³¹ P	2.0								
α -GlcN1P A, LPS-Hy	¹ H	5.50	3.90	3.85	3.80	3.98	3.95	4.18		
	¹³ C	95.0	55.0	71.4	70.0	72.9	69.2			
	³¹ P	0.3								
β -GlcN4P B, 1a	¹ H	4.89	3.15	3.93	3.97	3.79	3.54	3.67		
	¹³ C	100.5	56.9	72.9	75.4	75.0	63.8			
	³¹ P				3.1					
β -GlcN4P B, LPS-Hy	¹ H	4.67	3.90	3.94	3.92	3.77	3.55	3.74		
	¹³ C	103.4	56.6	74.6	74.8	75.0	64.3			
	³¹ P				2.14					
α -aminoKdo C, 1a	¹ H		2.16	2.26	4.61	4.36	3.86	4.11	3.08	3.51
	¹³ C	175.4	101.3	35.3	71.3	74.9	75.1	67.0	44.4	
	³¹ P				2.9					
α -aminoKdo C with GalA Y, LPS-Hy var. 1	¹ H		2.12	2.34	4.65	4.40	3.92	4.13	3.24	3.55
	¹³ C			35.6	71.9	75.5	75.7	67.0	44.3	
	³¹ P				-0.6					
α -aminoKdo C no GalA Y, LPS-Hy var. 1	¹ H		2.12	2.34	4.62	4.33	3.92	4.13	3.24	3.55
	¹³ C			35.6	71.1	75.2	75.7	67.0	44.3	
	³¹ P				1.4					
α -aminoKdo C' with GalA Y, LPS-Hy var. 2	¹ H		2.12	2.34	4.66	4.49	3.92	4.13	3.24	3.55
	¹³ C			35.6	71.9	75.0	75.7	67.0	44.3	
	³¹ P				-0.6					
α -aminoKdo C' no GalA Y, LPS-Hy var. 2	¹ H		2.12	2.34	4.62	4.43	3.92	4.13	3.24	3.55
	¹³ C			35.6	71.1	74.7	75.7	67.0	44.3	
	³¹ P				1.4					
α -DDHep E, 1a	¹ H	5.12	4.30	4.36	3.93	4.12	4.12	3.73	3.79	
	¹³ C	102.2	70.5	77.1	67.4	74.9	73.8	63.6		
	³¹ P				2.9					
α -DDHep E, LPS-Hy var 1	¹ H	5.11	4.32	4.47	3.92	4.07	4.04			
	¹³ C	102.2	70.9	76.2	66.5	74.6	73.2			
	³¹ P			-0.7						
α -DDHep E', LPS-Hy var 2	¹ H	5.39	4.72	4.55	4.10	4.18				
	¹³ C	99.0	76.5	80.6	66.9	74.4				
	³¹ P		-10.2	16.8						
α -Gal H, 3	¹ H	4.91	3.84	3.90	4.00	3.99	3.76	3.76		
	¹³ C	99.5	69.2	70.3	70.1	72.0	62.0			
	¹ H	4.96	3.82	3.89	3.99	4.03	3.72	3.82		
α -Gal H, 2 and LPS-Hy	¹³ C	100.2	69.9	71.0	70.9	71.4	62.4			
	¹³ C		3.87	3.91	4.07	4.25	3.69	3.89		
	³¹ P	100.8	69.2	69.9	70.1	70.7	67.9			
α -Gal G, LPS-Hy	¹³ C	5.24	3.91	4.14	4.06	4.21	3.72	3.95		
	³¹ P	94.3	69.6	70.5	71.0	70.8	68.4			
	¹ H	3.78	4.10	3.84	3.91	4.08	3.68			
Gal-ol F, 3	¹³ C	63.5	72.5	79.5	70.7	70.9	63.8			
	¹ H	5.69	4.07	3.94	4.30					
	¹³ C	97.7	70.0	73.7	65.7					
β -Fru/M, 2 and LPS-Hy	³¹ P	-0.7								
	¹ H	3.68/3.74		4.17	4.10	3.86	3.67	3.82		

Unit, compound	Atom	1	2 (3ax)	3 (3eq)	4	5	6(a)	7(a) (6b)	8a (7b)	8b
α -GalA Y, LPS Hy	^{13}C	61.7	105.2	78.2	76.0	82.6	63.8			
	^1H	5.64	3.93	4.03	4.38	4.50				
	^{13}C	96.6	69.8	70.6	72.0	74.2	176.5			
	^{31}P	-0.6								