Published in final edited form as:

J Am Chem Soc. 2006 February 22; 128(7): 2452–2462. doi:10.1021/ja0576684.

Experimental and Computational Studies on the Mechanism of *N*-Heterocycle C-H Activation by Rh(I)/PCy₃

Sean H. Wiedemann[‡], Jared C. Lewis, Jonathan A. Ellman, and Robert G. Bergman

Contribution from the Department of Chemistry, University of California, and Division of Chemical Sciences, Lawrence Berkeley National Laboratory, Berkeley, California 94720.

Abstract

Evidence is presented for a proposed mechanism of C-H activation of 3-methyl-3,4-dihydroquinazoline (**1**) by (PCy₃)₂RhCl. One intermediate (**3**), a coordination complex of **1** with (PCy₃)₂RhCl, was identified along the path to the Rh-*N*-heterocyclic carbene product of this reaction (**2**). Isotopic labeling and reaction-rate studies were used to demonstrate that C-H activation takes place intramolecularly on the reaction coordinate between **3** and **2**. Computational studies corroborate the proposed mechanism and suggest that the rate-limiting step is oxidative addition of the C-H bond to the metal center. The consequences of this mechanism for coupling reactions of *N*-heterocycles that occur via Rh-catalyzed C-H bond activation are discussed.

Introduction

Our group¹ and others² have been interested in the use of directing groups to promote and control regioselectivity in catalytic C-H additions to olefins. Murai and coworkers' method for Ru-catalyzed *ortho*-alkylation of acetophenones was the first general application of directed C-H activation toward C-C bond formation.³ Mechanistic studies carried out on that system showed that the keto group functions by coordinating to the catalyst and establishing a geometry that facilitates cyclometalation.^{4,5} Using mechanistic insight, the scope of competent directing groups was expanded to include many other functional groups that can act as Ru⁰ ligands.⁶

Contemporaneously with these investigations, it was revealed that a single atom, the ring heteroatom of an *N*-heterocycle, can perform the same function as a pendant directing group—promoting selective functionalization of an adjacent C-H bond. Since predictable methods for heterocycle elaboration are critical to the synthesis of pharmaceuticals, there is great potential benefit in understanding the mechanistic pathways unique to this substrate class. The first example of a direct catalytic coupling between a heterocycle and an olefin was reported by Jordan and coworkers in 1989.⁷ Using a zirconocene catalyst, the C6-H bond of 2-picoline was added to propene in high yield. Despite a thorough investigation of that reaction's mechanism, its substrate scope could not be significantly enlarged.⁸ In 1992, Moore and coworkers reported heterocycle/olefin coupling with the incorporation of CO.⁹ The development of this reaction by other investigators led to limited gains in terms of scope and utility; direct mechanistic information remains scarce because metal-carbonyl clusters are probable intermediates.¹⁰

Correspondence to: Jonathan A. Ellman; Robert G. Bergman, bergman@cchem.berkeley.edu.

[‡]Present address: Graduate School of Pharmaceutical Sciences, University of Tokyo, Hongo 7-3-1, Bunkyo-ku, Tokyo 113-0033, Japan

Olefin/heterocycle coupling has made a resurgence recently with the advent of potent electrophilic catalysts. Intramolecular coupling reactions targeting the electronically activated C-H bonds of indole, for example, have been demonstrated for Ru,¹¹ Pt,¹² and an organo-catalyst.¹³ Still, the direct coupling of diverse and cheap alpha-olefins to minimally activated heterocycles remains a tantalizingly underdeveloped transformation.¹⁴

In 2001, we described a Rh(I)/phosphine catalyst system capable of promoting the C-H coupling of *N*-heterocycles to olefins.¹⁵ Initially, azoles and benzazoles were shown to be efficiently activated.^{15a} The substrate specificity of this reaction was illuminated by a mechanistic study of one benzimidazole derivative.^{15b} Remarkably, in the principal catalyst resting state, the benzimidazole substrate has undergone rearrangement, the resulting product binding to rhodium as an *N*-heterocyclic carbene (NHC) ligand.

With a Rh-NHC complex as the starting point, rate studies and DFT calculations were carried out and used as the basis for a proposed mechanism of Rh-catalyzed heterocycle/alkene coupling (Scheme 1). Subsequent reaction development focused on heterocycles capable of forming and stabilizing an NHC tautomer. Using insight from mechanistic study, the substrate scope of Rh-catalyzed olefin coupling was expanded to include azolines¹⁶ and, more recently, dihydroquinazolines¹⁷ (eq 1).

Yet, portions of the proposed mechanism were still not fully understood. In fact, the most remarkable step, the formation of a Rh-NHC complex from heterocycle and Rh(I) via H-migration, is without literature precedent. Being a unique C-H bond activation process and a mild method for the preparation of NHC ligands, we felt that a deeper understanding of this fundamental transformation could lead to new opportunities for reaction discovery. The results of experimental and computational studies of Rh(I) mediated heterocycle C-H activation reactions are described in this report. On the basis of these studies a detailed mechanism for C-H activation is presented and evaluated in the context of catalysis.

Results

Structural studies

Following some initial work, 3-methyl-3,4-dihydroquinazoline (**1**) was chosen from among the heterocycles active toward Rh-catalyzed C-H functionalization as our platform for mechanistic study. It reacted cleanly with stoichiometric amounts of the catalyst mixture used in olefin coupling reactions, [(coe)₂RhCl]₂/PCy₃ (coe = *cis*-cyclooctene, Cy = cyclohexyl), to give carbene complex **2** (eq 2).¹⁸ These conditions were previously used to synthesize an analogous NHC-complex from *N*-methylbenzimidazole.¹⁹ As in that case, the structure of **2** was elucidated by single crystal X-ray analysis (Figure 1).

The irreversible conversion of **1** to **2** occurs between 45–75 °C, well below the temperature required for olefin coupling (150 °C). At temperatures below 35 °C, carbene formation is sluggish and an orange precipitate forms over 1–3 d. Based on ¹H and ³¹P NMR analysis, this precipitate (**3**) was found to be a new Rh-complex of **1**. Further *in situ* NMR study of carbene formation reactions revealed the presence of **3** at a variety of temperatures. Importantly, throughout all reactions, the mass balance of added **1** could be accounted for as either free **1**, carbene **2**, or complex **3**.

The structure of **3** was assigned to be a *trans*-2:1 PCy₃/**1** coordination complex of Rh(I)Cl based on peak integration in ¹H NMR spectra and peak splitting in ³¹P NMR spectra. Single crystal X-ray analysis of **3** was not possible because, regardless of solvent system, **3** crystallized with too much interstitial solvent to survive even gentle handling. NMR studies of isotopically labeled materials, on the other hand, did provide details about the solution structure of **3**. Using a convergent three-step procedure, ¹⁵N**1-1** was prepared from ¹⁵N-benzamide and used to synthesize ¹⁵N**1-3** (Scheme 2). The magnitude of ¹⁵N-³¹P coupling observed in the ³¹P spectrum of ¹⁵N**1-3** was consistent with a Rh(I) complex bearing ¹⁵N-heteroarene and phosphine ligands in a *cis* orientation (3.0 Hz).²⁰ Given this strong evidence that **1** binds to rhodium through N1, rather than N3, we proposed the structure of **3** depicted in Scheme 2.

In order to confirm that the heterocycle was indeed σ-bound to rhodium we synthesized and formed the Rh(I)-complex of ¹⁵N**1**, ¹³C**2-1** (Scheme 2). The ¹³C chemical shift change of the amidine carbon upon complexation was found to be small and upfield (Δδ = +5.1 ppm). Gladysz and others have shown that small chemical shift changes are expected for σ-complexes of carbonyls to electron-rich late metals, whereas large downfield shift changes (Δδ = −90–140 ppm) are observed for the corresponding π-complexes.²¹

With confirmation of σ-bonding in **3** in hand, DFT calculations were performed on a model system (using PMe₃ in place of PCy₃, **5**) in order to obtain more detailed structural information. After energy minimization, **5** adopted a calculated geometry that can be described as an intersection of two orthogonal molecular planes, where one plane is made up of the dihydroquinazoline framework and the other is defined by the Rh-square plane (Figure 2).²² In this conformation, the C8-H atom of bound **1** resides 2.7 Å above the square-planar rhodium center. At that distance, the M-H close contact falls into the upper range of “pre-agostic” interactions.²³ If the true geometry of **3** in solution were analogous to the calculated geometry of **5** in the gas phase, then the resulting pre-agostic interaction would cause the ¹H NMR resonance for C8-H to be strongly shifted downfield. Indeed, upon complexation of **1** to (PCy₃)₂RhCl, a single Ar-H resonance attributable to C8-H was observed 3 ppm downfield of the others (δ = 10 vs. 6.9 ppm) supporting the structure of **3** depicted in Figure 2.²⁴

Reactivity Studies

Before proceeding further it became necessary to ascertain the role of **3** in the formation of **2**. Two C-H activation reactions were monitored concurrently at 60 °C: one starting from isolated **3**, and the other starting from **1** and [(coe)₂RhCl]₂/2PCy₃. In both of these reactions an observable equilibrium between **1** and **3** was rapidly established. From this maximum point, the concentration of **3** dropped continuously to zero as **1** was consumed to form **2**. Thus, either **3** is a reactive intermediate lying on the path to **2** (Scheme 3, Path A), or the equilibrium between **1** and **3** is a kinetic “dead end”, with C-H activation arising from a separate bimolecular pathway (Scheme 3, Path B).

This sort of uncertainty is typically resolved by comparing the rate that product is formed by the reaction of starting materials to the rate that product is formed by subjecting the isolated intermediate to the same reaction conditions. Unfortunately, that exact experiment is not possible in the present case, because it relies on k₁/k_{−1} equilibration being sufficiently slow.

At the temperatures required to dissolve **3** for NMR reaction analysis (>45 °C) there is rapid equilibration between **1** and **3**. Once equilibrium is established, Path A and Path B become kinetically indistinguishable—by applying the equilibrium approximation, the rate expression for either path can be rewritten so that it differs from the rate expression for the other path only in the way that k_{obs} is defined (eq 3).

$$d[\mathbf{2}]/dt (\text{Path A}) = k_2[\mathbf{3}] \quad (3a)$$

$$d[\mathbf{2}]/dt (\text{Path B}) = k_2'[\mathbf{1}][(\text{PCy}_3)_2\text{RhCl}] \quad (3b)$$

$$K_{\text{eq}} = [\mathbf{3}]/([\mathbf{1}][(\text{PCy}_3)_2\text{RhCl}]) \quad (3c)$$

$$d[\mathbf{2}]/dt (\text{Path B}) = (k_2'/K_{\text{eq}})[\mathbf{3}] = k_{\text{obs}}[\mathbf{3}] = k_2[\mathbf{3}] = d[\mathbf{2}]/dt (\text{Path A}) \quad (3d)$$

Eventually, the two reaction paths were discriminated by monitoring a reaction between **1**, $[(\text{coe})_2\text{RhCl}]_2$ and PCy_3 at a temperature low enough for **1** and **3** to slowly approach equilibrium, yet high enough for the rate of formation of **2** ($d[\mathbf{2}]/dt$) to be accurately measured. Accordingly, at 37 °C the concentrations of **1**, **2**, and **3** were measured continuously as a function of time using *in situ* ^1H NMR analysis (Figure 3). If Path A were operative, $d[\mathbf{2}]/dt$ would be expected to increase over time as $[\mathbf{3}]$ increases, resulting in an upward curving graph of $[\mathbf{2}]$ vs. time. In contrast, Path B predicts that as $[\mathbf{1}]$ and $[(\text{PCy}_3)_2\text{RhCl}]$ are depleted to form **3**, $d[\mathbf{2}]/dt$ should decrease, resulting in a downward curving graph of $[\mathbf{2}]$ vs. time. The observed plot of $[\mathbf{2}]$ vs. time showed slight upward curvature at early conversion, when the approach to equilibrium was most rapid, providing support for Path A over Path B (Figure 4). To further confirm this qualitative result, the observed concentration vs. time profiles were fitted to the mechanistic pathways illustrated in Scheme 3 with the aid of the Gepasi kinetics modeling program.²⁵ The simulated $[\mathbf{2}]$ vs. time profile generated from Path A matched experimental data far better than the profile derived from Path B (Figure 4).

Given that **3** is an intermediate on the pathway to **2**, and has the same stoichiometry as **2**, the simplest plausible mechanism for C-H activation would seem to be intramolecular H-transfer. Such a mechanism would lead to a first-order dependence of $d[\mathbf{2}]/dt$ on the concentration of **3**. The simulated reaction profiles generated for Path A by kinetics modeling software (Figure 3, smooth curves), presuppose first-order dependence on **3** and precisely fit observed data. While this does constitute evidence in favor of an intramolecular mechanism, goodness of fit is generally regarded as a weak measure of reaction order. We posed two crucial questions, therefore, in order to clarify the role of bimolecular processes in the seemingly intramolecular conversion of **3** to **2** via Path A: (1) Is the H removed from C2 the same atom as that added to N1? and if so, (2) Does that H exchange with any others as it is transferred? A deuterium tracer experiment was performed to shed light on these questions.

C2-D-1 reacted with the usual $\text{Rh(I)}/\text{PCy}_3$ mixture to produce **N1-D-2** with 88% isotopic incorporation at the N1 position (eq 4). Some minor deuterium loss can be explained by invoking transient Rh-D intermediates. If such intermediates underwent reversible cyclometalation, a well-known process for $\text{Rh(I)}/\text{PCy}_3$ complexes,²⁶ deuterium exchange into phosphine cyclohexyl groups could occur (Scheme 4). Because rhodium hydride intermediates were indeed located computationally (vide infra) and a reaction pathway involving the dissociation of PCy_3 was recognized experimentally (vide infra) this mechanism seems reasonable. Moreover, the extent of deuterium loss is small, and thus does not jeopardize the conclusion that the C2-H of **1** is the primary source of N1-H in **2**.²⁷

(4)

In an effort to answer the question of whether H-transfer from C2 to N1 occurs directly or with intermolecular exchange, a double-labeling crossover experiment was carried out using equal amounts of C2-D-1 and ¹⁵N1-1 (Table 1). The amount of observable crossover product (¹⁴N1-H-2, **2**_{cross}) produced in this experiment was no greater than the amount expected to arise from minor deuterium loss via the pathway discussed above. Furthermore, the fact that percent crossover stayed nearly constant at first, and then slowly rose over long reaction times suggests that some of the observed crossover product may have arisen from isotopic scrambling between NHC products after they were formed. Hence, we conclude that the conversion of **3** to **2** results from intramolecular H-transfer,²⁸ and rule out mechanisms involving the exchange of hydrides between metal centers or the reductive elimination and recapture of hydrides as HCl.²⁹

Rate studies

Assuming that C-H activation takes place via intramolecular H-transfer, according to the mechanism presented in Path A of Scheme 3, an overall rate law can be derived for the conversion of **1** to **2** (eq 5). By applying this expression, d[**2**]/dt and the concentrations of **1** and (PCy₃)₂RhCl can be used to calculate k₂, which provides information about the C-H activation step in isolation. Unfortunately, it is difficult to ascertain the concentration of (PCy₃)₂RhCl necessary for this calculation because, (1) (PCy₃)₂RhCl lacks diagnostic resonances for quantification by ¹H NMR, (2) it exists in equilibrium with its dimer and other complexes (Scheme 5),³⁰ and (3) it undergoes decomposition over the course of the carbene formation reaction.³¹ In addition to causing uncertainty in the concentration of “active Rh” the tendency of (PCy₃)₂RhCl toward decomposition readily explains the observed reaction yields of not greater than 85% for 1:1 reactions between **1** and Rh(I)/PCy₃.³² Thus, in order to compute k₂ given the three quantities measurable by ¹H NMR analysis ([**1**], [**3**], [**2**]), kinetics simulation software was used. By allowing the software to freely vary [(PCy₃)₂RhCl] and the rate constants defined in Path A of Scheme 3, good fits to empirical data were achieved (relative errors for k₂ < 1%).³³

$$d[\mathbf{2}]/dt = k_2[\mathbf{3}] = K_{eq}k_2[\mathbf{1}][(\text{PCy}_3)_2\text{RhCl}] \quad (5)$$

In order to obtain activation parameters for C-H activation, a series of reactions, starting from isolated **3**, were monitored continuously by *in situ* ¹H NMR analysis over a range of temperatures (45–75 °C). In addition to values for k₂, kinetics simulations provided values for K_{eq}. From these values a Van't Hoff plot was constructed and fitted to a linear function (Figure S-1c). Probably as a result of the aforementioned uncertainty in [(PCy₃)₂RhCl], there was considerable scatter of data about the best-fit Van't Hoff line (r² = 0.82). Nevertheless, the slope and intercept of the fitting function did correspond to thermodynamic parameters that are consistent with reversible binding of a neutral ligand (ΔH° = −5.9 ± 0.8 kcal/mol and ΔS° = −10 ± 2 cal/mol·K).

Initially, when data collected for k₂ were transformed into an Eyring plot, deviation from linear behavior was once again observed. This imprecision was puzzling to us because the parameters governing k₂, [**2**] and [**3**], are accurately measurable. In an effort to trace the source of what seemed to be systematic error, we hypothesized that C-H activation might proceed through a

rate-limiting transition structure with only one bound PCy₃. In such a sequence, rapid, post-rate-limiting recoordination of free PCy₃ would complete the conversion to **2** (eq 6). This mechanism predicts an inverse dependence of d[**2**]/dt on [free PCy₃]. Systematic error would thus occur if small but differing amounts of free PCy₃ were introduced, for example, from different batches of reaction starting material (**3**).

(6)

In agreement with our hypothesis, we found that a small amount of added PCy₃ (13 mM = 0.5 equiv vs. [Rh]_T) generally did lead to a 2-2.5 fold reduction in the rate of carbene formation. However, for [free PCy₃] > 13 mM the rate of carbene formation was found to be independent of [PCy₃] (Figure 5). Taken together, these data reveal the existence of two competing reaction pathways between **3** and **2**—one that is dissociative in phosphine and one that is not.

Between the two operative pathways, the dissociative one is the more difficult to study due to its extreme sensitivity to free PCy₃. We elected to study the non-dissociative pathway instead. In subsequent studies, the dissociative pathway was effectively suppressed by the addition of a small amount of excess PCy₃ (13 mM) to all reaction solutions.³⁴ The deuterium kinetic isotope effect on *k*₂ was measured under these conditions. The observed *k*_H/*k*_D (1.8 ± 0.1) is consistent with cleavage of the C2-H bond during or prior to the rate determining step. When the Eyring plot of *k*₂ was redrawn using data acquired from reactions employing excess PCy₃, a very precise linear fit was obtained (*r*² = 0.9995) (Figure 6). The resulting activation parameters (Δ*H*[‡] = 26.0 ± 0.3 kcal/mol and Δ*S*[‡] = -10.3 ± 0.8 cal/mol·K) reveal the dramatic extent to which metal-mediation facilitates heterocycle to NHC tautomerization.³⁵

Computational Studies

In an effort to gain further insight into the microscopic steps of C-H activation, our experimental mechanistic data were augmented with DFT calculations.³⁶ Aided by ample structural information about **3** and **2**, a model system, using 3-methyl-3,4-dihydropyrimidine in place of **1** and PMe₃ in place of PCy₃, was constructed. After energy minimization, all bond distances common to both isolated carbene complex **2** and model carbene complex **I** differed by no more than 5%, thus ensuring the applicability of the chosen basis set and level of theory (B3LYP, LACVP**++). Starting complex **A**, the structure of which is consistent with all spectroscopic observables for **3**,³⁷ was connected to **I** via two possible non-dissociative, intramolecular H-transfer reaction coordinates differing from one another in the manner by which they effect spatial interchange of the reacting Rh and H atoms (Figure 7).

Both pathways initially pass through transition state **B**, which involves migration of the Rh-center toward the C2-H bond, and into minimized C-H σ-complex **C**. C-H σ-complexes have generally been proposed and identified along reaction coordinates for C-H activation³⁸ and have been specifically located along calculated pathways for the C-H oxidative addition of azoliums to (PMe₃)₂RhCl and IrH₃(PMe₃)₂.³⁶ Consistent with these reports, we were readily able to locate a transition state for C-H oxidative addition (**D**) originating from the C-H σ-bound intermediate **C**. Contrary to our expectations, the resultant minimized anti-hydride complex (**E**) could not be connected to carbene **I**. Intramolecular H-transfer from Rh to N3 is prohibited due to the anti orientation of N and H atoms about C2-Rh bond of **E**. Nor could a transition state be located to connect **C** to a different hydride complex with a more favorable N-C-Rh-H dihedral angle (**G**), presumably due to improper orientation of the Rh-center with

respect to the C2-H bond. Nevertheless, we were able to determine a productive reaction coordinate from anti-hydride complex **E** by recognizing that a 180° C-Rh bond rotation would bring N3 and the hydride into alignment for intramolecular H-transfer. A corresponding rotation transition state (**F**, 20 kcal/mol above **E**) and the desired *syn*-hydride (**G**) were successfully minimized. Finally, the transition state for H-transfer between Rh and N3 was located along the path from **G** to the final product, **I**. To our knowledge, this result constitutes the first instance of NHC formation by β -hydride insertion.

A second path to **I** was identified diverging from the “rotation” pathway at the C-H σ -complex **C**. Instead of ascending to the high-energy transition state for C-H oxidative addition, **C** was shown to undergo a low-energy Rh-migration to a second C-H σ -complex (**K**) via transition state **J**. During the migration, the Rh-center moves about the C2-H bond to a position closer N3 than N1. A transition state for oxidative addition (**L**) was located from C-H σ -complex **K**. Just as the geometry of **K** is inverse of **C**, the geometry of the resulting hydrides were also found to be inverse. Transition state **L** provides *syn* hydride **G** directly. The remainder of the reaction coordinate for the “migration” pathway is identical to that described above for the “rotation” pathway.

These mechanisms provide two possible microscopic reaction coordinates for the Rh-mediated intramolecular H-transfer leading from coordination complex **A** to NHC complex **I**. By either pathway, the computed reaction is exergonic by 4.3 kcal/mol at 298 K. This value corresponds to an equilibrium constant favoring product by $>10^3:1$, in agreement with the observation that **1** reacts irreversibly with $(\text{PCy}_3)_2\text{RhCl}$ to give **3**. Also, as in the observed reaction of **1**, neither the “migration” nor the “rotation” mechanisms predict observable intermediates other than the N1- σ -bound complex (**A**).³⁹ Both calculated pathways are non-dissociative in phosphine and involve rate-limiting C-H activation steps, consistent with the significant primary kinetic isotope effect measured for the non-dissociative pathway of **2**-formation. The activation parameters calculated for both mechanistic variants compare well with experimental values (Table 2).⁴⁰

In the model system, the “rotation” and “migration” mechanisms seem nearly interchangeable (the highest overall barriers differ by an insignificant amount, <2 kcal/mol at 298 K). In the real system, however, when the PMe_3 ligands are replaced by considerably more bulky PCy_3 ligands, the same is not likely to be true. As a result of non-bonded interactions between the phosphine and heterocycle ligands, we expect the transition state for C2-Rh bond rotation (**F**) to be especially high in energy in the real system, possibly prohibitively so. The “migration” pathway avoids obvious steric interactions with PR_3 because the heterocycle and the Rh-square plane stay relatively perpendicular throughout the reaction coordinate. In light of this divergence, the extreme preference for bulky phosphines that we have consistently observed in Rh(I)/ PR_3 -catalyzed heterocycle coupling reactions can now potentially be understood in terms of the limitations that such phosphines place on accessible (deleterious) reaction pathways.

Discussion

By combining kinetic, structural and computational data it is possible to draw a plausible general mechanism for the C-H activation of *N*-heterocycles by Rh(I)/ PCy_3 (Scheme 6). The reaction is initiated by the association of a heterocycle to $(\text{PCy}_3)_2\text{RhCl}$ as a dative ligand through an sp^2 hybridized nitrogen. Then, the adjacent C-H bond is cleaved by rate-limiting intramolecular oxidative addition to give a transient rhodium hydride. A final, rapid H-transfer step restores rhodium to the +1 oxidation state and the transformation from heterocycle to NHC is complete. Less information is available about a competing PCy_3 -dissociative pathway. Since the experiment verifying the intermediacy of **3** was run under conditions in the which both

pathways make substantial contributions to $d[2]/dt$, the PCy₃-dissociative pathway can be said to proceed via intermediate **3**. Given such a ground state, the stoichiometry of the rate-limiting transition structure is likely to be as depicted in Scheme 6. However, since the dissociative reaction could not be studied in isolation, the validity of depicted bonding cannot be verified at this time.

This mechanism of Scheme 6 bears similarities to related directed C-H activation reactions. As in the Murai system, the directing group functions to bring the reactive metal fragment into proximity with the scissile C-H bond via pre-coordination.⁴¹ There are substantial differences, however, between that system, which relies on a pendant directing group, and a heterocycle C-H activation system, like the one discussed here. Pendant directing groups can stay coordinated to a metal fragment during C-H oxidative addition. Often described as taking place by cyclometalation, these reaction steps are promoted by the formation of stable 5- or 6-membered chelate rings. Calculated models of our system, in contrast, show that the directing heteroatom (N1) must detach from the metal fragment prior to C-H cleavage.⁴² Thus, catalytic functionalization of *N*-heterocycles by Rh(I)/PR₃ can be said to operate via a complex-induced proximity effect. Only by proceeding through an *N*-bound intermediate can the reactants gain access to an energetically accessible transition state for C-H activation.

Cavell and coworkers have recently reported a Ni⁰-catalyzed C-H coupling of olefins to the 2-position of imidazolium salts (eq 7).⁴³ Similar to the Rh(I) C-H activation, this reaction probably proceeds through a well-characterized metal-NHC intermediate.⁴⁴ However, the proposed mechanism is most similar to that of other group 10 metal-catalyzed cross coupling reactions that operate within the M⁰/M²⁺ redox couple. No experimental evidence was found for directed C-H activation via a catalyst-substrate precomplex. Despite the differences between the Rh- and Ni-based mechanisms, both seem to benefit from a well of stability along the energy surface of the catalytic cycle provided by an M-NHC intermediate. Taken together, these studies strongly contradict the notion that NHC's tend to act only as dative ligands toward transition metals.

(8)

Since the Rh(I)-catalyzed functionalization reactions of dihydroquinazolines, such as **1**, occur under similar conditions and with similar selectivity patterns to the reactions of other *N*-heterocycles, the mechanism for C-H activation of **1** is likely general for Rh-NHC-mediated C-H processes. Strong predictions can thus be made about the substrate scope for Rh-catalyzed coupling reactions. Earlier work by our group showed that all competent substrates have in common the ability to form and stabilize an NHC tautomer. If our newly outlined mechanism is a general one, then it reveals the further requirement that the site of dative coordination (the *sp*²-hybridized nitrogen) and the scissile C-H bond be *cis* to one another about a C=N double bond. Only then can facile intramolecular oxidative addition take place. All competent substrates recognized so far are cyclic, and therefore meet this requirement. Potential acyclic substrates, such as formamidines or formimides, on the other hand, are unlikely to exhibit the necessary geometry for activation, because they exist predominately in the *S*-trans form. Indeed, in our hands, acyclic substrates have never produced coupling products. Nor have substrates requiring a hydride shift of farther than two atoms produced functionalized products to date.

The proposed reaction mechanism of Scheme 6 does provide some guidance concerning the expansion of substrate scope. Heterocycles capable of forming pyrazolyl⁴⁵ or “abnormal”-type⁴⁶ carbenes should, in principle, be compatible with this mechanism. More enticingly, our investigations reveal that the other atom (besides the *sp*²-nitrogen) bonded to the carbene carbon does not directly interact with catalyst during C-H activation. While this atom may be important to the electronic structure of active substrates, our previous work has already shown that, in addition to N, S and O are tolerated at this position.^{15d} Given recent reports by Bertrand and coworkers of stable amino-alkyl NHC's,⁴⁷ catalytic functionalization might now also be expected to proceed with C adjacent to the carbene position.⁴⁸ If successful, such a coupling reaction between cyclic imines and alkenes would constitute a rare example of catalytic hydroiminoacylation.⁴⁹

Conclusion

In order to better understand the C-H activation step governing the Rh(I)-catalyzed coupling of *N*-heterocycles and olefins, stoichiometric reactions between (PCy₃)₂RhCl and heterocycle **1** were studied. The product of these reactions, a Rh-NHC complex analogous to those previously reported as catalytic intermediates (**2**), was isolated and characterized. Furthermore, a datively bound catalyst-substrate complex (**3**) was identified as an intermediate in the formation of **2**. The Rh-mediated 1,2-hydrogen shift responsible for C-H cleavage and NHC formation was studied in detail using experimental and computational methods. Based on these studies, a directed intramolecular hydrogen transfer pathway proceeding via Rh-H intermediates was implicated as the operative reaction mechanism. Similarities between the reactivity patterns of dihydroquinazolines and other *N*-heterocycles activated by (PCy₃)₂RhCl suggest that this mechanism may be a common route to reactive M-NHC intermediates. Insights gained from this mechanistic study are being applied toward the ongoing development of Rh (I)-catalyzed C-H functionalization reactions in our laboratories.

Experimental

General Experimental

Unless otherwise noted, all reagents were obtained from commercial suppliers and degassed prior to use. Tetrahydrofuran (THF) and hexanes were passed through a column of activated alumina (A1, 12 × 32, Purifry Co.) under nitrogen pressure and sparged with nitrogen prior to use.⁵⁰ The 1,4-dioxane was purchased anhydrous, stored over 3 Å activated molecular sieves, and degassed prior to use. Benzene-*d*₆ was degassed by three freeze-pump-thaw cycles and stored over 3 Å activated molecular sieves. THF-*d*₈ was vacuum transferred from sodium/benzophenone ketyl prior to use. All compounds were stored and handled in a N₂-filled IT Braun inert atmosphere glove box. All reactions were carried out in oven-dried glassware. Thin layer chromatography was performed on Merck 60 F254 250-μm silica gel plates. The developed chromatograms were visualized by UV (254 nm) light. Flash column chromatography was carried out using Merck 60 230–400 mesh silica gel. All infrared (IR) spectra were recorded on a Thermo Electron Avatar 370 spectrometer fitted with a single bounce ZnSe ATR plate. Only partial IR data are listed. The ¹H, ¹³C {¹H}, ³¹P {¹H}, ²H {¹H}, ¹⁵N NMR spectra were obtained on Bruker AV-300, AVQ-400, AVB-400, DRX-500 and AV-500 instruments and referenced to residual solvent. When necessary, *J* coupling was deconvoluted using ¹³C {¹H, ¹⁵N} NMR experiments and ¹³C assignments were made based on DEPT experiments. Elemental analyses were performed by the University of California, Berkeley elemental analysis facility. The X-ray crystal structure was obtained at the UC Berkeley CHEXRAY. [(coe)₂RhCl]₂ was prepared according to literature procedures and stored at –30 °C under N₂.⁵¹

3-Methyl-3,4-dihydroquinazoline (1)—Following a modified literature procedure⁵² 3,4-dihydroquinazoline⁵³ (907 mg, 6.86 mmol) and iodomethane (974 mg, 6.86 mmol) were sealed into a glass-walled vessel equipped with a vacuum stopcock with MeOH (7 mL) and allowed to stand at 25 °C for 4 d until crystallization of salts was complete. Then the reaction suspension was treated with 30 mL of 0.6 M NaOH and extracted with EtOAc (3x 30 mL). The combined organic phases were washed with brine, dried over MgSO₄, and concentrated onto a small amount of silica. Column chromatography on silica (5:4:0.5:0.5 EtOAc/Acetone/MeOH/Et₃N) afforded a white powder that was redissolved in THF, filtered to remove traces of silica, and concentrated under vacuum. The white crystals obtained from this procedure (380 mg, 38%) exhibited spectroscopic properties consistent with literature data.⁵⁴ ¹H NMR (500 MHz, C₆D₆): δ 7.49 (d, 1H, *J* = 8.0 Hz, Ar-*H*), 7.07 (t, 1H, *J* = 7.5 Hz), 6.90 (t, 1H, *J* = 7.5 Hz, Ar-*H*), 6.58 (s, 1H, C2-*H*), 6.56 (d, 1H, *J* = 7.5 Hz, Ar-*H*), 3.74 (s, 2H, CH₂), 1.80 (s, 3H, CH₃). MS (EI); *m/z* (%): 145 [M⁺ - H] (100), 147 [M⁺ + (D-M⁺ - H)] (47.3).

trans-(3-Methyl-1,2,3,4-tetrahydro-1*H*-quinazolin-2-ylidene)(PCy₃)₂RhCl (2, 2_{cross} from crossover experiment)—In a N₂ atmosphere glovebox **1** (37 mg, 0.25 mmol), [(coe)₂RhCl]₂ (90 mg, 0.13 mmol) and PCy₃ (140 mg, 50 mmol) were combined in THF (5 mL) and sealed into a glass-walled vessel equipped with a vacuum stopcock. The reaction mixture was heated to 75 °C for 24 h and then returned to the glovebox. After the reaction solvent was removed under vacuum **2** was obtained as a yellow solid. This material could be further purified by crystallization (slow diffusion of hexanes into THF at -30 °C under N₂) to give clear yellow blocks (153 mg, 72%). IR: 3422, 2913, 2845, 1603, 1514, 1472, 1445, 1403, 1058, 947, 732 cm⁻¹. ¹H NMR (300 MHz, *d*₈-THF): δ 7.99 (s, 1H, N-*H*), 7.16 (t, 1H, *J* = 7.6 Hz, Ar-*H*), 7.02 (d, 1H, *J* = 7.1 Hz, Ar-*H*), 6.92 (t, 1H, *J* = 7.4 Hz, Ar-*H*), 6.67 (d, 1H, *J* = 7.7 Hz, Ar-*H*), 4.29 (s, 2H, CH₂), 4.10 (s, 3H, CH₃), 2.3 - 0.9 (m, 66H, PCy₃). ¹³C NMR (75 MHz, *d*₈-THF): δ 134.4 (d, *J*_{Rh-C} = 1.6 Hz, C9), 129.5 (*C*_{Ar}), 127.0 (*C*_{Ar}), 123.5 (*C*_{Ar}), 118.6 (C10), 113.3 (*C*_{Ar}), 49.7 (CH₂), 48.4 (CH₃), 36.2 (t, *J*_{P-C} ≈ *J*_{Rh-C} = 8.2 Hz, PCH), 32.2 (bs, 1C, PCy), 31.0 (s, 1C, PCy), 29.1 (t, 2C, *J*_{P-C} ≈ *J*_{Rh-C} = 4.8 Hz, PCHCH₂), 28.0 (s, 1C, PCy), (missing C-2 signal revealed by ¹³C-labelling). ¹³C NMR (100 MHz, C₆D₆) (only partial data given): δ 32.0 (PCy), 30.8 (PCy), 28.9 (t, *J*_{P-C} ≈ *J*_{Rh-C} = 4.7 Hz, PCHCH₂), 28.8 (t, *J*_{P-C} ≈ *J*_{Rh-C} = 4.9 Hz, PCHCH₂), 27.7 (PCy). ³¹P NMR (200 MHz, *d*₈-THF): δ 28.0 (d, *J*_{Rh-P} = 152 Hz). Anal. Calc'd for C₄₅H₇₆ClN₂P₂Rh: C, 63.93; H, 9.06; N, 3.31. Found: C, 64.02; H, 9.14; N, 3.59. See Supporting Information for X-ray crystal structure.

trans-N1-(3-Methyl-3,4-dihydroquinazoline)(PCy₃)₂RhCl (3)—In a N₂ atmosphere glovebox **1** (73 mg, 0.50 mmol), [(coe)₂RhCl]₂ (180 mg, 0.25 mmol) and PCy₃ (280 mg, 1.00 mmol) were dissolved in 10 mL of THF. The solution was allowed to stand for 5 d. Then the solid was separated by filtration, washed with pentane, and the remaining solvent was removed under vacuum. The resulting orange powder (202 mg, 46%) was stored at -30 °C under N₂. Under these conditions this compound is relatively stable with color change to pale green (decomposition) occurring slowly (>1 month). ¹H NMR (400 MHz, *d*₈-THF): δ 10.04 (d, 1H, *J* = 8.4 Hz, C8-*H*), 7.61 (s, 1H, C2-*H*), 7.20 (t, 1H, *J* = 7.6 Hz, Ar-*H*), 6.97 (t, 1H, *J* = 7.4 Hz, Ar-*H*), 6.71 (d, 1H, *J* = 7.2 Hz, Ar-*H*), 4.51 (s, 2H, CH₂), 2.97 (s, 3H, CH₃), 2.2-0.9 (m, 66H, PCy₃). ³¹P NMR (160 MHz, *d*₈-THF): δ 22.1 (d, *J*_{Rh-P} = 143 Hz). Inconsistent amounts of bound solvent prevented this complex from giving reliable microanalysis data.

General procedure for the synthesis of isotopically labeled 3-methyl-3,4-dihydroquinazolines

To an NMR tube was added 2-methylaminomethyl-phenylamine (100 mg, 0.73 mmol) and 0.5 mL of formic acid. The tube was flame sealed under vacuum and heated for 1 h until ¹³C NMR analysis indicated that a complete reaction had occurred. After cooling, the tube contents were diluted with 2 mL of water, basified with 0.5 mL NH₄OH, and extracted with CH₂Cl₂ (3x

2 mL). The combined organic phases were washed with brine, dried over MgSO₄ and concentrated under vacuum to give a white solid. Purification by crystallization (slow diffusion of hexanes into THF at -30 °C under N₂) gave white needles.

2-Methylaminomethyl-phenyl-¹⁵N-amine (4)—The reaction was run according to a modified general procedure for the amidation of aryl bromides.⁵⁵ In an N₂ atmosphere glovebox CuI (94 mg, 0.49 mmol) and *N,N'*-dimethyl-ethane-1,2-diamine (87 mg, 0.98 mmol) were combined in 1,4-dioxane (5 mL), and this mixture was stirred until the CuI completely dissolved. The resulting solution was added to a glass-walled vessel equipped with a vacuum stopcock containing (2-bromo-benzyl)-methyl-amine⁵⁶ (983 mg, 4.91 mmol), ¹⁵N-benzamide (600 mg, 4.91 mmol), K₂CO₃ (1.36 g, 9.83 mmol) and a magnetic stirbar. The reaction vessel was sealed and heated at 110 °C for 5.5 h until TLC (5% Et₃N in EtOAc) indicated complete consumption of the aryl bromide. The crude reaction mixture was filtered through a pad of silica eluting with CH₂Cl₂ containing 5% of a mixture of 10% NH₄OH in MeOH to give a yellow oil. This material was suspended in 5 mL of water and then 10 mL of conc. HCl was added dropwise. The resulting solution was washed (3x 2 mL of Et₂O) and then resealed into the original reaction vessel and heated at 105 °C for 24 h. After cooling to 0 °C the reaction solution was basified by dropwise addition of NH₄OH and extracted with Et₂O (3x 40 mL). The combined ethereal phases were washed with brine, dried over MgSO₄, and concentrated under vacuum to give a yellow oil. After silica gel chromatography (50% → 95% EtOAc in hexanes with 5% Et₃N), **4** (324 mg, 48%) was obtained as a yellow oil. Its spectroscopic properties were consistent with data reported for 2-methylaminomethyl-phenylamine.⁵⁷ ¹H NMR (500 MHz, C₆D₆): 7.11 (t, 1H, *J* = 7.6 Hz, Ar-*H*), 6.96 (d, 1H, *J* = 7.3 Hz, Ar-*H*), 6.73 (t, 1H, *J* = 7.4 Hz, Ar-*H*), 6.43 (d, 1H, *J* = 7.8 Hz, Ar-*H*), 4.43 (bd, 2H, *J*_{N-H} = 79.0 Hz, ¹⁵NH₂), 3.47 (s, 2H, CH₂), 2.01 (s, 3H, CH₃), 0.29 (bs, 1H, NH). ¹³C NMR (125 MHz, C₆D₆): 148.3 (d, *J*_{N-C} = 10.4 Hz, C-¹⁵N), 130.5, 129.0, 124.4 (d, *J*_{N-C} = 2.0 Hz), 117.8, 116.0 (d, *J*_{N-C} = 1.8 Hz), 56.1 (CH₂), 36.1 (CH₃). ¹⁵N {¹H} NMR (40 MHz, C₆D₆): 55.8.

3-Methyl-3,4-dihydro-1-¹⁵N-quinazoline (¹⁵M-1)—The reaction was performed as described in the general procedure using **4** (100 mg, 0.73 mmol). The white solid obtained after crystallization (55 mg, 51%) exhibited spectroscopic properties consistent with data collected for **1**. ¹H NMR (400 MHz, CDCl₃): δ 7.14 (t, 1H, *J* = 7.6 Hz, Ar-*H*), 7.04 (d, 1H, *J* = 7.9 Hz, Ar-*H*), 6.98 (t, 1H, *J* = 7.4 Hz, Ar-*H*), 6.96 (d, 1H, *J*_{N-H} = 13.1 Hz, C2-*H*), 6.83 (d, 1H, *J* = 7.5 Hz, Ar-*H*), 4.49 (s, 2H, CH₂), 2.88 (s, 3H, CH₃). ¹³C NMR (100 MHz, CDCl₃): δ 150.3 (d, *J*_{N-C} = 3.0 Hz, C2), 141.5, 128.3 (d, *J*_{N-C} = 3.0 Hz), 125.5, 124.6, 124.5 (d, *J*_{N-C} = 8.9 Hz, C9), 120.0, 48.9, 39.7. ¹⁵N NMR (40 MHz, CDCl₃): δ 203.4 (d, *J*_{H-N} = 13.2 Hz). MS (EI); *m/z* (%): 145 [¹⁴N-M⁺ - H] (2.0), 146 [(¹⁴N-M⁺) + (¹⁵N-M⁺ - H)] (100), 147 [(¹⁵N-M⁺) + (¹⁵N, D-M⁺ - H)] (45.6). Calc'd: 99% ¹⁵N by comparison to the fragmentation pattern of **1**.

3-Methyl-3,4-dihydro-1-¹⁵N,2-¹³C-quinazoline (¹⁵M, ¹³C2-1)—The reaction was performed as described in the general procedure using **4** (75 mg, 0.55 mmol) with ¹³C-formic acid (250 mg, 5.3 mmol). The white solid obtained after crystallization (27 mg, 33%) exhibited spectroscopic properties consistent with data collected for ¹⁵N1-**1**. ¹H NMR (400 MHz, C₆D₆): δ 7.49 (d, 1H, *J* = 7.8 Hz, Ar-*H*), 7.08 (t, 1H, *J* = 7.7 Hz, Ar-*H*), 6.90 (t, 1H, *J* = 7.4 Hz, Ar-*H*), 6.59 (dd, 1H, *J*_{C-H} = 190 Hz, *J*_{N-H} = 13.4 Hz, C2-*H*), 6.57 (d, 1H, *J* = 7.6 Hz, Ar-*H*), 3.75 (s, 2H, CH₂), 1.81 (d, 3H, *J*_{C-H} = 4.0 Hz, CH₃). ¹³C NMR (100 MHz, THF) (only partial data given): δ 150.1 (d, *J*_{N-C} = 3.0 Hz, C2). MS (EI); *m/z* (%): 146 [singly labeled-M⁺ - H] (3.2), 147 [(singly labeled-M⁺) + (¹⁵N, ¹³C-M⁺ - H)] (100), 148 [(¹⁵N, ¹³C-M⁺) + (¹⁵N, ¹³C, D-M⁺ - H)] (44.8). Calc'd: 96% ¹⁵N, ¹³C by comparison to the fragmentation pattern of **1**.

3-Methyl-3,4-dihydro-2-²H-quinazoline (C2-D-1)—The reaction was performed as described in the general procedure using 2-methylaminomethyl-phenyl-¹⁴N-amine⁵⁸ (125 mg, 0.92 mmol) with D₂-formic acid (500 mg, 10.4 mmol). The white solid obtained after crystallization (48 mg, 35%) exhibited spectroscopic properties consistent with data collected for **1**, except that the ¹H NMR signal at δ 6.6 ppm was absent, as expected. IR: 2240 cm⁻¹ (C-D). ¹H NMR (400 MHz, C₆D₆): δ 7.51 (d, 1H, *J* = 7.8 Hz, Ar-*H*), 7.08 (t, 1H, *J* = 8.3 Hz, Ar-*H*), 6.89 (t, 1H, *J* = 7.4 Hz, Ar-*H*), 6.56 (d, 1H, *J* = 7.5 Hz, Ar-*H*), 3.74 (s, 2H, CH₂), 1.79 (s, 3H, CH₃). ¹³C NMR (125 MHz, DCO₂D): δ 147.4 (t, *J*_{D-C} = 31 Hz, C2), 127.4, 126.9, 125.8, 124.8, 115.0, 114.9, 46.6, 39.8. ²H NMR (61 MHz, C₆H₆): δ 6.6. MS (ESI); *m/z* (%): 147 [¹H-M⁺ + H] (0.92), 148 [M⁺ + H] (100), 149 [D-M⁺ + H] (10.5). Calc'd: 99% ²H.

General procedure for observing isotopically labeled Rh-complexes

To an NMR tube was added a solution containing labeled-**1** (3.8 mg, 25 μmol), [(coe)₂RhCl]₂ (9.0 mg, 13 μmol) and PCy₃ (14.0 mg, 50 μmol) in *d*₈-THF (0.5 mL). The tube was flame sealed under vacuum, then heated between 45 – 75 °C for 1 – 24 h in order to observe varying ratios of free **1**, **3**, and **2**. The distinctive resonances corresponding to the various complexes are reported as part of these mixtures.

From ¹⁵N**1-1**

trans-(3-Methyl-1,2,3,4-tetrahydro-1-¹⁵NH-quinazolin-2-ylidene)(PCy₃)₂RhCl (¹⁵N1-2, 2_{dir} from crossover experiment): Its spectroscopic properties were consistent with data collected for **2**. ¹H NMR (400 MHz, *d*₈-THF): δ 7.98 (d, 1H, *J*_{N-H} = 94.0 Hz, N-*H*). This compound was not sufficiently soluble to obtain an ¹⁵N NMR spectrum.

trans-N1-(3-Methyl-3,4-dihydro-1-¹⁵N-quinazoline)(PCy₃)₂RhCl (¹⁵N1-3): Its spectroscopic properties were consistent with data collected for **3**. ¹H NMR (400 MHz, *d*₈-THF): δ 7.61 (d, 1H, *J*_{N-H} = 7.2 Hz, C2-*H*). ³¹P NMR (160 MHz, *d*₈-THF): δ 22.1 (d, *J*_{Rh-P} = 143 Hz, *J*_{N-P} = 3.0 Hz). This compound was not sufficiently soluble to obtain an ¹⁵N NMR spectrum.

From ¹⁵N**1**, ¹³C**2-1**

trans-(3-Methyl-1,2,3,4-tetrahydro-1-¹⁵NH,2-¹³C-quinazolin-2-ylidene)(PCy₃)₂RhCl (¹⁵N1, ¹³C2-2): Its spectroscopic properties were consistent with data collected for ¹⁵N1-2. ¹H NMR (500 MHz, *d*₈-THF): δ 7.99 (dd, 1H, *J*_{N-H} = 94.5 Hz, *J*_{C-H} = 3.9 Hz, N-*H*), 4.09 (d, 3H, *J*_{C-H} = 4.0 Hz, CH₃). ¹³C NMR (125 MHz, *d*₈-THF): δ 216.4 (dtd, *J*_{Rh-C} = 54.4 Hz, *J*_{P-C} = 12.1 Hz, *J*_{N-C} = 2.0 Hz, C2).

trans-N1-(3-Methyl-3,4-dihydro-1-¹⁵N,2-¹³C-quinazoline)(PCy₃)₂RhCl (¹⁵N1, ¹³C2-3): Its spectroscopic properties were consistent with data collected for ¹⁵N1-3. ¹H NMR (500 MHz, *d*₈-THF): δ 7.60 (dd, 1H, *J*_{C-H} = 193 Hz, *J*_{N-H} = 7.4 Hz, C2-*H*), 2.97 (d, 3H, *J*_{C-H} = 4.1 Hz, CH₃). ¹³C NMR (125 MHz, *d*₈-THF): δ 155.2 (d, *J*_{N-C} = 10.5 Hz, C2).

From C2-D-1

trans-(3-Methyl-1,2,3,4-tetrahydro-1H,2-²H-quinazolin-2-ylidene)-(PCy₃)₂RhCl (C2-D-2, 2_{dirD} from crossover experiment): Its spectroscopic properties were consistent with data collected for **2**, except that the ¹H NMR signal at δ 8.0 ppm (N-*H*) was greatly diminished, as expected. IR: 2543 cm⁻¹ (N-D).

trans-N1-(3-Methyl-3,4-dihydro-2-²H-quinazoline)(PCy₃)₂RhCl (C2-D-3): Its spectroscopic properties were consistent with data collected for **3**, except that the ¹H NMR signal at δ 7.6 ppm (C2-*H*) was absent, as expected.

Procedure for deuterium tracer experiment

In a N₂ atmosphere glovebox C2-D-1 (3.8 mg, 26 μmol), [(coe)₂RhCl]₂ (8.9 mg, 12 μmol), PCy₃ (14.4 mg, 51 μmol) and 2,6-dimethoxytoluene (1.1 mg, 7 μmol) were combined in *d*₈-THF (0.5 mL). This mixture was flame sealed into a medium-walled NMR tube and then heated at 70 °C overnight. After cooling to room temperature, analysis of the reaction mixture by ¹H NMR indicated 80% yield of **3** with 88% incorporation of deuterium at the N1-H position.

Procedure for double labeling crossover experiment (with C2-D-1 and ¹⁵N1-1)

In a N₂ atmosphere glovebox C2-D-1 (0.9 mg, 6 μmol), ¹⁵N1-1 (0.9 mg, 6 μmol), [(coe)₂RhCl]₂ (4.5 mg, 6 μmol), PCy₃ (8.8 mg, 31 μmol) and 2,6-dimethoxytoluene (1.9 mg, 12 μmol) were combined in *d*₈-THF (0.5 mL). This mixture was flame sealed into a medium-walled NMR tube that was then heated at 45 °C for 1 d. Periodically, the reaction tube was removed from its heating bath, cooled to room temperature, and a one-pulse ¹H NMR spectrum was acquired prior to reheating to 45 °C. Based on these spectra, the relative abundance of the various isotopomers of **2** was determined (Table 1).

General procedures for measurements of reaction rate

The reactions were carried out in medium-walled NMR tubes flame sealed with a solid glass rod inside to minimize solvent reflux during heating. Reaction progress was monitored by *in situ* NMR analysis in the temperature-controlled probe of a Bruker DRX-500 MHz spectrometer. Instrument temperature readings were verified by reference to an ethylene glycol standard. Single pulse ¹H NMR spectra were collected at regular intervals of at least five times the longest measured T₁. The peak area of methyl and methylene resonances for the dihydroquinazoline moiety were measured relative to the internal standard (2,6-dimethoxytoluene) methoxy group peak area in order to obtain absolute concentrations. Initially, a reaction was run to complete conversion and the resulting conc. vs. time data were satisfactorily fit to our kinetic model (*vide infra*) (Figure S-1a). Subsequent reactions were monitored to between 10–40% conversion (Figure S-1b). Concentration vs. time data were fitted using the Gepasi 3.0 software package (mass action kinetics) in order to obtain rate constants corresponding to Scheme 3, Path A. The initial concentration of free (PCy₃)₂RhCl in solution was set based on the weight of added reagents because it could not be independently measured by NMR. While the actual concentration of (PCy₃)₂RhCl is determined by complex equilibria (Scheme 5), for the sake of data fitting, approximating [(PCy₃)₂RhCl] as the total concentration of free Rh(I) proved to be an acceptable simplification. A term for slow unimolecular decomposition of (PCy₃)₂RhCl (*k*_d < 6.5 × 10⁻⁵ s⁻¹) was included in all fits.

General methods for preparing reaction solutions for rate studies

Method A (*in situ* generation)—In an N₂ atmosphere glovebox **1** (1.8 mg, 13 μmol), [(coe)₂RhCl]₂ (4.5 mg, 6 μmol), PCy₃ (8.8 mg, 31 μmol) and 2,6-dimethoxytoluene (1.9 mg, 12 μmol) were combined in *d*₈-THF (0.5 mL). The resulting dark red solution was transferred to an NMR tube, frozen using liquid N₂, flame sealed under vacuum and kept frozen until use.

Method B (using isolated intermediate)—In an N₂ glovebox **3** (11 mg, 13 μmol), PCy₃ (1.8 mg, 6 μmol) and 2,6-dimethoxytoluene (1.9 mg, 12 μmol) were transferred to an NMR tube using *d*₈-THF (0.5 mL). After sealing under vacuum, the reaction tube was heated with agitation using a warm water bath (45 °C) until the light orange suspension became dark red and homogeneous. The reaction tube was maintained at 45 °C until use.

Method for measuring deuterium kinetic isotope effect—In an N₂ atmosphere glovebox [(coe)₂RhCl]₂ (9.0 mg, 12 μmol), PCy₃ (17.5 mg, 62 μmol) and 2,6-dimethoxytoluene (1.9 mg, 12 μmol) were combined in *d*₈-THF (1 mL). This solution was

divided into two portions, one of which was combined with **1** (1.8 mg, 13 μ mol) while the other was combined with **C2-D-1** (1.8 mg, 13 μ mol). The resulting dark red solutions were transferred to separate NMR tubes, frozen using liquid N₂, flame sealed under vacuum, and kept frozen until use. Reaction rates were otherwise measured as usual.

Computation Methodology

The B3LYP/LACVP**+//B3LYP/LACVP** level of theory⁵⁹ with “medium” grid density as implemented in the Jaguar 4.0 quantum chemistry program package has been utilized throughout this study.⁶⁰ For N, C, Cl, P and H, the 6–31G basis set of Pople and co-workers was used.⁶¹ For rhodium, a Hay-Wadt small core effective core potential replaces the 28 innermost core electrons.⁶² A single asterisk(*) indicates the addition of polarization functions to atoms Li through Ar; a double asterisk indicates the addition of polarization functions to atoms H through Ar. Similarly, a single plus (+) indicates the addition of diffuse functions to atoms Li through Ar; and double pluses indicate the addition of diffuse functions to atoms H through Ar. The basis set on rhodium has double ζ quality [contraction scheme {331/311/31}]. A full geometry optimization and analytical vibrational frequency calculation were performed for all simplified models. Stationary points are characterized by exactly zero imaginary vibrations; transition structures are characterized by exactly one imaginary vibration. Gibbs free energies ΔG for 298.15 K and 1 atm are based on unscaled molecular vibrations.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgement

The authors thank Dr. Fred Hollander at the UCB X-ray diffraction facility for determination of the crystal structure of **2**. Dr. Jeff Pelton of the UCB structural biology NMR facility is thanked for providing assistance for multi-nuclear decoupling experiments. We acknowledge Dr. Katherine Durkin of the Berkeley Molecular Graphics and Computation Facility for her helpful advice on the use of Jaguar. This work was supported by NIH grant GM069559 (to JAE) and by the Director and Office of Energy Research, Office of Basic Energy Sciences, Chemical Sciences Division, U. S. Department of Energy, under contract DE-AC02-05CH11231 (to RGB).

References

1. Wiedemann, SH.; Bergman, RG.; Ellman, JA. Handbook of C-H Transformations. Dyker, G., editor. Vol. 1. Weinheim: Wiley-VCH; 2005. p. 187
2. (a) Guari Y, Sabo-Etienne S, Chaudret B. Eur. J. Inorg. Chem 1999;1047. (b) Ritleng V, Sirlin C, Pfeffer M. Chem. Rev 2002;102:1731. [PubMed: 11996548] Kakiuchi, F.; Murai, S. Activation of Unreactive Bonds in Organic Synthesis. Murai, S., editor. Vol. 3. London, England: Freund Pub. House; 1999. p. 47
3. Murai S, Kakiuchi F, Sekine S, Tanaka Y, Kamatani A, Sonoda M, Chatani N. Nature 1993;366:529.
4. For discussion of a cyclometalation mechanism involving direct oxidative addition, see: Kakiuchi F, Sekine S, Tanaka Y, Kamatani A, Sonoda M, Chatani N, Murai S. Bull. Chem. Soc. Jpn 1995;68:62.
5. For discussion of a cyclometalation mechanism involving conjugate addition of Ru⁰, followed by alpha-elimination, see: (a) Matsubara T, Koga N, Musaev DG, Morokuma K. J. Am. Chem. Soc 1998;120:12692. (b) Matsubara T, Koga N, Musaev DG, Morokuma K. Organometallics 2000;19:2318.
6. Kakiuchi F, Murai S. Acc. Chem. Res 2002;35:826. [PubMed: 12379135]
7. Jordan RF, Taylor DF. J. Am. Chem. Soc 1989;111:778.
8. Jordan RF, Taylor DF, Baenziger NC. Organometallics 1990;9:1546.
9. Moore EJ, Pretzer WR, O'Connell TJ, Harris J, Labounty L, Chou L, Grimmer SS. J. Am. Chem. Soc 1992;114:5888.

10. (a) Chatani N, Fukuyama T, Kakiuchi F, Murai S. J. Am. Chem. Soc 1996;118:493. (b) Fukuyama T, Chatani N, Tatsumi J, Kakiuchi F, Murai S. J. Am. Chem. Soc 1998;120:11522. (c) Chatani N, Fukuyama T, Tatamidani H, Kakiuchi F, Murai S. J. Org. Chem 2000;65:4039. [PubMed: 10866622] (d) Szewczyk JW, Zuckerman RL, Bergman RG, Ellman JA. Angew. Chem., Int. Ed 2001;40:216.
11. (a) Youn SW, Pastine SJ, Sames D. Org. Lett 2004;6:581. [PubMed: 14961628] (b) Pittard KA, Lee JP, Cundari TR, Gunnoe TB, Petersen JL. Organometallics 2004;23:5514.
12. (a) Liu C, Han X, Wang X, Widenhoefer RA. J. Am. Chem. Soc 2004;126:3700. [PubMed: 15038708] (b) Liu C, Widenhoefer RA. J. Am. Chem. Soc 2004;126:10250. [PubMed: 15315430] (c) Liu C, Han XQ, Wang X, Widenhoefer RA. J. Am. Chem. Soc 2004;126:10493.
13. (a) Paras NA, MacMillan DWC. J. Am. Chem. Soc 2001;123:4370. [PubMed: 11457218] (b) Austin JF, MacMillan DWC. J. Am. Chem. Soc 2002;124:1172. [PubMed: 11841277]
14. Some oxidative protocols for heterocycle/olefin couple have appeared recently: (a) DeBoef B, Pastine SJ, Sames D. J. Am. Chem. Soc 2004;126:6556. [PubMed: 15161275] (b) Yi CS, Yun SY, Guzei IA. Organometallics 2004;23:5392. (c) Beccalli EM, Broggini G. Tetrahedron Lett 2003;44:1919. (d) Ferreira EM, Stoltz BM. J. Am. Chem. Soc 2003;125:9578. [PubMed: 12904010]
15. (a) Tan KL, Bergman RG, Ellman JA. J. Am. Chem. Soc 2001;123:2685. [PubMed: 11456947] (b) Tan KL, Bergman RG, Ellman JA. J. Am. Chem. Soc 2002;124:13964. [PubMed: 12440873] (c) Tan KL, Bergman RG, Ellman JA. J. Am. Chem. Soc 2002;124:3202. [PubMed: 11916392] (d) Tan KL, Vasudevan A, Bergman RG, Ellman JA, Souers AJ. Org. Lett 2003;5:2131. [PubMed: 12790546] (e) Tan KL, Park S, Ellman JA, Bergman RG. J. Org. Chem 2004;69:7329. [PubMed: 15471487]
16. Wiedemann SH, Bergman RG, Ellman JA. Org. Lett 2004;6:1685. [PubMed: 15128267]
17. Wiedemann SH, Ellman JA, Bergman RG. J. Org. Chem. 2005accepted (jo052345b)
18. For a previous synthesis of a **1**-derived M-NHC complex, see: Basato M, Benetollo F, Facchin G, Michelin RA, Mozzon M, Pugliese S, Sgarbossa P, Sbovata SM, Tassan A. J. Organomet. Chem 2004;689:454.
19. Lewis JC, Wiedemann SH, Bergman RG, Ellman JA. Org. Lett 2004;6:35. [PubMed: 14703344]
20. Carlton L, De Sousa G. Polyhedron 1993;12:1377.
21. (a) Mendez NQ, Seyler JW, Arif AM, Gladysz JA. J. Am. Chem. Soc 1993;115:2323. (b) Stark GA, Gladysz JA. Inorg. Chem 1996;35:5509. [PubMed: 11666738]
22. While we have chosen to depict the metal center of **3** as square planar we cannot rule out the presence of one weakly bound inner-sphere solvent ligand.
23. Yao WB, Eisenstein O, Crabtree RH. Inorg. Chim. Acta 1997;254:105.
24. Pre-agostic interactions have been identified as intermediates preceding C-H oxidative addition. The question of whether the C8-H bond of **3** can undergo metal-insertion is beyond the scope of this report. For a discussion of the this rare type of interaction in a related system, see: Lewis JC, Wu J, Bergman RG, Ellman JA. Organometallics 2005;24:5737.
25. (a) Mendes P. Comput. Appl. Biosci 1993;9:563. [PubMed: 8293329] (b) Mendes P. Trends Biochem. Sci 1997;22:361. [PubMed: 9301339] (c) Mendes P, Kell DB. Bioinformatics 1998;14:869. [PubMed: 9927716]
26. Cyclometalation into the alkyl groups of (PCy₃)₂RhCl has been implicated in the decomposition of this complex: Hietkamp S, Stufkens DJ, Vrieze K. J. Organomet. Chem 1978;152:347.
27. The ¹H-NMR signals of PCy₃ are too poorly resolved to permit the measurement of any minor deuterium incorporation there. Also, *NI-D-2* is not soluble enough for the same measurement to be made by ²H-NMR.
28. Another scenario that is consistent with the observed data is a large extent of crossover in conjunction with a large kinetic isotope effect. This interpretation is invalidated by the small measured kinetic isotope effect (vide infra).
29. Other evidence that free HCl is uninvolved in C-H activation is provided by our observation that the rate of formation of **2** is insensitive to added acid (HCl) or base (K₂CO₃).
30. (a) Van Gaal HLM, Van Den Bekerom FLA. J. Organomet. Chem 1977;134:237. (b) Van Gaal HLM, Moers FG, Steggerda JJ. J. Organomet. Chem 1974;65:C43.
31. For a description of the decomposition of (PCy₃)₂RhCl, see ref ^{30a}.

32. Incomplete conversion of **1** was also observed for reactions employing related trialkylphosphines (P*i*Pr₃, PtBu₂Me). Higher conversions can be obtained by using excess (PR₃)₂RhCl.
33. The omission of more complicated [Rh] equilibria was not found to affect data fitting.
34. Added PCy₃ does not affect the position of the pre-equilibrium because (PCy₃)₂RhCl has been shown not to coordinate additional PCy₃ to give (PCy₃)₃RhCl (ref 30b).
35. For treatments of uncatalyzed and proton catalyzed 1,2-hydrogen shifts at imidazol-2-yl carbenes, see: (a) Amyes TL, Diver ST, Richard JP, Rivas FM, Toth K. J. Am. Chem. Soc 2004;126:4366. [PubMed: 15053626] (b) Bourissou D, Guerret O, Gabbai FP, Bertrand G. Chem. Rev 2000;100:39. [PubMed: 11749234]
36. For detailed computational studies of the oxidative addition of 1,3-dimethylimidazolium to (PR₃)_xRhCl^{36a} and IrH₃(PMe₃)₂^{36b} see: (a) Hawkes KJ, McGuinness DS, Cavell KJ, Yates BF. Dalton Trans 2004;2505. [PubMed: 15303166] (b) Appelhans LN, Zuccaccia D, Kovacevic A, Chianese AR, Miecznikowski JR, Macchioni A, Clot E, Eisenstein O, Crabtree RH. J. Am. Chem. Soc 2005;127:16299. [PubMed: 16287324]
37. For the sake of simplicity no inner-sphere solvent molecules were included in our models. In a related system, reaction coordinates calculated with inner sphere solvent did not differ significantly from coordinates lacking solvation (ref 36a).
38. Crabtree RH, Holt EM, Lavin M, Morehouse SM. Inorg. Chem 1985;24:1986.
39. While the “migration” pathway does pass through a relatively low energy rhodium-hydride intermediate (**E**), based on its relative free energy with respect to **A** ($\Delta G^\circ = 5.2$ kcal/mol) it is not predicted to be observable. This is consistent with the lack of observed ¹H NMR resonances in the hydride region.
40. The large, negative experimental and calculated entropies of activation are similar in magnitude to that calculated for a related oxidative addition proceeding from the C-H σ -complex between 1,3-dimethylimidazolium and (PH₃)₂RhCl (−10.4 cal/mol·K, ref ^{36a}).
41. An instance of *ortho*-directed transition metal mediated C-H activation that is not chelation-assisted has been recently reported: Zhang XW, Kanzelberger M, Emge TJ, Goldman AS. J. Am. Chem. Soc 2004;126:13192. [PubMed: 15479051]
42. Maintaining *N*-coordination during C-H oxidative addition of an *N*-heterocycle would lead to a strained three-membered chelate ring. This type of coordination is known for exclusively for early metals. See ref ⁸ and references therein.
43. Clement ND, Cavell KJ. Angew. Chem., Int. Ed 2004;43:3845.
44. Clement ND, Cavell KJ, Jones C, Elsevier CJ. Angew. Chem., Int. Ed 2004;43:1277.
45. Kocher C, Herrmann WA. J. Organomet. Chem 1997;532:261.
46. (a) Grundemann S, Kovacevic A, Albrecht M, Faller JW, Crabtree RH. J. Am. Chem. Soc 2002;124:10473. [PubMed: 12197749] (b) Lebel H, Janes MK, Charette AB, Nolan SP. J. Am. Chem. Soc 2004;126:5046. [PubMed: 15099069](c) ref ^{36b}.
47. (a) Lavallo V, Mafhouz J, Canac Y, Donnadieu B, Schoeller WW, Bertrand G. J. Am. Chem. Soc 2004;126:8670. [PubMed: 15250717] (b) Lavallo V, Canac Y, Prasang C, Donnadieu B, Bertrand G. Angew. Chem., Int. Ed 2005;44:5705.
48. We have once observed functionalization at such a site in the case of the purine (ref 15b).
49. Efficient, versatile Rh-catalyzed hydroiminoacylation is possible when the imine *N*-substituent is a metal-coordinating group (2-pyridyl). See: Jun CH, Lee JH. Pure. Appl. Chem 2004;76:577.
50. Alaimo PJ, Peters DW, Arnold J, Bergman RG. J. Chem. Ed 2001;78:64.
51. Vanderent A, Onderdelinden AL. Inorg. Syn 1990;28:90.
52. Armarego WL. J. Chem. Soc 1961:2697.
53. The 3,4-dihydroquinazoline was prepared according to a literature procedure: Lewis JC, Wiedemann SH, Bergman RG, Ellman JA. Org. Lett 2004;6:35. [PubMed: 14703344]
54. Tietz H, Rademacher O, Zahn G. Eur. J. Org. Chem 2000:2105.
55. Klapars A, Huang XH, Buchwald SL. J. Am. Chem. Soc 2002;124:7421. [PubMed: 12071751]
56. Analytical data and a synthetic protocol have been reported for (2-bromo-benzyl)-methyl-amine: Philippe N, Denivet F, Vasse J-L, Santos JSO, Levacher V, Dupas G. Tetrahedron 2003;59:8049.
57. Coyne WE, Cusic JW. J. Med. Chem 1968;11:1208. [PubMed: 5680038]

58. Prepared according to the procedure for **4**.
59. (a) Becke AD. J. Chem. Phys 1993;98:5648–5652. (b) Vosko SH, Wilk L, Nusair M. Can. J. Phys 1980;58:1200. (c) Lee CT, Yang WT, Parr RG. Phys. Rev. B 1988;37:785. (d) Krishnan R, Binkley JS, Seeger R, Pople JA. J. Chem. Phys 1980;72:650.
60. Jaguar 4.0, release 23. Portland, OR, USA: Schrodinger, Inc.; 1998.
61. Frisch MJ, Pople JA, Binkley JS. J. Chem. Phys 1984;80:3265.
62. (a) Hay PJ, Wadt WR. J. Chem. Phys 1985;82:270–283. (b) Hay PJ, Wadt WR. J. Chem. Phys 1985;82:299.

Figure 1.
X-ray crystal structure (ORTEP diagram) of **2**.

Figure 2.
Calculated structure for **5** showing Rh-H “pre-agostic” interaction.

Figure 3.

Plot of concentration vs. time for the species present during the reaction of **1** with Rh(I)/PCy₃ at 36.7 °C. The solid curves depict kinetics simulations of the data (Path A: $k_1 = 1.0 \pm 0.2 \times 10^{-2} \text{ M}^{-1}\text{s}^{-1}$, $k_{-1} = 2.0 \pm 0.2 \times 10^{-4} \text{ s}^{-1}$, $k_d = 1.10 \pm 0.05 \times 10^{-5} \text{ s}^{-1}$).

Figure 4. Plot of $[2]$ vs. time from Figure 3, enlarged to show fits for two possible reaction pathways depicted in Scheme 3. (Path A: $k_2 = 1.809 \pm 0.003 \times 10^{-5} \text{ s}^{-1}$).

Figure 5.

Graph of k_2 vs. $[\text{free PCy}_3]$ (at 62.7 °C). The curve depicts a linear least-squares fit $k_2 = a [\text{PCy}_3] + b$ ($a = 4.7 \times 10^{-5}$, $b = 4.4 \times 10^{-4}$). The filled point is provided for reference only (see Supporting Information for details), and is not included in the linear fit.

Figure 6.

Eyring plot of **2** formation (k_2) in the presence of 13 mM excess PCy₃ between 42.2 °C and 72.9 °C. The curve depicts a linear least-squares fit to $\ln(k_2/T) = a[1/T] + b$ ($a = -1.31 \pm 0.01 \times 10^4$, $b = 18.6 \pm 0.4$).

Figure 7.
Calculated reaction coordinates for C-H activation (zero-point calculations, 298 K).

Scheme 1.

The proposed mechanism for rhodium-catalyzed benzimidazole/alkene coupling.

Scheme 2.
 Synthesis of isotopically labeled Rh-complexes.

Scheme 3.

Possible mechanisms for carbene formation via C-H activation.

Scheme 4.

Proposed mechanism for isotopic scrambling of C2-D-3.

Scheme 5.
Equilibria governing $[(\text{PCy}_3)_2\text{RhCl}]$.

Scheme 6.
Proposed mechanism of C-H activation of **1**.

Table 1

Double labeling crossover experiment data

				% Yield		% Crossover	
Time (h:min)	2 _{dir}	2 _{cross}	2 _{dirD}	2 _{crossD} ^a	+ Total	Observed ^b	
0:00	0	0	0	0	0	NA	
1:30	10	1	6	16	16	5	
4:30	24	1	14	39	39	4	
8:20	34	3	19	56	56	5	
21:10	41	5	26	72	72	7	
27:20	41	6	27	75	75	8	

^a 2_{dirD} and 2_{crossD} are not independently observable by ¹H NMR analysis

^b Observed %Crossover = %2_{cross}/Total. Because 2_{crossD} is not independently observable, the true extent of isotopic scrambling may be greater than the observed value.

Table 2

Comparison of measured and computed activation parameters.

Mechanism	$\Delta G^\ddagger$ (kcal/mol) ^a	$\Delta H^\ddagger$ (kcal/mol)	$\Delta S^\ddagger$ (cal/mol·K)
Calculated "Migration"	28.3	24.0	-14.5
Calculated "Rotation"	30.0	26.0	-13.2
Experimental ^b	29.1 ± 0.4	26.0 ± 0.3	-10.3 ± 0.8

^a at 298.15 K^b derived from Eyring Plot (Figure 6)