Published in final edited form as:

J Am Chem Soc. 2007 April 18; 129(15): 4508–4509. doi:10.1021/ja068189n.

Nucleophilic Acylation of *o*-Quinone Methides: An Umpolung Strategy for the Synthesis of α -Aryl Ketones and Benzofurans

Anita E. Mattson and Karl A. Scheidt*

Department of Chemistry, Northwestern University, 2145 Sheridan Road, Evanston, Illinois 60208

The inversion of normal reactivity patterns, or *Umpolung*, enables the development of unconventional bond-forming strategies for the synthesis of biologically active target molecules.¹ The generation of acyl anion equivalents by the polarity reversal of carbonyl functional groups is one valuable *Umpolung* tactic. Given our interest in unconventional bond-forming strategies,² we decided to concentrate on developing an acyl anion approach to synthesize α -aryl ketones. The preparation of α -aryl ketones is a challenging goal and the majority of recent progress has focused on the development of transition metal-catalyzed couplings of enolates and aryl halides.³ Herein, we report the direct synthesis of α -aryl ketones (**3**) by the fluoride promoted addition of protected thiazolium carbinols (**1**) to quinone methides that are generated *in situ* from silyl protected phenols (**2**, eq 1).

(1)

We recently disclosed that fluoride promotes the nucleophilic acylation of nitroalkenes when stable *O*-silyl thiazolium carbinols (**1**) are employed as carbonyl anion precursor.⁴ Interestingly, Rokita has reported that *O*-silylated phenols (such as **2**) produce *o*-quinone methides when exposed to fluoride.^{5,6} We envisioned that combining compounds with the general structures of **1** and **2** with an appropriate fluoride source would simultaneously generate the corresponding carbonyl anion and *o*-quinone methide species *in the same reaction flask*. The subsequent combination of the nucleophilic carbonyl anion and electrophilic *o*-quinone methide should provide the desired α -aryl ketones (**3**) in a single operation.⁷ The main challenge with this approach is the simultaneous generation of two highly reactive intermediates.

Our investigations of this new coupling strategy focused on the addition of protected thiazolium carbinol **1a** to *o*-quinone methide precursor **2** in the presence of tetramethylammonium fluoride (Table 1, eq 2).⁸ The exposure of **1a** and **2a** to this fluoride source at $-50\text{ }^{\circ}\text{C}$ afforded an encouraging 43% yield of the desired α -aryl ketone **3** (entry 1). Optimization of the reaction conditions revealed that the best yields of **3** are obtained with lower temperatures ($-78\text{ }^{\circ}\text{C}$, 23 h) followed by slowly warming to $23\text{ }^{\circ}\text{C}$ (entry 2).^{9,10} Another important factor affecting the yield of the reaction is concentration: the optimal value of 0.1 M affords 72% of **3** (entries 2–

4). Although bromide leaving groups on the *o*-quinone methide precursor gives the highest yields, the corresponding benzylic chloride (**2b**) is also a suitable reaction partner (entry 2 vs. entry 5), albeit in reduced yield (53%).¹¹ To lend support to the intervention of an *o*-quinone methide, a competition experiment was conducted with 1 equiv each of **1a**, **2a**, and 2-methoxybenzylbromide in the presence of 2 equiv of Me₄N·F. Using the reaction conditions from Table 1, *only* aryl ketone **3** is observed with none of the methyl ether product that would arise from direct alkylation of the carbonyl anion.⁹

With the optimal parameters established for this *Umpolung* reaction, we turned our attention to investigating the scope of the process (Table 2, eq 3). A brief survey of protected thiazolium carbinols reveals the reaction accommodates additional aromatic substituents (entries 1–3). When carbinols derived from alkyl and α,β -unsaturated aldehydes are employed as the acyl anion precursor, the desired α -aryl ketones are formed in moderate yields (not shown 40–53%). While these observations are encouraging, further investigations of this new strategy are necessary to accommodate these particular substrates.

An examination of the *o*-quinone methide component of the reaction indicates that variously substituted protected phenols are competent electrophile progenitors. In addition to the unsubstituted methides (e.g. derived from **2**), alkyl and aryl substitution at R¹ provides good yields of the corresponding acylated products (60% and 68%, entry 5). Substitution on the phenyl ring is also well tolerated. The quinone methide precursor derived from *o*-vanillin was able to produce the desired α -aryl ketone in 68% yield (entry 6). A high yield of product was also observed when the quinone methide precursor **2f** was subjected to the reaction conditions (75%, entry 7). For certain structures, the benzylic bromide reactants are unstable and could not be easily isolated. For these cases, the corresponding benzylic chlorides can be successfully employed in the reaction (entries 5 and 7).

This fluoride-induced *Umpolung* strategy can be extended beyond *o*-quinone methides. For example, α -indoyl ketone compounds such as **12** can be accessed in good yield (70%) from the combination of a silyl protected gramine derivative **11** and thiazolium carbinol **1a** in the presence of 2.5 equivalents of Me₄N·F (Scheme 1).¹²

This new bond-forming process facilitates a total synthesis of the naturally occurring aromatase inhibitor, demethylmoracin I.¹³ In the key step, highly substituted α -aryl ketone **14** is prepared in 62% yield from thiazolium carbinol **13**⁹ and **2f** with Me₄N·F. The acid-catalyzed cyclization to the corresponding benzofuran **15** is accomplished in 79% yield. The prenyl group is installed using a Stille cross coupling reaction and global demethylation with lithium diphenylphosphanide¹⁴ in refluxing THF affords demethylmoracin I.

In summary, the synthesis of α -aryl ketones via the direct nucleophilic acylation of *o*-quinone methides has been reported. In this *Umpolung* process, a reactive carbonyl anion and *o*-quinone methide are generated in the same flask by addition of fluoride ion and then undergo productive bond formation to yield the desired adducts. Investigations into the utility and applications of thiazolium carbinols as unique acyl anion precursors are ongoing in our laboratory.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

This work has been generously supported by Northwestern, NIH (RO1 GM73072), Abbott Laboratories, Amgen, Boehringer-Ingelheim, 3M, and Wacker Chemical Corp. A.E.M. acknowledges an ACS Division of Organic

Chemistry Fellowship sponsored by Eli Lilly. We thank a reviewer for suggesting the competition experiment employing **1a**, **2a**, and 2-methoxy-benzylbromide with Me₄N⁺F⁻.

References

1. (a) Grobel BT, Seebach D. *Synthesis* 1977:357–402. (b) Seebach D. *Angew Chem Int Ed* 1979;18:239–258.
2. (a) Mattson AE, Bharadwaj AR, Scheidt KA. *J Am Chem Soc* 2004;126:2314–2315. [PubMed: 14982429] (b) Mattson AE, Scheidt KA. *Org Lett* 2004;6:4363–4366. [PubMed: 15524484] (c) Chan A, Scheidt KA. *Org Lett* 2005;7:905–908. [PubMed: 15727471] (d) Myers MC, Bharadwaj AR, Milgram BC, Scheidt KA. *J Am Chem Soc* 2005;127:14675–14680. [PubMed: 16231921]
3. (a) Fox JM, Huang XH, Chieffi A, Buchwald SL. *J Am Chem Soc* 2000;122:1360–1370. (b) Hamada T, Chieffi A, Ahman J, Buchwald SL. *J Am Chem Soc* 2002;124:1261–1268. [PubMed: 11841295] (c) Culkin DA, Hartwig JF. *Acc Chem Res* 2003;36:234–245. [PubMed: 12693921]
4. Mattson AE, Zuhl AM, Reynolds TE, Scheidt KA. *J Am Chem Soc* 2006;128:4932–4933. [PubMed: 16608309]
5. For a review on o-quinone methides, see: Van de Water RW, Pettus TRR. *Tetrahedron* 2002;58:5367–5405.
6. (a) Pande P, Shearer J, Yang JH, Greenberg WA, Rokita SE. *J Am Chem Soc* 1999;121:6773–6779. (b) Rokita SE, Yang JH, Pande P, Greenberg WA. *J Org Chem* 1997;62:3010–3012. [PubMed: 11671669] (c) Génisson Y, Tyler PC, Ball RG, Young RN. *J Am Chem Soc* 2001;123:11381–11387. [PubMed: 11707114] (d) Barrero AF, del Moral JFQ, Herrador MM, Arteaga P, Cortés M, Benites J, Rosellón A. *Tetrahedron* 2006;62:6012–6017.
7. For a singular example of a lithiated dithiane undergoing an addition to an ortho-oxygenated benzylic halide, see: *Tetrahedron Lett* 1988;29:6885–6889.
8. (a) Christe KO, Wilson WW, Wilson RD, Bau R, Feng JA. *J Am Chem Soc* 1990;112:7619–7625. (b) Menand ML, Dalla V. *Synlett* 2005:95–98.
9. The yield of **3** is lower (45%) when incubating the reaction at –78 °C for 1 h (vs. 23 h) followed by warming to 23 °C.
10. See Supporting Information for details.
11. The combination of **1a** and Me₄N⁺F⁻ with benzyl bromide affords only low yields of benzyl ketone under the best conditions from Table 1.
12. For examples of fluoride-induced elimination-addition reactions of N-triisopropylsilyl gramine methyl iodide, see: Iwao M, Motoi O. *Tetrahedron Lett* 1995;36:5929–5932.
13. Lee D, Bhat KPL, Fong HHS, Farnsworth NR, Pezzuto JM, Kinghorn AD. *J Nat Prod* 2001;64:1286–1293. [PubMed: 11678652]
14. McAllister GD, Hartley RC, Dawson MJ, Knaggs AR. *J Chem Soc, Perkin Trans 1* 1998:3453–3457.

Scheme 1.

Scheme 2.

Synthesis of Demethylmoracin I

a. $\text{Me}_4\text{N}^+\text{F}^-$, CH_2Cl_2 , -78 to 23 $^\circ\text{C}$, 62% b. Amberlyst 15, $4\text{\AA}$ sieves, CH_2Cl_2 79% c. prenyl tributyltin, $\text{PdCl}_2(\text{dppf})$, DMF, 100 $^\circ\text{C}$, 76% d. $n\text{BuLi}$, PPh_3 , THF, 61 %.

Table 1

Optimization of Carbonyl Anion Additions^a

Reaction scheme: **1a**, Ar = 4Cl-Ph + **2** $\xrightarrow[\text{CH}_2\text{Cl}_2]{\text{Me}_4\text{N}^+\text{F}^-}$ **3**, Ar = 4Cl-Ph (2)

entry	temp. (°C)	X	concentration	yield ^b
1	-50 to 23	Br (2a)	0.1 M	43%
2	-78 to 23	Br (2a)	0.1 M	72%
3	-78 to 23	Br (2a)	0.3 M	23%
4	-78 to 23	Br (2a)	0.05 M	45%
5	-78 to 23	Cl (2b)	0.1 M	53%

^aSee Supporting Information for details.^bIsolated yield

Table 2 α -Aryl Ketones Prepared from Thiazolium Carbinols (**1**) and Silylated Phenols (**2**)^a

Entry	Carbinol ^b	Halide	Aryl ketone
1			
2			
3			
4			
5			
6			
7			

^aSee Supporting Information for reaction details,^bTz = 2-substituted 3,4,5-trimethylthiazol-3-ium iodide.