Published in final edited form as:

Horm Behav. 2008 August ; 54(3): 455–462. doi:10.1016/j.yhbeh.2008.05.010.

Differential Effects of Acute Progesterone Administration on Spatial and Object Memory in Middle-Aged and Aged Female C57BL/6 Mice

Michael C. Lewis¹, Patrick T. Orr¹, and Karyn M. Frick^{1,2*}

¹Department of Psychology, Yale University, New Haven, CT 06520

²Interdisciplinary Neuroscience Program, Yale University, New Haven, CT 06520

Abstract

The present study examined the effects of acute progesterone administration on hippocampal-dependent memory consolidation in ovariectomized middle-aged (16 month) and aged (22 month old) female mice. Spatial memory was tested in a 2-day Morris water maze task and object memory was tested using an object recognition task with 24- and 48-hour delays. Immediately after water maze training, mice received i.p. injections of vehicle, or 5.0, 10.0, or 20.0 mg/kg of water-soluble progesterone. Twenty-four hours later, retention of the platform location was tested. No overnight forgetting of the platform location was observed in middle-aged vehicle-treated mice. Acute progesterone administration had no effect on spatial memory in middle-aged mice. However, aged vehicle-treated mice demonstrated impaired memory for the platform location on Day 2 relative to Day 1. Twenty mg/kg, but not 5 or 10 mg/kg, progesterone reversed these deficits, suggesting that 20 mg/kg progesterone can improve spatial memory in aged females. In the object recognition task, mice explored two identical objects and then immediately received vehicle or progesterone injections. In middle-aged mice, 10 and 20 mg/kg progesterone enhanced object memory consolidation, relative to chance, after 24-hours, but all doses were ineffective after 48-hours. In aged mice, 10 mg/kg progesterone enhanced object memory consolidation, relative to chance, after 24 hours, whereas both 5 and 10 mg/kg progesterone enhanced memory after 48 hours. Together, these results indicate that acute progesterone differentially enhances hippocampal-dependent memory in middle-aged and aged females.

Keywords

Progesterone; Object Recognition; Morris Water Maze; Steroid Hormones; Aging

1. Introduction

Interest in the effects of sex-steroid hormones, including estrogen and progesterone, on cognitive function has increased substantially in recent years due, in part, to clinical findings

*Corresponding Author: Karyn M Frick Ph.D., Department of Psychology, Yale University, 2 Hillhouse Ave. Rm. 106, New Haven, CT 06511, Phone: 203-432-4673, E-mail: Karyn.frick@yale.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Disclosure Statement: The author's declare that there are no actual or potential conflicts of interest.

suggesting that the reduction in hormone levels that occurs during menopause may contribute to age-related memory decline (Sherwin, 1999; Yaffe, 1998). Thus, a thorough understanding of the roles these hormones play in modulating memory formation could lead to improved treatment for age-related memory dysfunction. The effects of estrogen on regions of the brain involved in memory, including the hippocampus, have been well studied. One of the most potent forms of estrogen, 17 β -estradiol, has numerous effects on hippocampal physiology including increased CA1 dendritic spine density (Woolley and McEwen, 1992; Woolley and McEwen, 1993) and synaptic proteins (Choi et al., 2003; Stone et al., 1998), reduced GABAergic inhibition of CA1 pyramidal neurons (Murphy et al., 1998), and enhanced long-term potentiation (Foy et al., 1999; Warren et al., 1995) and neurogenesis (Tanapat et al., 1999). Estradiol also phosphorylates hippocampal signal transduction proteins such as extracellular signal-regulated kinase (ERK) (Carlstrom et al., 2001; Kuroki et al., 2000) and cyclic AMP response element binding protein (CREB) (Murphy and Segal, 2000). Most studies that have examined the effects of estradiol administration on hippocampal-dependent memory (e.g., spatial memory, object recognition) typically report improvements in young ovariectomized female rodents (Bimonte and Denenberg, 1999; Daniel et al., 1997; Fader et al., 1998; Farr et al., 1995; Gibbs, 1999; Gresack and Frick, 2004; Heikkinen et al., 2002; Luine and Rodriguez, 1994; O'Neal et al., 1996; Rissanen et al., 1999; Sandstrom and Williams, 2001; 2006b).

In contrast, much less is known about the effects of progesterone on hippocampal physiology and memory. Similar to estradiol, progesterone increases levels of the presynaptic protein synaptophysin in the female hippocampus (Choi et al., 2003), and increases ERK activation in hippocampal cultures (Nilsen and Brinton, 2002; Nilsen and Brinton, 2003). However, progesterone does not appear to be as beneficial for hippocampal function as estradiol. For example, progesterone blocks estradiol-induced increases in CREB phosphorylation in hippocampal cultures (Murphy and Segal, 2000). Further, estradiol alone, but not progesterone alone, modulates hippocampal NMDA receptor expression after ovariectomy (Cyr et al., 2001; El-Bakri et al., 2004). Progesterone administration 48 hours after two estradiol injections (spaced 24 hours apart) initially increases CA1 dendritic spine density for 2–6 hours after treatment, but then sharply decreases spine density much more than is observed with estradiol alone (Woolley and McEwen, 1993). Interestingly, this same estradiol and progesterone treatment improves spatial memory if rats are tested 90 minutes, but not 24 hours, after progesterone injection (Sandstrom and Williams, 2001). A similar estradiol and progesterone treatment impaired spatial reference memory in the Morris water maze four hours after progesterone injection, whereas estradiol alone or progesterone alone had no effect (Chesler and Juraska, 2000). Other studies have reported that chronic administration of progesterone or its metabolite, allopregnanolone, alone impairs spatial memory and footshock avoidance in young female rodents (Farr et al., 1995; Johansson et al., 2002). On the other hand, acute post-training systemic injections of estradiol alone, progesterone alone, or both hormones together improve inhibitory avoidance and novel object recognition (Frye and Lacey, 2000; Walf et al., 2006). Further, treatment with estradiol plus progesterone reduces spatial memory impairments induced by the cholinergic agonist scopolamine (Tanabe et al., 2004) and protects against memory impairments induced by intra-hippocampal administration of the neurotoxin colchicine (Vongher and Frye, 1999). As such, the effects of progesterone in young females may be highly dependent on factors such as the timing of testing relative to treatment and whether it is given alone or in combination with estradiol.

Of particular relevance to the study of hormones and cognition are the effects of sex-steroid hormone loss during aging in females. Although hippocampal-dependent memory deteriorates with age in both male and female rodents, age-related deficits in spatial memory tested in the Morris water maze occur earlier in females than males, and this deficit is associated with declining estrous cyclicity (Frick et al., 2000; Markowska, 1999). Because estrogen functions

as a trophic factor in the adult hippocampus (Brinton, 2001), hormone deficiency during aging may render the hippocampus more vulnerable to deterioration and exacerbate emerging age-related memory deficits. Numerous studies have shown that estradiol treatment can reduce age-related memory decline in middle-aged (Daniel et al., 2006; Fernandez & Frick, 2004; Foster et al., 2003; Markham, et al, 2002) and aged (Frick et al., 2002; Gibbs, 2000; Heikkinen et al., 2002; Markowska and Savonenko, 2002; Vaucher et al., 2002) female rodents. However, very few studies have examined the effects of progesterone on memory decline in aging females. The handful of studies that have tested the effects of estradiol plus progesterone on memory have yielded inconsistent results: although chronic treatment with estradiol plus progesterone improved spatial reference and working memory in aged females (Gibbs, 2000; Markham et al., 2002), progesterone reversed the beneficial effects of estradiol on spatial reference memory in middle-aged (Bimonte-Nelson et al., 2006) and aged (Harburger et al., 2007) females. No study to date has examined the effects of progesterone alone on memory in aging females, and therefore, it is unclear whether progesterone itself can ameliorate age-related cognitive deficits.

As such, the current experiment examined whether progesterone alone could improve hippocampal-dependent spatial and object memory in ovariectomized middle-aged and aged female mice using Morris water-maze and object recognition tasks. In order to pinpoint effects of progesterone on memory consolidation specifically, a single injection of water-soluble progesterone was given immediately after training (post-training) in each task. Because progesterone was not in the circulation during training or testing, the post-training design allowed effects on memory consolidation to be observed in the absence of non-mnemonic confounds (e.g., effects on motivation, anxiety) that may influence performance (Gresack and Frick, 2006; Harburger et al., 2007). The results suggest that acute post-training progesterone administration differentially affects both spatial and non-spatial forms of hippocampal-dependent memory in middle-aged and aged female mice.

2. Methods

2.1 Subjects

Subjects were 16 and 22 month-old female C57BL/6 mice (N = 8–11/group) obtained from the National Institutes on Aging colony at Harlan Sprague Dawley (Indianapolis, IN). Mice were housed up to five per shoebox cage in a room with a 12:12 light/dark cycle (lights on at 07:00), with all testing performed during the light phase. Mice had ad libitum access to food and water. Animals were handled for 5 min/day for at least five days prior to ovariectomy surgery to habituate them to being picked up by the experimenter. All procedures were approved by the Institutional Animal Care and Use Committee of Yale University, and conformed to the guidelines established by the National Institutes of Health Guide for the Care and Use of Laboratory Animals.

2.2 Ovariectomy

All mice were ovariectomized to eliminate endogenous ovarian production of estrogen and progesterone prior to treatment. Surgeries were conducted one week prior to behavioral testing as previously described (Fernandez and Frick, 2004). Briefly, mice were anesthetized using 2% isoflurane gas in 100% oxygen. Bilateral dorsal incisions were made at the level of the pelvis, and the ovaries and tips of the uterine horns were isolated and removed. Uteri were then placed back into the body cavity, the muscle wall was sutured, and the skin closed with wound clips, which were removed one week after surgery. Mice were given one week to recover from surgery, during which they were housed singly and received 30 mg/kg children's ibuprofen in the drinking water for analgesia. Mice were then re-housed with their original cage mates after wound clip removal.

2.3 Hormone Administration

Water-soluble progesterone (Progesterone: 2-hydroxypropyl- β -cyclodextrin complex) and HBC vehicle (2-hydroxypropyl- β -cyclodextrin complex) were dissolved in physiological saline. Doses of 5, 10, or 20 mg/kg progesterone were administered and hormone solutions were prepared to ensure these doses reflected the progesterone content of the HBC complex. Drugs were administered intraperitoneally (i.p.) immediately upon completion of training in each behavioral task. All animals were run in each task (water maze & object recognition with 24 and 48 delays). Thus, to minimize any effects of previous behavioral testing and/or drug administration, a minimum of two weeks elapsed between all behavioral procedures, including between the 24 and 48-hour versions of the object recognition task. Further, the order in which the mice were run in the various tasks was pseudo-randomly varied such that some groups ran in the water maze first, while others were initially run in the object recognition task. The HBC conjugate enhances the water solubility of progesterone without affecting the pharmacokinetics of the hormone (Pitha and Pitha, 1985). The advantage of HBC preparations is that they are readily metabolized within 24 hours and, thus, are not present during testing (Pitha, Harman, and Michel, 1986). This minimizes potential confounding effects of the drug on non-mnemonic aspects of task performance. Further, although plasma progesterone levels were not measured in this study, previous work has demonstrated that doses of progesterone between 0.1 and 1.0 mg produce physiological levels of progesterone (Akinci and Johnston, 1997). Thus, given the average weight of the middle-aged (26.5 g) and aged (27.8 g) mice, the doses of progesterone administered in the current study (5, 10, and 20 mg/kg) equate to (0.13, 0.26, & 0.52 mg) for middle-aged mice and (0.14, 0.28, & 0.56 mg) for aged mice, which are all in the physiological range.

2.4 Morris water maze

Morris water maze testing was conducted as previously described (Gresack and Frick, 2006; Harburger et al., 2007). Briefly, testing took place in a white circular tank (97 cm in diameter) filled with water (24 ± 2 °C). The water was made opaque with white nontoxic paint and the maze was surrounded by various extramaze cues. Data were collected using an HVS 2020 (Hampton, England) automated tracking system.

Mice were shaped one day prior to testing using a four-trial procedure in which a smaller ring (55 cm) was placed inside of the larger ring (97 cm) to decrease the total swimming area. Mice were first placed on a visible 10×10 cm platform (covered in red tape) for 10 s and then removed. They were then placed at three distances progressively further from the platform and allowed to swim to the platform. If the mouse did not find the platform within 60 s, then the experimenter led the mouse to it. No data were collected during shaping.

During testing, a transparent Lucite platform (10×10 cm) was submerged just underneath the surface of the water and remained in the same location for all trials. On Day 1 of testing, eight consecutive spatial trials were conducted. Each mouse was placed into the tank at one of the four cardinal (N, S, E, or W) start positions, which varied for each trial. The mouse was given 60 s to find the platform in each trial. If the platform was not located, then the experimenter led the mouse to the platform where it remained for 10 s. The mouse was then placed in a holding cage under a heat lamp for an intertrial interval of 45 s. Immediately after the completion of trial 8, the mouse was removed from the platform and injected with vehicle or progesterone (see Drug Administration). Twenty-four hours later, mice were re-tested in four consecutive spatial trials to examine spatial memory consolidation. As in previous studies (Gresack and Frick, 2006; Harburger et al., 2007; Packard & Teather, 1997), spatial memory consolidation can only be measured during the first trial of Day 2 because re-exposure to the platform in this trial serves to remind the mouse of the platform location. The remaining trials ascertain whether the mice can re-learn the platform location. Swim distance (m) was used to

measure memory and swim speed (m/s) was used to measure general motor performance throughout testing.

2.5 Object Recognition

This task assesses hippocampal-dependent, non-spatial object memory and consists of habituation, sample, and choice phases (Baker and Kim, 2002; Clark et al., 2000; Frick & Gresack, 2003; Mumby et al., 2005; O'Brien et al., 2006). Mice were first habituated to an empty white box by allowing them to freely explore for 5 min. Twenty-four hours later, mice were placed in the empty box for 1 min of additional habituation. Mice were removed while two identical objects were placed in the northeast and northwest corners of the box, approximately 5 cm from the walls. Mice were then placed back in the box facing the middle of the south wall and allowed to freely investigate the objects until they accumulated a total of 30 s exploring the objects, at which point the sample phase was terminated. Object exploration was scored only when the mouse's nose was within approximately 1 cm of the object or if the front paws were in contact with the object. Immediately after the completion of the sample phase, mice received vehicle or progesterone injections. The use of 30 s of total exploration time rather than a fixed trial duration minimizes the effects of group differences in activity (Frick & Gresack, 2003). Twenty-four and forty-eight hours later, mice were tested in the choice phase. This phase was identical to the sample phase, except that a novel object was substituted for one of the objects used in the sample phase and different objects were used for each delay. Time spent with the objects during the choice phase was recorded and compared to chance performance of 15 s (see Data Analysis). Total time to complete the choice phase was also recorded. The location of the novel object (northeast or northwest corner) was counterbalanced across mice and different objects were used for each of the delays. For both phases, objects and the box were cleaned with 70% EtOH between mice.

2.6 Data Analysis

For the Morris water maze, a series of analyses were performed for swim distance. Acquisition of the spatial task was analyzed using both a two-way mixed design ANOVA with one between-factor (drug treatment) and one within-factor (training trials), as well as a trend analysis designed to examine the rate of acquisition across the training trials on Day 1. A one-way ANOVA without repeated-measures was then performed on the last training trial (trial 8) to examine whether the groups exhibited similar performance at the end of training. We next performed a repeated-measures ANOVA comparing performance in trial 8 of Day 1 and trial 1 of Day 2 as a measure of memory consolidation. In addition, a one-way ANOVA was performed on difference scores calculated between these two trials (Trial 8 of Day 1 - Trial 1 of Day 2) to further examine changes in performance between days. We performed a two-way mixed ANOVA on all trials of Day 2 to examine whether the mice could re-learn the platform location after a reminder trial. Finally, a one-way repeated-measures ANOVA was performed on the last trial of day 2 to examine reacquisition among treatments groups. For swim speed, identical analyses were run as were run for swim distance to examine motor performance between groups in each phase of the task.

For object recognition, time spent with the novel object during the choice phase was analyzed using separate one-sample t-tests for each group to determine if the time spent with the novel object differed from the chance value of 15 s. This type of analysis was used because the time spent with each object in a total time of 30 s is not independent; time spent with one object necessarily reduces the time spent with the other object (Frick and Gresack, 2003; Baker and Kim, 2002). Because each group was analyzed individually relative to chance, between-group comparisons of exploration were made relative to chance and not directly between groups. Total time to complete the choice phase at each delay was analyzed with one-way ANOVAs including treatment as the independent variable.

3. Results

3.1 Morris Water Maze

3.1.1 Middle-Aged Mice—A repeated measures ANOVA examining swim distances throughout training on Day 1 (Figure 1A) revealed a significant main effect of Trial ($F(7,273) = 3.30, p < 0.05$), a significant main effect of Treatment ($F(3,39) = 3.06, p < 0.05$), but no Treatment $\times$ Trial interaction ($F(21,273) = 0.49, p > 0.05$). Further, a trend analysis examining the curve-fit to an inverse function across the 8 training trials of Day 1 revealed a significant effect of trial ($F(1,342) = 16.43, p < 0.001$), suggesting that all groups decreased their path to the platform as a function of training trial. Despite the overall main effect of treatment, a one-way ANOVA performed on Trial 8 of Day 1 revealed no differences in swim distance among the groups ($F(3,42) = 0.87, p > 0.05$), demonstrating no significant differences among the groups in performance in the last training trial before hormone administration.

The effects of post-training progesterone administration are shown on the right side of Figure 1A. A repeated-measures ANOVA examining swim distance during the last trial of Day 1 and the first trial of Day 2 revealed no significant main effect of Trial ($F(1,39) = 0.01, p > 0.05$), no significant main effect of Treatment ($F(3,39) = 0.84, p > 0.05$), and no Trial $\times$ Treatment interaction ($F(3,39) = 0.90, p > 0.05$), suggesting that none of the groups exhibited significant overnight forgetting. In addition, a one-way ANOVA examining difference scores revealed no differences among any of the groups ($F(3,39) = 0.90, p > 0.05$). As such, no beneficial effect of progesterone administration on spatial memory consolidation was observed. The similarities in performance between trial 8 of Day 1 and trial 1 of Day 2 is illustrated in Figure 2A, which plots only these trials. A repeated-measures ANOVA on the four trials of Day 2 revealed no significant main effect of Trial ($F(3,117) = 0.18, p > 0.05$), no significant main effect of Treatment ($F(3,39) = 0.45, p > 0.05$) and no Treatment $\times$ Trial interaction ($F(9,117) = 1.28, p > 0.05$), which is likely attributable to the fact that no overnight forgetting was apparent and, thus, improved performance was not easily observable. Lastly, a one-way ANOVA examining performance on the last trial of Day 2 revealed no significant differences between groups ($F(3,39) = 0.50, p > 0.05$), demonstrating that all groups performed similarly at the end of Day 2.

A repeated measures ANOVA examining swim speed during training on Day 1 (Figure 1C) revealed a significant main effect of Trial ($F(7,273) = 28.63, p < 0.001$), a significant main effect of Treatment ($F(3,39) = 3.85, p < 0.05$), but no Trial $\times$ Treatment interaction ($F(21,273) = 0.92, p > 0.05$). There were no differences in swim speed on trial 8 of Day 1 ($F(3,42) = 1.39, p > 0.05$), suggesting that, while some differences in swim speed existed between groups during training on trials 1–7, all mice had decreased in swim speed to a similar extent by trial 8. A repeated-measures ANOVA examining swim speed on Trial 8 of Day 1 to Trial 1 of Day 2 revealed a significant main effect of Trial ($F(1,39) = 27.00, p < 0.05$), no significant main effect of Treatment ($F(3,39) = 1.87, p > 0.05$), and no Trial $\times$ Treatment interaction ($F(3,39) = 0.32, p > 0.05$), suggesting that all mice swam faster on the first trial of Day 2. A repeated-measures ANOVA examining the four trials of Day 2 revealed no significant main effect of Trial ($F(3,117) = 1.71, p > 0.05$), no significant main effect of Treatment ($F(3,39) = 1.43, p > 0.05$) and no Treatment $\times$ Trial interaction ($F(9,117) = 0.71, p > 0.05$), indicating that swim speed did not decrease significantly across trials. Finally, a one-way ANOVA examining swim speed on the last trial of Day 2 revealed no significant differences among groups ($F(3,42) = 1.20, p > 0.05$) suggesting that all mice swam similarly at the end of Day 2.

3.1.2: Aged Mice—A repeated-measures ANOVA examining swim distances during training on Day 1 (Figure 1B) revealed a significant main effect of Trial ($F(7,245) = 7.15, p < 0.001$), no significant main effect of Treatment ($F(3,36) = 0.76, p > 0.05$) and no Treatment $\times$ Trial interaction ($F(21,245) = 0.51, p > 0.50$). Further, a trend analysis examining the curve-

fit to an inverse function across the 8 training trials of Day 1 revealed a significant effect of trial ($F(1, 318) = 38.18, p < 0.001$), suggesting that all groups decreased their path to the platform as a function of training trial. A one-way ANOVA performed on trial 8 revealed no differences in swim distance among the groups ($F(3,36) = 0.62, p > 0.05$), suggesting that all groups performed similarly in the last training trial prior to hormone administration.

The effects of post-training progesterone administration are illustrated on the right side of Figure 1B. A repeated-measures ANOVA examining swim distance during the last trial of Day 1 and the first trial of Day 2 revealed a significant main effect of Trial ($F(1, 36) = 14.96, p < 0.001$) a trend towards a significant main effect of Treatment ($F(3,36) = 2.37, p = 0.08$) and a trend for a Trial $\times$ Treatment interaction ($F(3, 36) = 2.63, p = 0.06$). In addition, a one-way ANOVA examining difference scores revealed a trend toward significance ($F(3,36) = 2.63, p = 0.06$). To further examine treatment effects from the last trial of Day 1 to the first trial of Day 2, paired-samples t-tests were conducted for each group (Figure 2B). The paired samples t-tests revealed a significant 148% increase in swim distance from Day 1 to Day 2 in the vehicle treated group ($t(9) = 3.47, p < 0.05$), indicating that the mice did not remember the platform location after 24 hours. Further, swim distances for both the 5 mg/kg ($t(9) = 2.01, p < 0.05$) and 10 mg/kg ($t(9) = 2.11, p < 0.05$) progesterone groups were also significantly higher on Day 2 (112% and 82% for the 5 and 10 mg/kg groups, respectively), suggesting that these doses of progesterone did not enhance memory for the platform location. Importantly, swim distances for the 20 mg/kg progesterone group did not significantly increase between Day 1 and Day 2 ($t(9) = 0.13, p > 0.05$; swim distance increased by only 35%), which suggests that this dose of progesterone prevented the memory decline observed in the other groups, thereby enhancing spatial memory consolidation. A repeated-measures ANOVA on the four trials of Day 2 revealed a trend toward a significant main effect of Trial ($F(3, 108) = 2.45, p = 0.06$), a significant main effect of Treatment ($F(3,36) = 6.57, p < 0.05$), but no Treatment $\times$ Trial interaction ($F(9,108) = 0.85, p > 0.05$), suggesting re-learning of the platform location on Day 2. To further examine whether mice in each group were able to re-learn the platform position, two analyses were performed. A paired-samples t-test including all groups examining swim distance between trials 1 and 4 on Day 2 revealed a significant decrease across trials ($t(39) = 2.96, p < 0.05$), suggesting that overall swim distances were significantly lower by the end of Day 2. Further, a one-way ANOVA conducted on swim distance during the last trial of Day 2 revealed no significant main effect of Treatment ($F(3,79) = 2.13, p > 0.05$), suggesting that the groups performed similarly by the end of Day 2. These findings support the conclusion that all groups could re-learn the platform position after initial reminder trials.

A repeated measures ANOVA examining swim speed during training on Day 1 (Figure 1D) revealed a significant main effect of Trial ($F(7, 245) = 28.61, p < 0.001$), a significant main effect of Treatment ($F(3,36) = 3.50, p < 0.05$), but no Treatment $\times$ Trial interaction ($F(21, 245) = 0.98, p > 0.50$). A one-way ANOVA performed on trial 8 revealed no differences in swim speed among groups ($F(3,39) = 0.91, p > 0.05$), demonstrating that, while some groups differed in swim speed during training trials 1–7, all groups decreased to a similar degree by trial 8. A repeated measures ANOVA examining swim speed during the last trial of Day 1 and the first trial of Day 2 revealed a significant main effect of Trial ($F(1, 36) = 146.86, p < 0.001$), no significant main effect of Treatment ($F(3,36) = 1.22, p > 0.05$), and no Treatment $\times$ Trial interaction ($F(3, 36) = .40, p > 0.05$), suggesting that all groups swam faster on the first trial of day 2 relative to the last trial of Day 1. A repeated measures ANOVA on the four trials of Day 2 revealed a significant main effect of Trial ($F(3, 108) = 17.83, p < 0.05$), no significant main effect of Treatment ($F(3,36) = 1.50, p > 0.05$), and no Trial $\times$ Treatment interaction ($F(9, 108) = .47, p > 0.05$) suggesting that all mice swam similarly as testing trials progressed. Lastly, a one-way ANOVA conducted on swim speed during the last trial of Day 2 revealed no significant effect of Treatment ($F(3,39) = 2.13, p > 0.05$). Together, these findings suggest that decreased swim speed resulting from repeated swim trials were similar among all groups,

and suggests that the effects of post-training progesterone treatment were not due to alterations in swimming ability.

3.2 Novel Object Recognition

3.2.1: Middle-Aged Mice—Time spent with the novel object during the choice phases of the object recognition task is presented in Figure 3. At the 24-hour delay (Figure 3A), both the 10 mg/kg ($t(8) = 2.75, p < 0.05$) and 20 mg/kg ($t(9) = 2.56, p < 0.05$) progesterone groups spent significantly more time with the novel object than chance. Neither the vehicle nor the 5 mg/kg groups showed a preference for the novel object, suggesting that only 10 and 20 mg/kg progesterone can enhance object memory after 24 hours. At the 48-hour delay (Figure 3C), none of the groups demonstrated a significant preference for the novel object, suggesting that progesterone has no beneficial effect on object memory consolidation after 48-hours in middle-aged mice. There were no differences in total time to complete the choice phase during the 24- ($F(3,40) = 0.55, p > 0.05$) or 48-hour ($F(3,41) = 1.87, p > 0.05$) delays, indicating no effect of progesterone on the total time taken to accumulate 30 seconds of exploration during the choice phase (Table 1).

3.2.2: Aged Mice—Time spent with the novel object in the choice phases of the object recognition task is presented in Figure 3. At the 24-hour delay (Figure 3B), the 10 mg/kg progesterone group spent significantly more time with the novel object than chance ($t(10) = 3.74, p < 0.05$). Neither the vehicle, 5 mg/kg progesterone, or 20 mg/kg progesterone groups showed a significant preference for the novel object, suggesting that only 10 mg/kg progesterone enhances object memory after 24 hours. At the 48-hour delay (Figure 3D), the vehicle and 20 mg/kg groups exhibited chance performance, whereas the 5 mg/kg ($t(7) = 3.31, p < 0.05$) and 10 mg/kg ($t(10) = 2.4, p < 0.05$) progesterone groups spent significantly more time with the novel object. This finding suggests that both 5 and 10 mg/kg progesterone can enhance object memory consolidation after 48 hours. There were no differences in total time to complete the choice phase during the 24- ($F(3,44) = 0.59, p > 0.05$) or 48-hour ($F(3,44) = 1.22, p > 0.05$) delays, indicating no effect of progesterone on the total time taken to accumulate 30 seconds of exploration during the choice phase (Table 2).

4. Discussion

The present results demonstrate that acute post-training i.p. progesterone administration differentially affects spatial memory consolidation in the Morris water maze and hippocampal-dependent object memory consolidation in middle-aged and aged female mice. The 10 and 20 mg/kg doses of progesterone enhanced object memory consolidation, relative to chance, in middle-aged females when tested after 24, but not 48 hours, and no dose affected spatial memory consolidation in the water maze. The lack of effect in the Morris water maze was likely due to generally poor learning and a lack of overnight forgetting in vehicle-treated mice, which has been observed previously in middle-aged mice (Benice et al., 2006). Thus, the possibility remains that acute, post-training progesterone administration may have had beneficial effects on memory consolidation in middle-aged mice if task acquisition had been sufficient to observe significant overnight forgetting. Interestingly, several studies have observed memory deficits in middle-aged rats and mice (e.g. Harburger et al., 2007; Francia et al., 2006; mild impairment observed in Verbitsky et al., 2004). Several factors could account for the discrepancy between the present and previous studies, including the number of days of training (1 vs. 5 or 8 days), the specific age at testing (“middle-aged” is generally defined as 12–16 months of age), and differences in endogenous steroid production prior to ovariectomy. It has been well established that middle-aged rodents demonstrate differences in the transition from normal estrous cycling to persistent estrous, to acyclicity (Frick et al., 2000). Thus, middle-aged mice may demonstrate differences in memory and/or responses to ovarian

hormones based on the length of time since the transition from normal cycling, which was not directly examined in the present study. This possibility may also underlie the high variability and generally poor performance of middle-aged females in the current study. The high variability is highlighted by the significant main effect of Treatment observed prior to hormone administration, such that the group assigned to receive vehicle demonstrated a steeper learning curve than any of the groups assigned to receive progesterone. If middle-aged females are transitioning from regular cycling to acyclicity, then the middle-aged females might be expected to demonstrate more variable behavior relative to aged females, who have been persistently acyclic for a period of time (Markowska, 1999).

Our object recognition findings seem to suggest that middle-aged females are less sensitive to the memory enhancing effects of progesterone than aged females. Only 10 and 20 mg/kg progesterone were effective in middle-aged mice at the 24 hour delay (aged mice responded to 5 mg/kg), whereas no effect was observed after 48 hours (aged mice respond to 5 and 10 mg/kg). The reasons for this apparent dose-response shift in the effects of progesterone at the two ages are unclear. One possible mechanism could be an increase in the number of progesterone receptors in response to decreased levels of circulating progesterone, thus increasing the sensitivity to progesterone in aged females. While we are unaware of any studies specifically examining age-related changes in progesterone receptor expression in learning-related brain regions, previous studies have demonstrated increased glutamate receptor expression in the hippocampus as a function of age. For example, Adams et al., (2001) demonstrated increased NR1 and NR2B expression in the ventral hippocampus of aged rats, relative to both young and middle-aged female rats. Further, the distribution of NR2B receptors in the hippocampus was altered by estrogen administration (Adams et al., 2004). Thus, changes in the number or distribution of progesterone receptors in the hippocampus of middle-aged or aged females could underlie the dose-response shift observed in the present study, but this has yet to be tested directly.

In aged mice, we observed enhanced object memory consolidation, relative to chance, after acute progesterone administration. These results are in accord with previous research in young ovariectomized female rats demonstrating that post-training i.p. injections of 4 or 8 mg/kg non water-soluble progesterone can enhance 24-hour object recognition, working memory in a Y-maze, and inhibitory avoidance (Frye and Lacey, 2000; Walf et al., 2006). Our findings extend these results to demonstrate that, in aged female mice, post-training injections of 5 mg/kg water-soluble progesterone can enhance object recognition, relative to chance, after 48 hours, whereas 10 mg/kg progesterone can enhance OR after both 24 and 48 hours. The highest dose of progesterone, 20 mg/kg, had no effect on object recognition, but significantly improved spatial memory consolidation. Interestingly, another recent study in aged female mice found that the 10 mg/kg dose had no effect on the ability of 0.2 mg/kg estradiol to enhance spatial memory in the water maze, whereas 20 mg/kg progesterone, completely blocked the memory-enhancing effects of estradiol (Harburger et al., 2007). Together, these data suggest that post-training progesterone is beneficial for memory consolidation in aged females when given alone, but detrimental when given in combination with a beneficial dose of estradiol.

Two potential issues of concern relating to the aged water maze data are worth addressing. First, the Trial $\times$ Treatment interaction in the analysis from trial 8 of Day 1 to trial 1 of Day 2 only approached significance ($p = 0.06$), which might indicate no effect of any progesterone dose. However, the conclusion that 20 mg/kg progesterone can enhance spatial memory consolidation is supported by t-tests showing that all groups except the 20 mg/kg group swam significantly longer distances on trial 1 of Day 2 than on trial 8 of Day 2. The fact that three of the four groups exhibited similar performance during these two trials likely overshadowed the positive effect of the 20 mg/kg dose in the omnibus test. The second issue is that although re-learning of the platform location did occur during the four trials on Day 2, the Trial main

effect for this analysis only approached significance ($p = 0.06$). This lack of statistical significance is likely due to high variability during Day 2, particularly in trials 1–3 (Figure 1B). It is interesting to note that both vehicle and the 20 mg/kg progesterone groups appear to have reacquired the task earlier during testing than the 5 and 10 mg/kg groups, although the reason for this difference is unclear. Nevertheless, a t-test indicated that swim distance decreased significantly on Day 2 between trials 1 and 4 which, combined with the non-significant Treatment main effect on the final trial of Day 2, suggests that all groups re-learned the platform position to a similar extent after initial reminder trials.

Interestingly, the data suggest a shift in the aged dose-response curve for progesterone was observed in the Morris water maze relative to the novel object recognition task, with the former requiring a higher dose to produce an improvement. Although it is unclear what might cause such a shift, one possibility may involve differences in hippocampal involvement in each task. Spatial memory in the Morris water maze can be severely impaired by lesions of the dorsal hippocampus encompassing 30–50% of total hippocampal volume, whereas ventral hippocampal lesions have little effect on water maze performance (reviewed in Moser & Moser, 1998). Novel object recognition memory, on the other hand, is only disrupted after lesions that damage 75–100% of the hippocampus (Broadbent et al., 2004). Thus, whereas spatial memory in the Morris water maze is critically dependent on the dorsal hippocampus, various regions of the hippocampus can support novel object memory (reviewed in Mumby, 2001). This differential regional involvement may contribute to the differential sensitivity of the tasks to progesterone, as progesterone receptors are expressed in all sub regions of the hippocampus (Guerra-Araiza et al., 2000; Kato et al., 1994). It is unknown how this expression changes with age, but it is likely that receptor levels in the hippocampus decrease somewhat with age, as progesterone receptor expression decreases with age in the hypothalamus (Chakraborty & Gore, 2004) and progestin levels are reduced in aged female mice relative to young female mice (Nelson et al., 1992). Nevertheless, the fact that a smaller region of the hippocampus is necessary for spatial memory in the water maze than for object recognition may render the water maze less sensitive to progesterone and, thus, necessitate a higher dose to observe a beneficial effect. The more widespread involvement of the hippocampus in object recognition could lead to the increased sensitivity of this task to progesterone if progesterone modulates performance by binding to more receptors throughout the hippocampus. This hypothesis, however, would need to be further supported by future studies mapping the distribution of progesterone receptors in the hippocampus to the sub regions critical for each task.

It should be noted that some previous studies have failed to demonstrate that progesterone alone improves memory in females. For example, a single progesterone injection given 4 hours prior to training in the Morris water maze had no significant effect on spatial memory in young female rats, although neither did a single injection of estradiol alone (Chesler and Juraska, 2000). In contrast, injection of both estradiol and progesterone significantly improved performance (Chesler and Juraska, 2000). In addition, young female mice chronically implanted with pellets releasing either progesterone alone or estrogen plus progesterone were impaired in learning a footshock avoidance task relative to mice receiving placebo or estrogen pellets (Farr et al., 1995). The most likely explanation for the discrepancy between these two studies and the improvements observed in the present and previous studies (Frye and Lacey, 2000; Walf et al., 2006) is the timing of progesterone administration. The studies reporting no effect or an impairing effect of progesterone administered the hormone prior to training, whereas the three studies reporting improvements administered progesterone post-training. This timing is important given that progesterone and its metabolites have anxiolytic and analgesic effects (Bitran, Hilvers, and Kellogg, 1991; Frye and Duncan, 1994), which could interfere with mnemonic function if given prior to training. Post-training paradigms, such as the one used in the present study, eliminate this confound because the hormone is not present in the circulation during training or testing; HBC-conjugated progesterone is metabolized

within 24 hours (Pitha and Pitha, 1985). Thus, although the present study cannot address the more clinically relevant issue of how chronic progesterone treatment influences memory function, it makes an important contribution towards understanding how progesterone specifically affects memory consolidation in the absence of the confounding factors that influence task performance.

In summary, the present study demonstrates that acute post-training progesterone injections can differentially affect spatial memory consolidation in the Morris water maze and hippocampal-dependent object memory consolidation in middle-aged and aged female mice. These findings provide an important initial step in clarifying the role of progesterone in memory consolidation.

Acknowledgements

This work was supported by NIH grant AG022525 to K.M.F. The author's would like to thank Dr. Zenab Amin and Jason Peterson for assistance with behavioral procedures.

References

- Akinci MK, Johnston GA. Sex differences in the effects of gonadectomy and acute swim stress on GABAA receptor binding in mouse forebrain membranes. *Neurochem Int* 1997;31:1–10. [PubMed: 9185158]
- Adams MM, Fink SE, Janssen WG, Shah RA, Morrison JH. Estrogen modulates synaptic N-methyl-D-aspartate receptor subunit distribution in the aged hippocampus. *J Comp Neurol* 2004;23:419–426. [PubMed: 15174084]
- Adams MM, Shah RA, Janssen WG, Morrison JH. Different modes of hippocampal plasticity in response to estrogen in young and aged female rats. *Proc Natl Acad Sci U S A* 2001;98:8071–8076. [PubMed: 11427724]
- Baker KB, Kim JJ. Effects of stress and hippocampal NMDA receptor antagonism on recognition memory in rats. *Learn. Mem* 2002;9:58–65. [PubMed: 11992016]
- Benice TS, Rizk A, Kohama S, Pfankuch T, Raber J. Sex differences in age-related cognitive decline in C57BL/6J mice associated with increased brain microtubule-associated protein 2 and synaptophysin immunoreactivity. *Neuroscience* 2006;137:413–423. [PubMed: 16330151]
- Bimonte HA, Denenberg VH. Estradiol facilitates performance as working memory load increases. *Psychoneuroendocrinology* 1999;24:161–173. [PubMed: 10101725]
- Bimonte-Nelson HA, Francis KR, Umphlet CD, Granholm AC. Progesterone reverses the spatial memory enhancements initiated by tonic and cyclic oestrogen therapy in middle-aged ovariectomized female rats. *Eur J Neurosci* 2006;24:229–242. [PubMed: 16882019]
- Bitran D, Hilvers RJ, Kellogg CK. Anxiolytic effects of 3 alpha-hydroxy-5 alpha[beta]-pregnan-20-one: endogenous metabolites of progesterone that are active at the GABAA receptor. *Brain Res* 1991;561:157–161. [PubMed: 1686744]
- Brinton RD. Cellular and molecular mechanisms of estrogen regulation of memory function and neuroprotection against Alzheimer's disease: recent insights and remaining challenges. *Learn. Mem* 2001;8:121–133. [PubMed: 11390632]
- Broadbent NJ, Squire LR, Clark RE. Spatial memory, recognition memory, and the hippocampus. *Proc. Nat. Acad. Sci* 2004;101:14515–14520. [PubMed: 15452348]
- Carlstrom L, Ke ZJ, Unnerstall JR, Cohen RS, Pandey SC. Estrogen modulation of the cyclic AMP response element-binding protein pathway. Effects of long-term and acute treatments. *Neuroendocrinology* 2001;74:227–243. [PubMed: 11598379]
- Chakraborty TR, Gore AC. Aging-related changes in ovarian hormones, their receptors, and neuroendocrine function. *Exp. Biol. Med* 2004;229:977–987.
- Chesler EJ, Juraska JM. Acute administration of estrogen and progesterone impairs the acquisition of the spatial Morris water maze in ovariectomized rats. *Horm. Behav* 2000;38:234–242. [PubMed: 11104641]

- Choi JM, Romeo RD, Brake WG, Bethea CL, Rosenwaks Z, McEwen BS. Estradiol increases pre- and post-synaptic proteins in the CA1 region of the hippocampus in female rhesus macaques (*Macaca mulatta*). *Endocrinology* 2003;144:4734–4738. [PubMed: 12960039]
- Clark RE, Zola SM, Squire LR. Impaired recognition memory in rats after damage to the hippocampus. *J. Neurosci* 2000;20:8853–8860. [PubMed: 11102494]
- Cyr M, Ghribi O, Thibault C, Morissette M, Landry M, Di Paolo T. Ovarian steroids and selective estrogen receptor modulators activity on rat brain NMDA and AMPA receptors. *Brain Res. Brain Res. Rev* 2001;37:153–161. [PubMed: 11744083]
- Daniel JM, Fader AJ, Spencer AL, Dohanich GP. Estrogen enhances performance of female rats during acquisition of a radial arm maze. *Horm. Behav* 1997;32:217–225. [PubMed: 9454673]
- Daniel JM, Hulst JL, Berbling JL. Estradiol replacement enhances working memory in middle-aged rats when initiated immediately after ovariectomy but not after a long-term period of ovarian hormone deprivation. *Endocrinology* 2006;147:607–614. [PubMed: 16239296]
- El-Bakri NK, Adem A, Suliman IA, Mulugeta E, Karlsson E, Lindgren JU, Winblad B, Islam A. Estrogen and progesterone treatment: effects on muscarinic M(4) receptor subtype in the rat brain. *Brain. Res* 2002;948:131–137. [PubMed: 12383964]
- Fader AJ, Hendricson AW, Dohanich GP. Estrogen improves performance of reinforced T-maze alternation and prevents the amnesic effects of scopolamine administered systemically or intrahippocampally. *Neurobiol. Learn. Mem* 1998;69:225–240. [PubMed: 9707487]
- Farr SA, Flood JF, Scherrer JF, Kaiser FE, Taylor GT, Morley JE. Effect of ovarian steroids on footshock avoidance learning and retention in female mice. *Physiol. Behav* 1995;58:715–723.
- Fernandez SM, Frick KM. Chronic oral estrogen affects memory and neurochemistry in middle-aged female mice. *Behav. Neurosci* 2004;118:1340–1351. [PubMed: 15598143]
- Foster TC, Sharrow KM, Kumar A, Masse J. Interaction of age and chronic estradiol replacement on memory and markers of brain aging. *Neurobiol. Aging* 2003;24:839–852. [PubMed: 12927766]
- Foy MR, Xu J, Xie X, Brinton RD, Thompson RF, Berger TW. 17beta-estradiol enhances NMDA receptor-mediated EPSPs and long-term potentiation. *J. Neurophysiol* 1999;81:925–929. [PubMed: 10036289]
- Francia N, Cirulli F, Chiarotti F, Antonelli A, Aloe I, Alleva E. Spatial memory deficits in middle-aged mice correlate with lower exploratory activity and a subordinate status: role of hippocampal neurotrophins. *Eur. J. Neurosci* 2006;23:711–728. [PubMed: 16487153]
- Frick KM, Burlingame LA, Arters JA, Berger-Sweeney J. Reference memory, anxiety and estrous cyclicity in C57BL/6NIA mice are affected by age and sex. *Neuroscience* 2000;95:293–307. [PubMed: 10619486]
- Frick KM, Fernandez SM, Bulinski SC. Estrogen replacement improves spatial reference memory and increases hippocampal synaptophysin in aged female mice. *Neuroscience* 2002;115:547–558. [PubMed: 12421621]
- Frick KM, Gresack JE. Sex differences in the behavioral response to spatial and object novelty in adult C57BL/6 mice. *Behav. Neurosci* 2003;117:1283–1291. [PubMed: 14674847]
- Frye CA, Duncan JE. Progesterone metabolites, effective at the GABAA receptor complex, attenuate pain sensitivity in rats. *Brain Res* 1994;643:194–203. [PubMed: 8032914]
- Frye CA, Lacey EH. Progestins influence performance on cognitive tasks independent of changes in affective behavior. *Psychobiology* 2000;28:550–563.
- Gibbs RB. Estrogen replacement enhances acquisition of a spatial memory task and reduces deficits associated with hippocampal muscarinic receptor inhibition. *Horm. Behav* 1999;36:222–233. [PubMed: 10603286]
- Gibbs RB. Long-term treatment with estrogen and progesterone enhances acquisition of a spatial memory task by ovariectomized aged rats. *Neurobiol. Aging* 2000;21:107–116. [PubMed: 10794855]
- Gresack JE, Frick KM. Environmental enrichment reduces the mnemonic and neural benefits of estrogen. *Neuroscience* 2004;125:459–471. [PubMed: 15381276]
- Gresack JE, Frick KM. Post-training estrogen enhances spatial and object memory consolidation in female mice. *Pharmacol. Biochem. Behav* 2006;84:112–119. [PubMed: 16759685]

- Guerra-Araiza C, Cerbon MA, Morimoto S, Camacho-Arroyo I. Progesterone receptor isoforms expression pattern in the rat brain during the estrous cycle. *Life. Sci* 2000;66:1743–1752. [PubMed: 10809171]
- Harburger LL, Bennett JC, Frick KM. Effects of estrogen and progesterone on spatial memory consolidation in aged females. *Neurobiol. Aging* 2007;28:602–610. [PubMed: 16621169]
- Heikkinen T, Puoliväli J, Liu L, Rissanen A, Tanila H. Effects of ovariectomy and estrogen treatment on learning and hippocampal neurotransmitters in mice. *Horm. Behav* 2002;41:22–32. [PubMed: 11863380]
- Johansson IM, Birzniece V, Lindblad C, Olsson T, Bäckström T. Allopregnanolone inhibits learning in the Morris water maze. *Brain Res* 2002;934:125–131. [PubMed: 11955475]
- Kato J, Hirata S, Nozawa A, Yamada-Mouri N. Gene expression of progesterone receptor isoforms in the rat brain. *Horm. Behav* 1994;28:454–463. [PubMed: 7729814]
- Kuroki Y, Fukushima K, Kanda Y, Mizuno K, Watanabe Y. Putative membrane-bound estrogen receptors possibly stimulate mitogen-activated protein kinase in the rat hippocampus. *Eur. J. Pharmacol* 2000;400:202–209.
- Luine V, Rodriguez M. Effects of estradiol on radial arm maze performance of young and aged rats. *Behav. Neural. Biol* 1994;62:230–236. [PubMed: 7857245]
- Markham JA, Pych JC, Juraska JM. Ovarian hormone replacement to aged ovariectomized female rats benefits acquisition of the morris water maze. *Horm. Behav* 2003;42:284–293. [PubMed: 12460588]
- Markowska AL. Sex dimorphisms in the rate of age-related decline in spatial memory: relevance to alterations in the estrous cycle. *J Neurosci* 1999;19:8122–8133. [PubMed: 10479712]
- Markowska AL, Savonenko AV. Effectiveness of estrogen replacement in restoration of cognitive function after long-term estrogen withdrawal in aging rats. *J. Neurosci* 2002;22:10985–10995. [PubMed: 12486194]
- Moser MB, Moser EI. Functional differentiation in the hippocampus. *Hippocampus* 1998;8:608–619. [PubMed: 9882018]
- Mumby DG. Perspectives on object-recognition memory following hippocampal damage: lessons from studies in rats. *Hippocampus* 2001;127:159–181.
- Mumby DG, Tremblay A, Lecluse V, Lehmann H. Hippocampal damage and anterograde object-recognition in rats after long retention intervals. *Hippocampus* 2005;15:1050–1056. [PubMed: 16145694]
- Murphy DD, Cole NB, Greenberger V, Segal M. Estradiol increases dendritic spine density by reducing GABA neurotransmission in hippocampal neurons. *J. Neurosci* 1998;18:2550–2559. [PubMed: 9502814]
- Murphy DD, Segal M. Progesterone prevents estradiol-induced dendritic spine formation in cultured hippocampal neurons. *Neuroendocrinology* 2000;72:133–143. [PubMed: 11025407]
- Nelson JF, Felicio LS, Osterburg HH, Finch CE. Differential contributions of ovarian and extraovarian factors to age-related reductions in plasma estradiol and progesterone during the estrous cycle of C57BL/6J mice. *Endocrinology* 1994;130:805–810. [PubMed: 1733727]
- Nilsen J, Brinton RD. Impact of progestins on estrogen-induced neuroprotection: synergy by progesterone and 19-norprogesterone and antagonism by medroxyprogesterone acetate. *Endocrinology* 2002;143:205–212. [PubMed: 11751611]
- Nilsen J, Brinton RD. Divergent impact of progesterone and medroxyprogesterone acetate (Provera) on nuclear mitogen-activated protein kinase signaling. *Proc. Natl. Acad. Sci* 2003;100:10506–10511. [PubMed: 12925744]
- O'Brien N, Lehmann H, Lecluse V, Mumby DG. Enhanced context-dependency of object recognition in rats with hippocampal lesions. *Behav. Brain. Res* 2006;170:156–162. [PubMed: 16580742]
- O'Neal MF, Means LW, Poole MC, Hamm RJ. Estrogen affects performance of ovariectomized rats in a two-choice water-escape working memory task. *Psychoneuroendocrinology* 1996;21:51–65.
- Packard MG, Teather LA. Intra-hippocampal estradiol infusion enhances memory in ovariectomized rats. *Neuroreport* 1997;8:3009–3013. [PubMed: 9331907]
- Pitha J, Harman SM, Michel ME. Hydrophilic cyclodextrin derivatives enable effective oral administration of steroidal hormones. *J. Pharm. Sci* 1986;75:165–167. [PubMed: 3958926]

- Pitha J, Pitha J. Amorphous water-soluble derivatives of cyclodextrins: nontoxic dissolution enhancing excipients. *J. Pharm. Sci* 1985;74:987–990. [PubMed: 4067854]
- Rissanen A, Puoliväli J, van Groen T, Riekkinen P Jr. In mice tonic estrogen replacement therapy improves non-spatial and spatial memory in a water maze task. *Neuroreport* 1999;10:1369–1372. [PubMed: 10363955]
- Sandstrom NJ, Williams CL. Memory retention is modulated by acute estradiol and progesterone replacement. *Behav. Neurosci* 2001;115:384–393. [PubMed: 11345963]
- Sherwin BB. Can estrogen keep you smart? Evidence from clinical studies. *J Psychiatry Neurosci* 1999;24:315–321. [PubMed: 10516798]
- Stone DJ, Rozovsky I, Morgan TE, Anderson CP, Finch CE. Increased synaptic sprouting in response to estrogen via an apolipoprotein E-dependent mechanism: Implications for Alzheimer's disease. *J. Neurosci* 1998;18:3180–3185. [PubMed: 9547226]
- Tanabe F, Miyasaka N, Kubota T, Aso T. Estrogen and progesterone improve scopolamine-induced impairment of spatial memory. *J Med Dent Sci* 2004;51:89–98. [PubMed: 15137470]
- Tanapat P, Hastings NB, Reeves AJ, Gould E. Estrogen stimulates a transient increase in the number of new neurons in the dentate gyrus of the adult female rat. *J. Neurosci* 1999;19:5792–5801. [PubMed: 10407020]
- Vaucher E, Reymond I, Najaffe R, Kar S, Quirion R, Miller MM, Franklin KB. Estrogen effects on object memory and cholinergic receptors in young and old female mice. *Neurobiol. Aging* 2002;23:87–95. [PubMed: 11755023]
- Verbitsky M, Yonan AL, Malleret G, Kandel ER, Giliam TC, Pavlidis P. Altered hippocampal transcript profile accompanies an age-related spatial memory deficit. *Learn Mem* 2004;11:253–260. [PubMed: 15169854]
- Vongher JM, Frye CA. Progesterone in conjunction with estradiol has neuroprotective effects in an animal model of neurodegeneration. *Pharmacol. Biochem, Behav* 1999;64:777–785. [PubMed: 10593201]
- Walf AA, Rhodes ME, Frye CA. Ovarian steroids enhance object recognition in naturally cycling and ovariectomized, hormone-primed rats. *Neurobiol. Learn. Mem* 2006;86:35–46. [PubMed: 16529958]
- Warren SG, Humphreys AG, Juraska JM, Greenough WT. LTP varies across the estrous cycle: enhanced synaptic plasticity in proestrus rats. *Brain. Res* 1995;703:26–30. [PubMed: 8719612]
- Woolley CS, McEwen BS. Estradiol mediates fluctuation in hippocampal synapse density during the estrous cycle in the adult rat. *J. Neurosci* 1992;12:2549–2554. [PubMed: 1613547]
- Woolley CS, McEwen BS. Roles of estradiol and progesterone in regulation of hippocampal dendritic spine density during the estrous cycle in the rat. *J. Comp. Neurol* 1993;336:293–306. [PubMed: 8245220]
- Yaffe K, Sawaya G, Lieberburg I, Grady D. Estrogen therapy in postmenopausal women: effects on cognitive function and dementia. *JAMA* 1998;279:688–695. [PubMed: 9496988]

Figure 1.

Effects of post-training progesterone injection on water maze performance in middle-aged (A & C) and aged (B & D) females. For all graphs, each point represents the group mean for each trial. Error bars represent $\pm$ standard error of the mean (SEM). The filled arrow represents the progesterone injection immediately after training trial 8. (A) Spatial memory consolidation in middle-aged mice. Swim distance in all middle-aged groups decreased over the eight training trials on Day 1. The groups did not differ on trial 8 of Day 1, suggesting that all groups performed similarly during the last trial before training. None of the groups demonstrated significant overnight forgetting after 24 hours, as evidenced by no significant increase in swim distance between trial 8 of Day 1 and trial 1 of Day 2. There were no differences among groups across the 4 testing trials of Day 2, likely due to a lack of forgetting after 24 hours. (B). Spatial memory consolidation in aged mice. Swim distance in all aged groups decreased over the eight training trials on Day 1. The groups did not differ in trial 8 of Day 1, suggesting similar performance during the last training trial. During the first trial of Day 2, swim distances in the vehicle, 5 mg/kg, and 10 mg/kg groups were significantly higher than during trial 8 of Day 1 (148%, 112%, and 84%, respectively). In contrast, the 35% increase in swim distance exhibited by the 20 mg/kg group was not significant, suggesting that this dose of progesterone can enhance spatial memory consolidation in aged female mice. Swim speeds were not affected by progesterone in either the middle-aged (C) or aged (D) mice. Within an age, swim speeds decreased similarly during Day 1 and were significantly faster on the 1st trial of Day 2 relative to the last trial of Day 1. Further, swim speeds decreased similarly in all groups over the four testing trials on Day 2, suggesting that alterations in swimming ability did not contribute to the differences observed in swim distance.

Figure 2.

Change in performance from trial 8 of Day 1 to trial 1 of Day 2 in middle-aged (A) and aged (B) females. (A) There were no significant differences in swim distance from Day 1 to Day 2 in any of the middle-aged groups. Among aged females, (B), vehicle-treated mice demonstrated significantly increased swim distances on the first trial of Day 2 relative to the last trial of Day 1. The 20 mg/kg dose of progesterone, but not 5 or 10 mg/kg progesterone, prevented this decline in performance.

Figure 3.

Effects of post-training progesterone injection on novel object memory consolidation in middle-aged (A & C) and aged (B & D) females. For all graphs, error bars represent $\pm$ SEM and $*$ = $p < 0.05$ relative to chance (dotted line at 15 s). (A) In middle-aged females both 10 mg/kg and 20 mg/kg progesterone increased the time spent with the novel object relative to chance at the 24 hour delay. In contrast, progesterone in middle-aged females (C) did not affect time spent with the novel object at the 48 hour delay, relative to chance. In aged females, ten mg/kg significantly increased the time spent with the novel object relative to chance after 24 hours (B). After the 48 hour delay (D), both 5 and 10 mg/kg doses increased the time that aged females spent with the novel object, relative to chance.

Table 1

Total time to accumulate 30 sec of object exploration in middle-aged mice.

Treatment	24 hours	Delay
		48 hours
Vehicle (N = 10)	457.1 ± 114.3	437.8 ± 123.9
5 mg/kg (N = 9)	532.2 ± 110.7	568.4 ± 119.8
10 mg/kg (N = 10)	589.5 ± 124.6	704.1 ± 136.6
20 mg/kg (N = 10)	472.2 ± 121.2	464.4 ± 133.0

Values represent the mean ± SEM.

Table 2

Total time to accumulate 30 sec of object exploration in aged mice.

Treatment	24 hours	Delay
		48 hours
Vehicle (N = 9)	476.5 ± 89.1	511.5 ± 120.5
5 mg/kg (N = 10)	374.2 ± 92.1	598.0 ± 111.0
10 mg/kg (N = 11)	436.0 ± 84.7	369.1 ± 116.8
20 mg/kg (N = 10)	382.6 ± 96.2	452.5 ± 91.3

Values represent the mean ± SEM.