Published in final edited form as:

Org Lett. 2007 January 18; 9(2): 279–282. doi:10.1021/ol062728b.

Total Synthesis of (±)-Strychnine via a [4+2]-Cycloaddition/ Rearrangement Cascade

Hongjun Zhang, Jutatip Boonsombat, and Albert Padwa

Department of Chemistry, Emory University, Atlanta, Georgia 30322, chemap@emory.edu

Abstract

A new strategy for the synthesis of the *Strychnos* alkaloid (±)-strychnine has been developed and is based on an intramolecular [4+2]-cycloaddition/rearrangement cascade of an indolyl substituted amidofuran. The critical D-ring was assembled by an intramolecular palladium catalyzed enolate-driven cross-coupling of an *N*-tethered vinyl iodide.

Strychnos alkaloids belonging to the curane type constitute an important group of architecturally complex and widely distributed monoterpenoid indole alkaloids.^{1,2} The curane family is characterized by the presence of a pentacyclic 3,5-ethanopyrrolo[2,3-*d*]carbazole framework (*i.e.*, **1**) bearing a two-carbon appendage at C-20 and an oxidized one-carbon substituent (C-17) at the C-16 position (Figure 1).³ The synthesis of various members of the *Strychnos* family has been an area of interest ever since the structural elucidation of strychnine (**4**), the most famous of this group of alkaloids, by Robinson in 1946.⁴ Strychnine was first isolated in 1818 from the Indian poison nut⁵ (*Strychnos nux vomica*) and possesses highly toxic properties as a result of its interaction with the glycine receptor site thereby blocking the flux of chloride ions and disruption of nerve-cell signaling.⁶ Its complex heptacyclic structure, containing 24 skeletal atoms and six contiguous stereogenic centers, has fascinated organic chemists for the past sixty years. Nearly 40 years after Woodward's pioneering achievement of strychnine⁷, a number of other research groups have reported on its synthesis.⁸

Efficient approaches toward the *Strychnos* pentacyclic framework would allow not only the synthesis of other members of this family of natural products (*i.e.*, strychnopivotine (**2**) and akuammicine (**3**)), but also related non-natural products possessing biological activity. Along these lines, we have recently become involved in the development and optimization of a new approach for the construction of the pentacyclic framework of the *Strychnos* system. In this communication, we report a concise stereocontrolled total synthesis of (±)-strychnine (**4**) wherein an efficient [4+2]-cycloaddition/rearrangement method previously developed in our laboratories plays a crucial role.⁹

In Rawal and Iwasa's elegant synthesis of (±)-strychnine,^{8e} an ingenious palladium-catalyzed intramolecular Heck reaction was used as the key step for the creation of the D-ring. As in Woodward's original approach,⁷ isostrychnine (**5**) was the final critical intermediate used in this synthesis. Its Prelog-Taylor cyclization to strychnine¹⁰ (Scheme 1), however, suffers from an unfavorable 3:1-equilibration ratio of these two compounds. From this perspective, the alternative biomimetic route to strychnine involving condensation of the Wieland-Gumlich aldehyde (**6**) with an acetate equivalent for the formation of the G ring seemed to us to be the more attractive approach,¹¹ as it avoids the unfavorable equilibration ratio. As illustrated in Scheme 2, our retrosynthetic analysis of strychnine (**4**) relies upon the efficient construction

of the tetracyclic sub-structure **8**, containing the ABCE rings of **4**, by an intramolecular [4+2]-cycloaddition/rearrangement cascade of 2-amido-furan **7**. The synthetic plan also involves closure of the D-ring by a palladium-catalyzed intramolecular coupling of an amido tethered vinyl iodide with a keto enolate (*i.e.* **9** $\rightarrow$ **10**).^{12,13}

Some years ago, we commenced a program of research based on the intramolecular [4+2]-cycloaddition/rearrangement cascade of 2-amidofurans as a strategy for the synthesis of different classes of alkaloids.¹⁴ Intramolecular cycloaddition reactions often benefit from higher reactivity and greater control of stereoselectivity relative to their intermolecular counterparts. Specifically, connecting the two π -fragments *via* a “tether” generally facilitates the rate of the [4+2]-cycloaddition reaction. Our earlier experience with this domino sequence prompted us to apply the cascade approach toward ($\pm$)-strychnine. With this in mind, we prepared furanyl indole **7** by acylation of the mixed anhydride of 2-(1-acetyl-1*H*-indol-3-yl) acetic acid¹⁵ with the *N*-lithiate anion of *tert*-butyl furan-2-ylcarbamate followed by Boc deprotection¹⁶ with $\text{Mg}(\text{ClO}_4)_2$ and subsequent *N*-alkylation using 1-(bromomethyl)-2-methylbenzene in 55% overall yield for the 3-step procedure. We suspect that the presence of the large *o*-methylbenzyl group on the amido nitrogen atom causes the reactive *s*-trans rotamer to be more highly populated and probably helps promote the [4+2]-intramolecular cycloaddition. Indeed, the cycloaddition/rearrangement cascade of **7** was remarkably efficient given that two heteroaromatic systems are compromised in the reaction. Thus, heating a sample of **7** at 150 °C (toluene) in a microwave reactor for 30 min in the presence of catalytic MgI_2 afforded the desired aza-tetracycle **8** in 95% yield (Scheme 3). This reaction cascade can be rationalized by a nitrogen-assisted ring opening of the initially formed cycloadduct **11** to produce *N*-acyliminium ion **12**. A subsequent deprotonation followed by ketonization of the resulting enol accounts for the formation of **8**.

Our synthesis of the more advanced tetracycle **9** required for eventual D-ring cyclization of strychnine began by stereoselective reduction of the keto group of **8** with NaBH_4 followed by *N*-deacetylation using NaOMe (Scheme 4). The γ -lactam carbonyl group was then reduced with LiAlH_4 and the resulting enamine was further reduced using $\text{NaBH}(\text{OAc})_3$ to give alcohol **13** as the major diastereomer in 65% yield over the four-step sequence.¹⁷ Removal of the *o*-methylbenzyl group by catalytic hydrogenation furnished **14** in 70% isolated yield. This was followed by *N*-alkylation with *Z*-1-bromo-2-iodo-4-(methoxymethoxy)-but-2-ene¹⁸ to give alcohol **15** in 80% yield. Condensation of the indoline nitrogen present in **15** with 2,4-dimethoxybenzaldehyde in the presence of $\text{NaBH}(\text{OAc})_3$ ¹⁹ afforded the *N*-protected 2,4-dimethoxy-benzylamine (DMB) in 86% yield.²⁰ Oxidation of the resulting secondary alcohol to the corresponding ketone **9** occurred smoothly in 80% yield using tetrapropyl-ammonium perruthenate (TPAP).²¹ The stage was now set for the completion of the synthesis. The critical D-ring of strychnine was constructed by a palladium-catalyzed intramolecular coupling of the amino-tethered vinyl iodide with the keto enolate derived from **9**. The key palladium-catalyzed cyclization was carried out on a sample of **9** using $\text{Pd}(\text{PPh}_3)_4$ and PhOK . Gratifyingly, the reaction proceeded smoothly to furnish aza-pentacycle **16** in 56% unoptimized yield.²²

Our initial attempts to convert the keto group of **16** into the corresponding enol ether **10** using methoxymethylene-triphenylphosphorane ($\text{MeOCH}=\text{PPh}_3$) were unsuccessful, probably as a consequence of steric congestion about the ketone. Consequently, we turned our attention to the phosphine oxide reagent $\text{MeOCH}_2\text{P}(\text{O})\text{Ph}_2$ whose anion is sterically less demanding and more nucleophilic compared to the phosphorane $\text{MeOCH}=\text{PPh}_3$.^{23,24} Thus, treatment of $\text{MeOCH}_2\text{P}(\text{O})\text{Ph}_2$ with LDA in THF gave the lithio anion which reacted smoothly with ketone **16** at 0 °C to provide **10** as a single diastereomer in 72% isolated yield (Scheme 5). The last major hurdle involved an acid promoted deprotection/hydrolysis of **10** into the Wieland-Gumlich aldehyde (**6**). The hydrolysis was satisfactorily accomplished by the treatment of **10** with 3*N* HCl in THF at 55 °C for 10 h which gave **6** in 54% yield. By using shorter reaction

times and following the reaction by NMR spectroscopy, we found that the initially formed primary alcohol **17** undergoes a subsequent deprotection of the DMB group to furnish enol ether **18**. Further stirring of **18** under the acidic conditions eventually furnished the Wieland-Gumlich aldehyde (**6**). Although the conversion of **6** into strychnine had been reported by Robinson in 1953,¹¹ we decided to reproduce the described protocol for the sake of a complete total synthesis. Thus, the final biomimetic condensation of **6** with malonic acid, acetic anhydride, sodium acetate, and acetic acid provided ($\pm$)-strychnine (**4**) in 80% yield and with a 4.4% overall yield for the 13-step reaction sequence starting from furanyl indole **7**.

In summary, a concise synthesis of the *Strychnos* alkaloid ($\pm$)-strychnine is reported. A central step in the synthesis consists of an intramolecular [4+2]-cycloaddition/rearrangement cascade of an indolyl substituted amidofuran which delivers an aza-tetracyclic sub-structure containing the ABCE-rings of the natural product. Closure of the remaining D-ring was carried out from the aza-tetracyclic intermediate by an intramolecular palladium catalyzed enolate-driven cross-coupling between the *N*-tethered vinyl iodide and the keto group. The short sequence used to synthesize ($\pm$)-strychnine should also provide rapid entry into other *Strychnos* alkaloid natural products and is currently under active investigation in our laboratories.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

Dedicated to my colleague and friend David Goldsmith on the occasion of his 75th birthday. We appreciate the financial support provided by the National Institutes of Health (GM 059384) and the National Science Foundation (CHE-0450779). We wish to thank Professor Larry Overman (UC Irvine) for providing a proton NMR spectrum of the Wieland-Gumlich aldehyde (**6**) for comparison purposes.

References

1. For leading reviews of Strychnos alkaloids, see: (a) Bosch J, Bonjoch J, Amat M. The Alkaloids 1996;48:75. and references cited therein (b) Bonjoch J, Solé D. Chem Rev 2000;100:3455. [PubMed: 11777429]
2. Creasey, WA. The Monoterpene Indole Alkaloids. Saxton, JE., editor. Mterscience; New York: 1983. p. 800
3. The biogenetic numbering is used for pentacyclic structures such as **1**, see: Le Men J, Taylor WI. Experientia 1965;21:508. [PubMed: 5872609]
4. (a) Robinson R. Experientia 1946;2:28. (b) Holmes HL, Openshaw HT, Robinson R. J Chem Soc 1946:908. (c) Openshaw HT, Robinson R. Nature 1946;157:438.
5. Pelletier PJ, Caventou JB. Ann Chim Phys 1818;8:323.
6. Aprison, MH. Glycine Neurotransmission. Otterson, OP.; Strom-Mathisen, J., editors. Wiley; New York: 1990. p. 1-23. (b) Grenningloh G, Rienitz A, Schmitt B, Methfessel C, Zensen M, Beyreuther K, Gundelfinger ED, Betz H. Nature 1987;328:215. [PubMed: 3037383] (c) Kleckner NW, Dingledine R. Science 1988;241:835. [PubMed: 2841759]
7. Woodward RB, Cava MP, Ollis WD, Hunger A, Daeniker HU, Schenker K. J Am Chem Soc 1954;76:4749. Woodward RB, Cava MP, Ollis WD, Hunger A, Daeniker HU, Schenker K. Tetrahedron 1963;19:247.
8. (a) Magnus P, Giles M, Bonnert R, Kim CS, McQuire L, Merritt A, Vicker N. J Am Chem Soc 1992;114:4403. Magnus P, Giles M, Bonnert R, Johnson G, McQuire L, Deluca M, Merritt A, Kim CS, Vicker N. J Am Chem Soc 1993;115:8116. Stork, G. Presented at the Ischia Advanced School of Organic Chemistry; Ischia Porto, Italy. September 21, 1992; (c) Knight SD, Overman LE, Pairaudeau G. J Am Chem Soc 1993;115:9293. Knight SD, Overman LE, Pairaudeau G. J Am Chem Soc 1995;117:5776. (d) Kuehne ME, Xu F. J Org Chem 1993;58:7490. Kuehne ME, Xu F. J Org Chem 1998;63:9427. (e) Rawal VH, Iwasa S. J Org Chem 1994;59:2685. (f) Solé D, Bonjoch J, García-Rubio

- S, Peidró E, Bosch J. *Angew Chem, Int Ed* 1999;38:395. Solé D, Bonjoch J, García-Rubio S, Peidró E, Bosch J. *Chem-Eur J* 2000;6:655. (g) Eichberg MJ, Dorta RL, Lamottke K, Vollhardt KPC. *Org Lett* 2000;2:2479. [PubMed: 10956526] Eichberg MJ, Dorta RL, Grotjahn DB, Lamottke K, Schmidt M, Vollhardt KPC. *J Am Chem Soc* 2001;123:9324. [PubMed: 11562215] (h) Ito M, Clark CW, Mortimore M, Goh JB, Martin SF. *J Am Chem Soc* 2001;123:8003. [PubMed: 11506556] (i) Nakanishi M, Mori M. *Angew Chem, Int Ed* 2002;41:1934. Mori M, Nakanishi M, Kajishima D, Sato Y. *J Am Chem Soc* 2003;125:9801. [PubMed: 12904045] (j) Bodwell GJ, Li J. *Angew Chem, Int Ed* 2002;41:3261. (k) Ohshima T, Xu Y, Takita R, Shimizu S, Zhong D, Shibasaki M. *J Am Chem Soc* 2002;124:14546. [PubMed: 12465959] Ohshima T, Xu Y, Takita R, Shibasaki M. *Tetrahedron* 2004;60:9569. (l) Kaburagi Y, Tokuyama H, Fukuyama T. *J Am Chem Soc* 2004;126:10246. [PubMed: 15315428]
9. (a) Wang Q, Padwa A. *Org Lett* 2004;6:2189. [PubMed: 15200317] (b) Padwa A, Ginn JD. *J Org Chem* 2005;70:5197. [PubMed: 15960524] (c) Padwa A, Bur SK, Zhang H. *J Org Chem* 2005;70:6833. [PubMed: 16095303] (d) Zhang H, Padwa A. *Org Lett* 2006;8:247. [PubMed: 16408886]
10. Prelog V, Battagay J, Taylor WI. *Helv Chim Acta* 1948;31:2244. [PubMed: 18100059]
11. Anet FAL, Robinson R. *Chem Ind* 1953:245. See also references 8a and 8c
12. (a) Solé D, Peidró E, Bonjoch J. *Org Lett* 2000;2:2225. [PubMed: 10930249] (b) Solé D, Diaba J, Bonjoch J. *J Org Chem* 2003;68:5746. [PubMed: 12839475] (c) Bonjoch J, Diaba F, Puigbó G, Peidró E, Solé D. *Tetrahedron Lett* 2003;44:8387. (d) Solé D, Urbaneja X, Bonjoch J. *Adv Synth Catal* 2004;346:1646. (e) Solé D, Urbaneja X, Bonjoch J. *Org Lett* 2005;7:5461. [PubMed: 16288531]
13. For some other examples of intramolecular Pd-catalyzed vinylations, see: (a) Piers E, Oballa RM. *Tetrahedron Lett* 1995;36:5857. (b) Yu J, Wang T, Liu X, Deschamps J, Flippen-Anderson J, Liao X, Cook JM. *J Org Chem* 2003;68:7565. [PubMed: 14510528]
14. (a) Padwa A, Brodney MA, Dimitroff M. *J Org Chem* 1998;63:5304. (b) Bur SK, Lynch SM, Padwa A. *Org Lett* 2002;4:473. [PubMed: 11843569] (c) Ginn JD, Padwa A. *Org Lett* 2002;4:1515. [PubMed: 11975617]
15. Padwa A, Brodney MA, Lynch SM, Rashatasakhon P, Wang Q, Zhang H. *J Org Chem* 2004;69:3735. [PubMed: 15153003]
16. (a) Young FG, Frostick FC, Sanderson JJ, Hauser CR. *J Am Chem Soc* 1950;72:3635. (b) Mao C-L, Frostick FC, Man EH, Manyik RM, Wells RL, Hauser CR. *J Org Chem* 1969;34:1425.
17. Attempts to introduce the vinyl iodide moiety earlier in the synthesis were unsuccessful due to the competitive loss of the iodo group during the reduction step.
18. The synthesis of the required allylic bromide is described in the supplemental section.
19. Abdel-Magid AF, Carson KG, Harris BD, Maryanoff CA, Shah RD. *J Org Chem* 1996;61:3849. [PubMed: 11667239]
20. The DMP group was selected as it was readily removed under the acidic conditions employed in the hydrolysis of 10 to the Wieland-Gumlich aldehyde (6) (see Scheme 5).
21. Ley SV, Norman J, Griffith WP, Marsden SP. *Synthesis* 1994:639.
22. A similar approach was successfully employed in a synthesis of strychnopivotine (2) and this will be described at a later date.
23. Earnshaw C, Wallis CJ, Warren S. *J Chem Soc, Perkin Trans 1* 1979:3099.
24. Patel DV, Schmidt RJ, Gordon EM. *J Org Chem* 1992;57:7143.

1; curane skeleton

2; strychnopivotine

3; akuammicine

4; strychnine

Figure 1.
Some representative *Strychnos* alkaloids

Scheme 1.

Scheme 2.

Scheme 3.

Scheme 4.

Scheme 5.