Case Report

Il Choi, M.D.

Soo-Bin Im, M.D.

Bum-Tae Kim, M.D.

Won-Han Shin, M.D.

Department of Neurosurgery
College of Medicine
Soonchunhyang University
Bucheon, Korea

Radiculopathy Caused by Internal Iliac Artery Pseudoaneurysm Managed with Endovascular Embolization

The authors describe a case of pseudoaneurysm arising from internal iliac artery presented with radiculopathy mimicking the symptoms of lumbar disc disease or spinal cord tumor. Among the several preoperative evaluation including CT, MRI, electrophysiologic study and ultrasonography, important diagnostic clue was obtained by ultrasonographic findings of turbulence flow at the core of partially enhanced mass in the pelvic cavity. The patient was managed with endovascular coil embolization successfully. The current case makes us remind that assessment of neurological symptoms on lower extremity should include consideration of extraspinal cause in pelvis.

KEY WORDS : Pseudoaneurysm · Internal iliac artery · Radiculopathy · Endovascular embolization.

INTRODUCTION

The most common cause of the neurological symptoms on the lower extremities comes from the area around the spinal canal or spinal root. When the clinicians examine such patients, the lumbar disc herniation or spinal cord tumor are initially considered. However, a lesion on extraspinal area could be a cause of the neurological presentation on lower extremity^{1,7,10)}. The authors describe a case of pseudoaneurysm in the pelvic cavity presenting radiating pain and motor weakness on lower extremity like symptoms of lumbar disc herniation or spinal cord tumor.

CASE REPORT

A 58-year-old woman presented with radiating pain on L5 and S1 dermatomes and weakness of right lower extremity, especially the ankle dorsiflexion grade II. The symptom had started one year before and aggravated gradually. On lumbar and thoracic spine magnetic resonance image (MRI), however we could not find any lesion which could explain her neurological symptoms (Fig. 1). The electrophysiologic examination showed multiple lumbosacral radiculopathy such as cauda equina syndrome mainly on the right side. She had a history of high-dose pelvic irradiation for the cervical cancer of uterus 10 years ago. Pelvis MRI, checked at gynecology department, showed a 5.5 × 7 cm sized heterogeneously enhancing mass on the left presacral area. The mass eroded the sacral bone, resulting

Fig. 1. Lumbar and thoracic spine T2-weighted magnetic resonance image showing no abnormality on thoracolumbar area.

- Received : August 10, 2007
- Accepted : November 12, 2007
- Address for reprints :
Soo-Bin Im, M.D.
Department of Neurosurgery
College of Medicine
Soonchunhyang University
1174 Jung-dong, Wonmi-gu
Bucheon 420-767, Korea
Tel : +82-32-621-5290
Fax : +82-32-621-5016
E-mail : isbrzw@sch.ac.kr

Fig. 2. The pelvic magnetic resonance image showing 5.5×7 cm sized partially enhancing heterogenous mass which is eroding the sacral bone and widening the right S1 neural foramen.

Fig. 3. Ultrasonography on pelvis area showing turbulent vascular flow in the center of mass.

Fig. 4. Angiography showing pseudoaneurysm (A) and after coil embolization at the neck of pseudoaneurysm and proximal branch of right internal iliac artery (B).

in widening of the S1 neural foramen (Fig. 2). Initial radiological impression was ovarian tumor or tumor of neural origin. Further evaluation with ultrasonography was performed to obtain more information about the mass, preoperatively. The turbulent vascular flow was detected in the central part of the mass by ultrasonography suggesting pseudoaneurysm (Fig. 3). We changed treatment plan from direct surgery to endovascular embolization. On aortography approached from the left common femoral artery, a 5 cm sized pseudoaneurysm supplied from the right internal iliac artery was found. The proximal branch of right internal iliac artery and neck of

Fig. 5. Pelvic computed tomography checked 2 months (A) and 1 year (B) after embolization showing significant decreased size of mass.

pseudoaneurysm was embolized with 8 coils of 10 mm and 12 mm rolling diameter delivered from the 5 F DAV's catheter (Fig. 4). After embolization, the radiating pain was improved immediately and the motor weakness on the right lower extremity recovered gradually during 3 months except weakness of ankle dorsiflexion. The weakness of ankle dorsiflexion was not improved significantly until one year follow up. The follow-up computed tomography, obtained 2 months and 1 year after embolization, showed significant decrease in the size of pseudoaneurysm (Fig. 5).

DISCUSSION

Although radiculopathy is usually attributed to degenerative disc disease such as a herniated intervertebral disc in the lower lumbar spine, it may be related by other less common causes. The reported cases of extraspinal cause of lumbosacral radiculopathy were sporadic and overall incidence was seldom reported. Kleiner et al.¹⁰ reported that the incidence of extraspinal cause of lumbosacral radiculopathy was about 0.09% from the reviewing study of 12,125 patients who had spinal disorder.

Recently reported extraspinal causes include tumor of bone or soft tissue^{1,10,12,13,18}, pseudoaneurysm or aneurysm of the internal iliac artery⁷, inferior gluteal artery¹⁵, abdominal aorta¹⁹ or other pelvic artery³, and compartment syndrome or entrapment neuropathy¹¹. The various iatrogenic causes of pseudoaneurysm or arteriovenous fistulas in the pelvis were also reported, such as transvaginal needle biopsy¹⁵, renal transplantation^{6,8,14}, lumbar disc surgery⁹, and radiation therapy^{4,18} like current case.

The vascular lesion in the deep pelvic cavity could present radiating pain and motor weakness mimicking symptoms of herniation of lumbar disc or spinal cord tumor. The importance of clinical history and examination in the diagnosis of these lesions cannot be overemphasized. The current case makes us remind that assessment of neurological symptoms on lower extremity include consideration of

Fig. 6. Retrograde reviewing of lumbar MRI showing presacral mass (arrow) which was difficult to distinguish from the pelvic content.

extraspinal cause though it is rare.

Reviewing the previous lumbar spine MRI retrospectively, we could find out the mass at the extreme parasagittal section. However, it was difficult to distinguish partially sectioned mass from the pelvic organ or bowel signal. Moreover, the routine axial section didn't cover to the bottom of sacrum (Fig. 6).

MR imaging with

CT or MR angiography may not provide a valuable data for diagnosis in chronic contained pseudoaneurysm filled with blood products. For evaluation of mass or vascular lesion in the deep pelvic structure, the role of ultrasonography should be emphasized. In this case, the ultrasonography provided crucial clue in the diagnosis and the decision for the treatment plan. Treatment strategies for the pseudoaneurysm in the deep pelvis could be either surgical excision or endovascular technique. In some of surgery cases, however, because of incorrect preoperative diagnosis, the surgeons may be confronted with severe and unexpected hemorrhage and the surgery can be aborted without effective treatment^{2,4,18}. Also, ligation of proximal artery during surgery may cause ischemic necrosis of leg¹⁸. Many of successful endovascular treatments for internal iliac artery pseudoaneurysm have been reported^{4,5,9,16,17}. Based on ultrasonographic diagnosis, we changed treatment plan from surgical excision to endovascular embolization that could avoid potentially lethal situation.

CONCLUSION

The vascular lesion in the deep pelvic cavity could present radiating pain and motor weakness mimicking herniation of lumbar disc or spinal cord tumor. The ultrasonographic

evaluation and endovascular embolization are considered as valuable tools for diagnosis and management for the vascular lesion for the deep pelvic area.

References

1. Bickels J, Kahanovitz N, Rubert CK, Henshaw RM, Moss DP, Meller I, et al : Extraspinal bone and soft-tissue tumors as a cause of sciatica. Clinical diagnosis and recommendations : analysis of 32 cases. *Spine* 24 : 1611-1616, 1999
2. Buckley JG, Salimi Z, George E : Late onset renal allograft anastomotic pseudoaneurysm with absent Doppler signal. *J Urol* 148 : 392-394, 1992
3. Chapman EM, Shaw RS, Kubik CS : Sciatic pain from arteriosclerotic aneurysm of pelvic arteries *N Engl J Med* 271 : 1410-1411, 1964
4. de Baere T, Ousehal A, Kuoch V, Sapoval M, Lagrange C, Roche A : Endovascular management of bleeding iliac artery pseudoaneurysms complicating radiation therapy for pelvic malignancies. *AJR Am J Roentgenol* 170 : 349-353, 1998
5. de Godoy JM, Reis LF, Casagrande M, Braile DM : Treatment of internal iliac artery pseudoaneurysm by an endovascular approach. *J Chin Med Assoc* 68 : 435-436, 2005
6. Donckier V, De Pauw L, Ferreira J, Hanquinet S, Hooghe L, Janssen F, et al : False aneurysm after transplant nephrectomy. Report of two cases. *Transplantation* 60 : 303-304, 1995
7. Dudeney S, O'Farrell D, Bouchier-Hayes D, Byrne J : Extraspinal causes of sciatica. A case report. *Spine* 23 : 494-496, 1998
8. Fleschner NE, Johnston KW : Repair of an autotransplant renal artery aneurysm : case report and literature review. *J Urol* 148 : 389-391, 1992
9. Jeong CA, Park KH, Jo YG : An iliac arteriovenous fistula and pseudoaneurysm after lumbar disc surgery. *J Korean Neurosurg Soc* 36 : 415-418, 2004
10. Kleiner JB, Donaldson WF 3rd, Curd JG, Thorne RP : Extraspinal causes of lumbosacral radiculopathy. *J Bone Joint Surg Am* 73 : 817-821, 1991
11. Kopell HP, Thompson WA, Postel AH : Entrapment neuropathy of the ilioinguinal nerve. *N Engl J Med* 266 : 16-19, 1962
12. Ladha SS, Spinner RJ, Suarez GA, Amrami KK, Dyck PJ : Neoplastic lumbosacral radiculoplexopathy in prostate cancer by direct perineural spread : an unusual entity. *Muscle Nerve* 34 : 659-665, 2006
13. Larson J, O'Malley JE, Cohen TI : Case report : solitary pelvic osteochondroma presenting with L3 nerve root compression. *Alaska Med* 44 : 35-37, 2002
14. Luzzio CC, Waclawik AJ, Gallagher CL, Knechtle SJ : Iliac artery pseudoaneurysm following renal transplantation presenting as lumbosacral plexopathy. *Transplantation* 67 : 1077-1078, 1999
15. Papadopoulos SM, McGillicuddy JE, Messina LM : Pseudoaneurysm of the inferior gluteal artery presenting as sciatic nerve compression. *Neurosurgery* 24 : 926-928, 1989
16. Parodi JC : Endovascular repair of aortic aneurysms, arteriovenous fistulas, and false aneurysms. *World J Surg* 20 : 655-663, 1996
17. Todd GJ, DeRose JJ, Martin EC : Complementary surgical/interventional techniques for nonresective management of "inoperable" aneurysms : a second look. *Ann Vasc Surg* 12 : 248-254, 1998
18. Vetto JT, Culp SC, Smythe TB, Chang AE, Sindelar WF, Sugarbaker PH, et al : Iliac arterial-enteric fistulas occurring after pelvic irradiation. *Surgery* 101 : 643-647, 1987
19. Wilberger Jr JE : Lumbosacral radiculopathy secondary to abdominal aortic aneurysms. Report of three cases. *Neurol Clin* 58 : 965-967, 1983