P_{BAD}-Based Shuttle Vectors for Functional Analysis of Toxic and Highly Regulated Genes in *Pseudomonas* and *Burkholderia* spp. and Other Bacteria[▽]

Dongru Qiu,^{1,‡} F. Heath Damron,^{1,‡} Takehiko Mima,⁴ Herbert P. Schweizer,⁴ and Hongwei D. Yu^{1,2,3,*}

Departments of Biochemistry and Microbiology¹ and Pediatrics,² Joan C. Edwards School of Medicine at Marshall University, Huntington, West Virginia 25755-9320; Progenesis Technologies, LLC, Bldg. 740, Rm. 4136, Dow Technology Park, 3200 Kanawha Turnpike, South Charleston, West Virginia 25303³; and Department of Microbiology, Immunology and Pathology, Rocky Mountain Regional Center of Excellence for Biodefense and Emerging Infectious Diseases Research, Colorado State University, Fort Collins, Colorado 80523-1690⁴

Received 18 June 2008/Accepted 6 October 2008

We report the construction of a series of *Escherichia-Pseudomonas* broad-host-range expression vectors utilizing the P_{BAD} promoter and the *araC* regulator for routine cloning, conditional expression, and analysis of tightly controlled and/or toxic genes in pseudomonads.

Gene cloning, disruption, deletion, complementation analysis, and allelic exchange are central to prokaryotic molecular genetics. In *Pseudomonas aeruginosa*, Schweizer and colleagues developed the pUCP family of general-purpose vectors for cloning and gene expression (24, 29) based on the well-characterized pUC18/19 vectors (32) and the cryptic mini-plasmid pRO1614 (19). Other promoters are also in routine use, such as the *tac* (4, 6), T7 (28), and *araBAD* promoter-based (8, 11) vectors for regulated expression in *Escherichia coli* and many other bacterial species (e.g., see references 2, 18, and 25). In *E. coli*, AraC represses the *araBAD* promoter (P_{BAD}) and the expression of a cloned gene is induced by the addition of L-arabinose. *Pseudomonas* researchers have used the inducible properties of the *araC* regulator and the P_{BAD} promoter cassette for the controlled gene expression by integrating the *araC*-P_{BAD}-specific transcription fusion into the chromosome by using a suicide vector or an integration-proficient vector (1, 3, 13, 17, 30, 31). In the present study, we modified the existing *Escherichia-Pseudomonas* shuttle vectors pUCP20T, -26, -28T, and -30T by replacing the *lac* promoter with the *araC*-P_{BAD} cassette to allow conditional expression in pseudomonads and other bacteria, e.g., *Burkholderia* spp.

Construction and features of pHERD vectors. Functional genetic analysis requires vectors capable of conditional expression. The P_{BAD} promoter has been used for gene expression extensively in *E. coli* and some in *P. aeruginosa* and *Burkholderia* spp. (12, 27, 31). We first constructed three shuttle vectors, pHERD20T, -28T, and -30T (Fig. 1), based on *Escherichia-Pseudomonas* shuttle vectors pUCP20T, pUCP28T, and pUCP30T (29) and the commercial expression vector pBAD/Thio-TOPO (Invitrogen). The 368-bp fragment of the pUCP vectors spanning two restriction sites, AflII and EcoRI, was re-

placed with the *araC*-P_{BAD} fragment (1.3 kb), produced via PCR with pBAD/Thio-TOPO as the template and primers pBAD-F and pBAD-R (Table 1). The PCR product was purified and directly digested with AflII and EcoRI, and the two fragments were ligated into the pUCP vectors, creating pHERD20T (Fig. 1). The EcoRI/AflII regions of these vectors were sequenced to confirm that no mutations were introduced during the cloning pro-

FIG. 1. Construction of an *Escherichia-Pseudomonas* shuttle vector, pHERD20T, an arabinose-inducible expression vector. pHERD20T is a pUCP20T-based, conjugatable vector with pBR322- and pRO1600-derived replicons which support replication in *E. coli*, *P. aeruginosa*, and other bacteria, respectively. The P_{BAD} promoter was derived from the expression vector pBAD/Thio-TOPO (Invitrogen). The P_{lac} promoter in pUCP20T was replaced with the P_{BAD} promoter-containing segment with an EcoRI-AflII fragment generated via PCR containing the *araC* gene and P_{BAD}. Black arrows indicate the region transferred from pBAD/Thio-TOPO into pUCP20T. pHERD20T contains a multiple cloning site within *lacZα* encoding the β-galactosidase α peptide.

* Corresponding author. Mailing address: Robert C. Byrd Biotechnology Science Center, 1 John Marshall Drive, Huntington, WV 25755-9320. Phone: (304) 696-7356. Fax: (304) 696-7207. E-mail: yuh@marshall.edu.
[‡] D.Q. and F.H.D. contributed equally to this publication.
[▽] Published ahead of print on 10 October 2008.

TABLE 1. Bacterial strains and plasmids used in this study

Strain, plasmid, or primer	Genotype, phenotype, or sequence ^a	Source or reference
<i>P. aeruginosa</i>		
PAO1	Alg ^{wt} prototroph	P. Phibbs
PAO1VE2Δ <i>algW</i>	Alg ⁺ PAO1 <i>mucE</i> ^{+oe} (<i>himar1</i> Gm ^r :: <i>P</i> _{GM} :: <i>mucE</i>) Δ <i>algW</i>	20
<i>P. fluorescens</i> Pf-5	Alg ^{wt} prototroph	ATCC
<i>E. coli</i> DH5α	F [−] ϕ80 <i>dlacZ</i> Δ <i>M15</i> Δ(<i>lacZYA-argF</i>) <i>U169</i> <i>deoR</i> <i>recA1</i> <i>endA1</i> <i>hsdR17</i> (r _K [−] m _K ⁺) <i>phoA</i> <i>supE44</i> λ [−] <i>thi-1</i> <i>gyrA96</i> <i>relA1</i>	Laboratory strain
<i>B. pseudomallei</i> Bp50	Δ(<i>amrRAB-oprA</i>) derivative of wild-type strain 1026b	5
Plasmids		
pBAD/Thio-TOPO	<i>araC</i> -P _{BAD} <i>ori</i> _{pUC} Ap ^r (4,454 bp)	Invitrogen
pHERD20T	pUCP20T P _{lac} replaced with 1.3-kb AflII-EcoRI fragment of <i>araC</i> -P _{BAD} cassette (5,087 bp)	This study
pHERD30T	pUCP30T P _{lac} replaced with 1.3-kb AflII-EcoRI fragment of <i>araC</i> -P _{BAD} cassette (5,216 bp)	This study
pHERD26T	pUCP26 P _{lac} replaced with 2.4-kb AdhI-EcoRI fragment of <i>araC</i> -P _{BAD} cassette and <i>oriT</i> (6,166 bp)	This study
pHERD28T	pUCP28T P _{lac} replaced with 1.3-kb AflII-EcoRI fragment of the <i>araC</i> -P _{BAD} cassette (4,993 bp)	This study
pHERD30T- <i>mucE</i>	<i>mucE</i> in pHERD30T EcoRI/HindIII	This study
pHERD20T- <i>algU</i>	<i>algU</i> in pHERD20T EcoRI/HindIII	This study
pHERD20T- <i>oprF</i>	<i>oprF</i> (PA1777) in pHERD20T EcoRI/HindIII	This study
pHERD20T- <i>oprF</i> -WVF	<i>oprF</i> allele encoding OprF ending with the WVF motif cloned in pHERD20T EcoRI/HindIII	This study
Primers		
pBAD-F	AGTATACCTTAAGGAATCCCCAAATTATGACAACCTTGACGGCTACATCAT	This study
pBAD-R	AGGATCCCCGGGTACCGAGCTCGAATTCCTATCAGATCCCATGGGTATG TATA	This study
pHERD-SF	ATCGCAACTCTCTACTGTTTCT	This study
pHERD-SR	TGCAAGGCGATTAAGTTGGGT	This study
algU-F	AGAATTCGATGCTAACCCAGGAACAGGA	This study
algU-R	CAAGCTTTTCAGGCTTCTCGCAACAAAGGCTGCA	This study
algW-F	AGAATTCGATGCCCAAGGCCCTGCGTTTCCT	This study
algW-R	TGCCAAGCTTTCACTCGCCGCCGCTCTGTTT	This study
mucE-F	AGAATTCGATGGGTTTCCGGCCAGTTA	This study
mucE-R	GAAGCTTCAAAACACCCAGCGCAACTCGTC	This study
oprF-F	AGAATTCGATGAAACTGAAGAACACCTTA	This study
oprF-R	CAAGCTTTTACTTGGCTTCAGCTTCTACTTCGGCT	This study
oprF-WVF-R	AAGCTTAAACACCCAGCGCTTGGCTTCAGCTTCTACTTCGGCT	This study
lacZ-RT-For2	GTCGTGACTGGGAAAACC	This study
lacZ-RT-Rev2	GCCTCTCGCTATTACG	This study
Bp23S_F	GTAGACCCGAAACCAGGTGA	This study
Bp23S_R	CACCCCTATCCACAGCTCAT	This study

^a Alg^{wt}, wild-type nonmucoid phenotype; Alg[−], nonmucoid phenotype; Alg⁺, mucoid phenotype. Primers used for cloning carried built-in restriction sites (underlined), with F denoting forward and R denoting reverse primers, respectively.

cess. We next transferred the 2.4-kb AdhI-EcoRI fragment from pHERD20T to pUCP26, generating pHERD26T (Tet^r, 6,166 bp), which includes the *araC*-P_{BAD} cassette and the *oriT* sequence.

The pHERD vectors have the features of the pUCP vector family, including the pBR322 origin, four different antibiotic resistance markers, the *oriT* region for conjugation-mediated plasmid transfer (23), *ori*₁₆₀₀, and the *rep* gene encoding the replication-controlling protein (24, 29). However, the main advantage for cloning into the pHERD vectors is low expression, which occurs from the P_{BAD} promoter when it is not induced (Fig. 2). α complementation is inducible for blue-white screening, which facilitates the identification of recom-

binants on a 5-bromo-4-chloro-3-indolyl-β-D-galactopyranoside (X-Gal)-containing plate supplemented with arabinose (0.01%). The P_{BAD} promoter responds in a dose-dependent manner (Fig. 2). Two sequencing and PCR primers were designed that anneal to regions on both sides of the multiple cloning site, pHERD-SF 78 bp upstream of the EcoRI site and pHERD-SR 49 bp downstream of the HindIII site. If a gene is cloned in frame into the EcoRI site, a fusion protein with an additional seven NH₂-terminal amino acids (MGSDKNS) derived from thioredoxin of pBAD-TOPO/Thio will result. Thioredoxin acts as a translation leader to facilitate high-level expression and, in some cases, increase solubility in *E. coli* (9). These amino acids at the N terminus of the target protein may

FIG. 2. Arabinose-regulated *lacZα* expression in *B. pseudomallei*. RNA was extracted from log-phase *B. pseudomallei* Bp50 cells harboring pHERD30T that either had no arabinose added (None) or were induced for 2 h by the addition of the indicated amounts of L-arabinose. Quantitative real-time PCR was performed by using *lacZα*-specific primers. Data were normalized by using the 23S rRNA gene as the housekeeping control.

also serve as an epitope tag for protein analysis. pHERD vectors can be readily transferred from *E. coli* into *Pseudomonas* species and other bacteria via triparental conjugation (7) or by electroporation. It has been shown that the progenitor plasmid pRO1614 could replicate in a series of bacterial species, including *P. aeruginosa*, *P. putida*, *P. fluorescens*, *Klebsiella pneumoniae* (19), and *Burkholderia* spp. (5, 26). Therefore, the pHERD vectors are most likely functional in these bacteria. Another feature of the P_{BAD} promoter is catabolite repression of expression in the presence of glucose in the growth medium, which reduces intracellular cyclic AMP concentrations in *E. coli* cells, preventing the transcriptional activation of many genes by the cyclic AMP-binding protein (8).

Validation of pHERD20T in *P. aeruginosa* by modulating alginate production. We have observed that pHERD vectors can be used for the high-fidelity cloning and conditional expression of P_{BAD} transcription in the absence of L-arabinose (10). Initial attempts to clone the open reading frame of *P. aeruginosa* alternative sigma factor *algU* into pUCP20T were not successful. All of the *algU* alleles cloned were not functional, and sequence analysis showed that only mutant *algU* alleles were cloned into pUCP20T. This was consistent with the previous observations that *algU/T* cannot be cloned into the common expression vectors (16, 21). However, the *algU* gene was readily cloned into pHERD20T. Upon the expression of *algU* from P_{BAD} on pHERD20T, we observed dose-dependent alginate production or mucoidy in *P. aeruginosa* strain PAO1 in response to arabinose in the growth medium (Fig. 3).

Overexpression of the small peptide encoded by *mucE* activates AlgW, inducing alginate production (Fig. 4) in *P. aeruginosa* PAO1 and PA14 (20). Overexpression of *mucE* caused mucoidy in *P. aeruginosa* PAO1 and *P. fluorescens* Pf-5 (Table 2). The C-terminal WVF signal encoded by *mucE* is required for the activation of AlgW. The outer membrane protein OprF does not activate alginate production (Fig. 4); however, addition of the MucE WVF signal motif to the C terminus of OprF did cause alginate production (Table 2). Some genes are not highly expressed, and therefore expression in *trans* for complementation needs to be conditional. Expression of *algW* from P_{BAD} can complement an *algW* mutant back to alginate pro-

FIG. 3. Arabinose-dependent induction of alginate production in *P. aeruginosa* PAO1 carrying pHERD20T-*algU*. PAO1 with pHERD20T-*algU* was grown at 37°C for 24 h on *Pseudomonas* isolation agar and LB plates supplemented with carbenicillin and 0, 0.1, and 1.0% arabinose, respectively. The empty pHERD20T vector was used as the control (open box). Bars indicate means with standard errors. PAO1/pHERD20T with 0.1 and 1.0% arabinose does not increase alginate production (data not shown). OD, optical density.

duction due to titratable expression (Table 2). In addition to PAO1, we have used the pHERD vectors in PA14, CF149, environmental *P. fluorescens* isolates, and *P. putida* (data not shown). We have successfully employed pHERD30T for complementation of the $\Delta(amrAB-oprA)$ efflux pump mutation in *Burkholderia pseudomallei* strain Bp50 (5). In this case, however, complementation was also observed in uninduced cells, presumably because of basal transcription from the P_{BAD} promoter, which could not be overcome by growing cells in the

FIG. 4. Regulated alginate production in *P. aeruginosa*. Regulation of alginate production in *P. aeruginosa* involves many genes coding for products with many different functions. Mucoidy or alginate production is directed by the alternative σ^{22} factor AlgU (14). MucA is the cognate anti-sigma factor that negatively regulates AlgU activity by sequestering AlgU to the inner membrane (IM) (22). Sequestering of AlgU by MucA can be relieved by either mutation of *mucA* (15) or proteolytic degradation of MucA by the intramembrane protease AlgW (20). Derepression of MucA causes AlgU activation and alginate production. OM, outer membrane.

TABLE 2. Modulation of mucoidy in *P. aeruginosa* and *P. fluorescens* by pHERD20T-borne alginate regulators

Strain (genotype)	Colony morphology ^a	Plasmid	Colony morphology with plasmid ^a				
			1% Glucose	0% Arabinose	0.1% Arabinose	1% Arabinose	2.5% Arabinose
<i>P. aeruginosa</i>							
PAO1	NM	pHERD20T- <i>algU</i>	NM	NM	M	M	NM
		pHERD30T- <i>mucE</i>	NM	NM	M	M	M
		pHERD20T- <i>oprF</i>		NM	NM		
		pHERD20T- <i>oprF</i> -WVF		NM	M		
		pHERD20T- <i>algW</i>	NM	M	M	NM	NM
PAO1VE2Δ <i>algW</i> (P _{Gm} - <i>mucE</i> Δ <i>algW</i>)	NM						
<i>P. fluorescens</i> Pf-5	NM	pHERD30T- <i>mucE</i>		NM	M		

^a NM, nonmucoid; M, mucoid.

presence of 0.2% glucose and was not dependent on the growth medium used for the MIC assays (LB versus Mueller-Hinton broth) (data not shown).

In summary, we constructed a series of small *Escherichia-Pseudomonas* shuttle vectors with the *E. coli* *araC* and P_{BAD} promoter for highly regulated expression of cloned genes in *Pseudomonas* species and other bacteria and confirmed their utility by modulation of alginate production. Results presented here demonstrate that pHERD vectors are useful tools for bacterial physiological research and gene function studies with pseudomonads, as well as other bacteria, including medically significant *Burkholderia* spp.

Nucleotide sequence accession numbers. The GenBank accession numbers for the nucleotide sequences of pHERD20T, -26T, -28T, and -30T are EU603324, EU603327, EU603325, and EU603326, respectively.

The pHERD vectors described here are dedicated to the memory of the 1970 Marshall University Thundering Herd football team as depicted in the 2006 Warner Bros. film *We Are Marshall*.

This work was supported by a research grant (NNA04CC74G) from the National Aeronautics and Space Administration (NASA) and research grants from the NASA West Virginia Space Grant Consortium to H.D.Y. H.P.S.'s *Burkholderia* research was supported by NIH grant AI065357. F.H.D. was supported by a training grant (NNX06AH20H) from the NASA Graduate Student Researchers Program (GSRP).

We thank N. E. Head for the initial analysis of the mutant *algU* alleles in pUCP20T, V. M. Eisinger for the generation of the VE mutants, and K. D. Dillon for technical assistance with the alginate assay.

REFERENCES

- Baynham, P. J., D. M. Ramsey, B. V. Gvozdyev, E. M. Cordonnier, and D. J. Wozniak. 2006. The *Pseudomonas aeruginosa* ribbon-helix-helix DNA-binding protein AlgZ (AmrZ) controls twitching motility and biogenesis of type IV pili. *J. Bacteriol.* **188**:132–140.
- Ben-Samoun, K., G. Leblon, and O. Reyes. 1999. Positively regulated expression of the *Escherichia coli* *araBAD* promoter in *Corynebacterium glutamicum*. *FEMS Microbiol. Lett.* **174**:125–130.
- Boles, B. R., M. Thoendel, and P. K. Singh. 2005. Rhamnolipids mediate detachment of *Pseudomonas aeruginosa* from biofilms. *Mol. Microbiol.* **57**:1210–1223.
- Brosius, J., M. Erfle, and J. Storella. 1985. Spacing of the –10 and –35 regions in the *tac* promoter. Effect on its *in vivo* activity. *J. Biol. Chem.* **260**:3539–3541.
- Choi, K.-H., T. Mima, Y. Casart, D. Rholl, A. Kumar, I. R. Beacham, and H. P. Schweizer. 2008. Genetic tools for select agent compliant manipulation of *Burkholderia pseudomallei*. *Appl. Environ. Microbiol.* **74**:1064–1075.
- de Boer, H. A., L. J. Comstock, and M. Vasser. 1983. The *tac* promoter: a functional hybrid derived from the *trp* and *lac* promoters. *Proc. Natl. Acad. Sci. USA* **80**:21–25.
- Figurski, D. H., and D. R. Helinski. 1979. Replication of an origin-containing derivative of plasmid RK2 dependent on a plasmid function provided in trans. *Proc. Natl. Acad. Sci. USA* **76**:1648–1652.
- Guzman, L.-M., D. Belin, M. J. Carson, and J. Beckwith. 1995. Tight regulation, modulation, and high-level expression by vectors containing the arabinose P_{BAD} promoter. *J. Bacteriol.* **177**:4121–4130.
- LaVallie, E. R., E. A. DiBlasio, S. Kovacic, K. L. Grant, P. F. Schendel, and J. M. McCoy. 1993. A thioredoxin gene fusion expression system that circumvents inclusion body formation in the *E. coli* cytoplasm. *Bio/Technology* **11**:187–193.
- Lee, N., C. Francklyn, and E. P. Hamilton. 1987. Arabinose-induced binding of AraC protein to *araI2* activates the *araBAD* operon promoter. *Proc. Natl. Acad. Sci. USA* **84**:8814–8818.
- Lee, N., W. O. Gielow, and R. G. Wallace. 1981. Mechanism of *araC* autoregulation and the domains of two overlapping promoters, P_C and P_{BAD}, in the L-arabinose regulatory region of *Escherichia coli*. *Proc. Natl. Acad. Sci. USA* **78**:752–756.
- Lefebvre, M. D., and M. A. Valvano. 2002. Construction and evaluation of plasmid vectors optimized for constitutive and regulated gene expression in *Burkholderia cepacia* complex isolates. *Appl. Environ. Microbiol.* **68**:5956–5964.
- Ma, L., K. D. Jackson, R. M. Landry, M. R. Parsek, and D. J. Wozniak. 2006. Analysis of *Pseudomonas aeruginosa* conditional *psl* variants reveals roles for the *psl* polysaccharide in adhesion and maintaining biofilm structure postattachment. *J. Bacteriol.* **188**:8213–8221.
- Martin, D. W., B. W. Holloway, and V. Deretic. 1993. Characterization of a locus determining the mucoid status of *Pseudomonas aeruginosa*: AlgU shows sequence similarities with a *Bacillus* sigma factor. *J. Bacteriol.* **175**:1153–1164.
- Martin, D. W., M. J. Schurr, M. H. Mudd, J. R. Govan, B. W. Holloway, and V. Deretic. 1993. Mechanism of conversion to mucoidy in *Pseudomonas aeruginosa* infecting cystic fibrosis patients. *Proc. Natl. Acad. Sci. USA* **90**:8377–8381.
- Mathee, K., C. J. McPherson, and D. E. Ohman. 1997. Posttranslational control of the *algT* (*algU*)-encoded σ^{22} for expression of the alginate regulon in *Pseudomonas aeruginosa* and localization of its antagonist proteins MucA and MucB (AlgN). *J. Bacteriol.* **179**:3711–3720.
- Mdluli, K. E., P. R. Witte, T. Kline, A. W. Barb, A. L. Erwin, B. E. Mansfield, A. L. McClerren, M. C. Pirrung, L. N. Tumey, P. Warren, C. R. Raetz, and C. K. Stover. 2006. Molecular validation of LpxC as an antibacterial drug target in *Pseudomonas aeruginosa*. *Antimicrob. Agents Chemother.* **50**:2178–2184.
- Newman, J. R., and C. Fuqua. 1999. Broad-host-range expression vectors that carry the L-arabinose-inducible *Escherichia coli* *araBAD* promoter and the *araC* regulator. *Gene* **227**:197–203.
- Olsen, R. H., G. DeBusscher, and W. R. McCombie. 1982. Development of broad-host-range vectors and gene banks: self-cloning of the *Pseudomonas aeruginosa* PAO chromosome. *J. Bacteriol.* **150**:60–69.
- Qiu, D., V. M. Eisinger, D. W. Rowen, and H. D. Yu. 2007. Regulated proteolysis controls mucoid conversion in *Pseudomonas aeruginosa*. *Proc. Natl. Acad. Sci. USA* **104**:8107–8112.
- Schurr, M. J., D. W. Martin, M. H. Mudd, and V. Deretic. 1994. Gene cluster controlling conversion to alginate-overproducing phenotype in *Pseudomonas aeruginosa*: functional analysis in a heterologous host and role in the instability of mucoidy. *J. Bacteriol.* **176**:3375–3382.
- Schurr, M. J., H. Yu, J. M. Martinez-Salazar, J. C. Boucher, and V. Deretic. 1996. Control of AlgU, a member of the sigma E-like family of stress sigma factors, by the negative regulators MucA and MucB and *Pseudomonas aeruginosa* conversion to mucoidy in cystic fibrosis. *J. Bacteriol.* **178**:4997–5004.
- Schweizer, H. P. 1992. Allelic exchange in *Pseudomonas aeruginosa* using

- novel ColE1-type vectors and a family of cassettes containing a portable *oriT* and the counter-selectable *Bacillus subtilis* *sacB* marker. *Mol. Microbiol.* **6**:1195–1204.
24. **Schweizer, H. P.** 1991. *Escherichia-Pseudomonas* shuttle vectors derived from pUC18/19. *Gene* **97**:109–121.
 25. **Schweizer, H. P.** 2001. Vectors to express foreign genes and techniques to monitor gene expression in pseudomonads. *Curr. Opin. Biotechnol.* **12**:439–445.
 26. **Schweizer, H. P., T. R. Klassen, and T. Hoang.** 1996. Improved methods for gene analysis and expression in *Pseudomonas*, p. 229–237. *In* T. Nakazawa, K. Furukawa, D. Haas, and S. Silver (ed.), *Molecular biology of pseudomonads*. ASM Press, Washington, DC.
 27. **Suh, S. J., L. A. Silo-Suh, and D. E. Ohman.** 2004. Development of tools for the genetic manipulation of *Pseudomonas aeruginosa*. *J. Microbiol. Methods* **58**:203–212.
 28. **Tabor, S., and C. C. Richardson.** 1985. A bacteriophage T7 RNA polymerase/promoter system for controlled exclusive expression of specific genes. *Proc. Natl. Acad. Sci. USA* **82**:1074–1078.
 29. **West, S. E., H. P. Schweizer, C. Dall, A. K. Sample, and L. J. Runyen-Janecky.** 1994. Construction of improved *Escherichia-Pseudomonas* shuttle vectors derived from pUC18/19 and sequence of the region required for their replication in *Pseudomonas aeruginosa*. *Gene* **148**:81–86.
 30. **Woolwine, S. C., A. B. Sprinkle, and D. J. Wozniak.** 2001. Loss of *Pseudomonas aeruginosa* PhpA aminopeptidase activity results in increased *algD* transcription. *J. Bacteriol.* **183**:4674–4679.
 31. **Wyckoff, T. J., B. Thomas, D. J. Hassett, and D. J. Wozniak.** 2002. Static growth of mucoid *Pseudomonas aeruginosa* selects for nonmucoid variants that have acquired flagellum-dependent motility. *Microbiology* **148**:3423–3430.
 32. **Yanisch-Perron, C., J. Vieira, and J. Messing.** 1985. Improved M13 phage cloning vectors and host strains: nucleotide sequences of the M13mp18 and pUC19 vectors. *Gene* **33**:103–119.