Published in final edited form as:

Org Lett. 2005 June 23; 7(13): 2711–2713. doi:10.1021/ol050910r.

Practical and Highly Enantioselective Synthesis of β -Alkynyl- β -Amino Esters Through Ag-Catalyzed Asymmetric Mannich Reactions of Silylketene Acetals and Alkynyl Imines

Nathan S. Josephsohn, Emma L. Carswell, Marc L. Snapper, and Amir H. Hoveyda

Department of Chemistry, Merkert Chemistry Center, Boston College Chestnut Hill, Massachusetts 02467

Abstract

A readily available *iso*-leucine-based phosphine ligand is used to promote Ag-catalyzed Mannich reactions between silylketene acetals and various alkynyl imines. Reactions can be effected in the presence of 5 mol % catalyst, without the need for rigorous exclusion of air, and with commercially available solvents (without purification) to afford the desired β -alkynyl- β -amino esters in 84–94% ee and 61–91% isolated yield.

β -Amino acids and derivatives constitute an important class of organic molecules critical to the preparation of a variety of biologically significant molecules.¹ Accordingly, in recent years a number of efforts have focused on the development of efficient and stereoselective methods that allow access to such compounds.¹ In terms of related catalytic asymmetric methods, enantioselective hydrogenations of β -amino acrylates² and conjugate additions of N-based nucleophiles to unsaturated carbonyls³ have been introduced. Moreover, several investigations have been concerned with catalytic asymmetric additions of different enolate equivalents to imines (Mannich reactions).⁴

Although significant advances in the development of methods for the synthesis of β -amino esters have occurred, a number of challenges in connection with substrate generality persist. Regarding catalytic asymmetric Mannich reactions, one critical issue arises from the fact that nearly all the investigations to date have focused on transformations that deliver β -amino esters that bear a β -ester^{5,6} or a β -aryl group⁴ (reactions of imines derived from aromatic aldehydes). There are only a few cases where α,β -unsaturated imines are used, and methods that yield β -alkyl amino esters are relatively scarce, particularly those that deliver high enantioselectivities (>90% ee).^{6,7} Finally, to the best of our knowledge, none of the catalytic asymmetric methods disclosed so far deliver β -alkynyl- β -amino esters.⁸

amir.hoveyda@bc.edu.

Supporting Information Available. Experimental and analytical data for substrates and products. This material is available free of charge via the Internet at <http://pubs.acs.org>.

As summarized in Scheme 1, we recently disclosed an efficient Ag-catalyzed method for enantioselective addition of silyl enol ethers to a variety of imines to afford the desired β -amino ketones in high enantioselectivity (typically >90% ee).⁹ One notable attribute of the above protocol is that it is not limited to reactions of aryl imines; alkyl, alkenyl and alkynyl imines can be used as well.

Based on such recent advances, and considering the above mentioned methodological deficiencies, we set out to establish whether the catalytic system depicted in Scheme 1 can be used to effect catalytic asymmetric Mannich reactions between non-aryl imines and silyl ketene acetals.

Herein, we report a highly efficient Ag-catalyzed enantioselective method for the addition of silyl ketene acetals to a variety of alkynyl imines (eq 1). The catalytic asymmetric protocol delivers the desired β -alkynyl- β -amino esters in up to 94% ee; optically enriched amino esters are readily converted to a number of synthetically useful derivatives (e.g., β -alkyl- β -amino esters). The Ag-catalyzed transformation requires a chiral phosphine ligand (**1**) that can be easily prepared from commercially available materials, including an inexpensive amino acid (*iso*-Leu). Moreover, the enantioselective processes can be carried out in air without the need for purified solvent.

As the data summarized in Table 1 indicate, Ag-catalyzed addition of silyl ketene acetal **3** to a wide range of alkynyl imines (**2a–h**) proceeds smoothly at -60°C to afford β -amino esters **4a–h** in 61–91% isolated yield and 84–94% ee (>98% conv in all cases). It is noteworthy that the Ag-catalyzed protocol is equally efficient and enantioselective with alkynyl imines that bear silyl (entry 1), aryl (entries 2–4), alkyl (entries 5–6, 8) and alkenyl (entry 7) substituents.

Our investigation focused on transformations with phenyl ester **3** for several reasons: (a) Initial studies indicated that other ester derivatives (e.g., *t*-butyl esters) undergo efficient addition but in lower enantioselectivity (see Scheme 2 for an example). (b) Preparation of silylketene acetal **3** is typically more convenient and efficient than ketene acetals derived from alkyl esters (vs phenyl ester). (c) Phenyl esters can be more readily converted to a variety of other esters. (d) Preliminary screening indicated that catalytic asymmetric additions with *t*-butyl(dimethyl)silyl ketene acetals affords significantly lower enantioselectivities (<50% ee).

Next, we examined several key functionalizations that may be carried out with optically enriched β -alkynyl amino esters. Thus, as illustrated by the example in eq (2), Pd-catalyzed hydrogenation of **4d** (obtained in 92% ee) leads to the formation of β -amino ester **5** in 84% isolated yield after silica gel chromatography. It is important to note that, under the conditions outlined in Table 1, imines derived from aliphatic aldehydes do not provide the desired product in the presence of silyl ketene acetal **3** (<2% conv by 400 MHz ^1H NMR analysis).

As demonstrated by preparation of Boc-protected β -amino ester **6** from **4b** (68% isolated yield), shown in Scheme 2, removal of the N-activating group is effected oxidatively through a procedure developed in these laboratories,¹⁰ concomitant with installment of an amide unit. In a similar fashion, as also depicted in Scheme 2, oxidative removal of the anisidyl unit of **4b** (94% ee) in the presence of activated optically pure amino acid **7** leads to the formation of **8** in 73% yield and >98% ee after silica gel chromatography.

Synthesis of *t*-butyl ester **10**, which proceeds in 61% isolated yield to afford the desired product in >98% ee is noteworthy for two reasons. (a) It provides an illustration of the lower enantioselectivity of Ag-catalyzed reactions involving silyl ketene acetals of alkyl esters (72% ee vs 94% ee obtained with ketene acetal **3**). (b) The oxidative conditions used for the removal of the anisidyl unit are compatible with an acid-sensitive group such as a *t*-butyl ester. Optically enriched β -amino esters that contain a β -alkyne group can serve as versatile building blocks in the synthesis of biologically active molecules.¹¹

In summary, we have developed an efficient Ag-catalyzed method for enantioselective synthesis of β -alkynyl- β -amino acids, a class of compounds that cannot be accessed by other catalytic enantioselective Mannich protocols. The optically enriched (84–94% ee) products obtained can be functionalized in a number of ways to afford a range of useful β -amino esters. Hydrogenation of the alkyne group delivers difficult-to-access β -alkyl- β -amino esters. Removal of the N-activating group can be effected en route to optically enriched unprotected β -amino esters or, alternatively, after in situ coupling with activated esters, to a variety of amides.

Future studies will focus on the development of other catalytic asymmetric Mannich reactions and related applications to the synthesis of biologically significant molecules.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This paper is dedicated to Professor Richard R. Schrock on the occasion of his sixtieth birthday. Financial support was generously provided by the NIH (GM-57212). N. S. J. is grateful to Schering-Plough Research Institute for a graduate fellowship.

References

- For related reviews, see: a) Juaristi E. *Enantioselective Synthesis of β -Amino Acids* 1997 Wiley-VCH New York b) Ma J-A. *Angew. Chem., Int. Ed* 2003;42:4290–4299. c) Kleinmann F, Trost B, Fleming I. *Comprehensive Organic Synthesis* 1991 Pergamon Press New York Chapter 4.1 d) Cardillo G, Tomasini C. *Chem. Soc. Rev* 1996:117–128. e) Juaristi E, Lopez-Ruiz H. *Curr. Med. Chem* 1999;6:983–1004. [PubMed: 10519909] e) Beddow JE, Davies SG, Smith AD, Russell AJ. *Chem. Commun* 2004:2778–2779. and references cited therein
- For recent examples, see: a) Tang W, Zhang X. *Org. Lett* 2002;4:1594–161 [PubMed: 12423111] b) Zhou Y-G, Tang W, Wang W-B, Li W, Zhang X. *J. Am. Chem. Soc* 2002;124:4952–4953. [PubMed: 12423111]

- 11982347] c Tang W, Wang W, Chi Y, Zhang X. *Angew. Chem., Int. Ed* 2003;42:3509–3511. c Hsiao Y, Rivera NR, Rosner T, Krska SW, Njolito E, Wang F, Sun Y, Armstrong JD, Grabowski EJJ, Tillyer RD, Spindler F, Malan C. *J. Am. Chem. Soc* 2004;126:9918–9919. [PubMed: 15303855]
3. a Myers JK, Jacobsen EN. *J. Am. Chem. Soc* 1999;121:8959–8960. b Horstmann TE, Guerin DJ, Miller SJ. *Angew. Chem., Int. Ed* 2000;39:3635–3638. c Sibi MP, Asano Y. *J. Am. Chem. Soc* 2001;123:9708–9709. [PubMed: 11572706] d Guerin DJ, Miller SJ. *J. Am. Chem. Soc* 2002;124:2134–2136. [PubMed: 11878965] e Sibi MP, Prabakaran N, Ghorpade SG, Jasperse CP. *J. Am. Chem. Soc* 2003;125:11796–11797. [PubMed: 14505383]
4. For examples of catalytic asymmetric Mannich reactions with ester-derived nucleophiles and non-activated imines, see: a Isgitani H, Ueno M, Kobayashi S. *J. Am. Chem. Soc* 1997;119:7153–7154. b Xue S, Yu S, Deng Y, Wulff WD. *Angew. Chem., Int. Ed* 2001;40:2271–2274. c Wenzel AG, Jacobsen EN. *J. Am. Chem. Soc* 2002;124:12964–12965. [PubMed: 12405820] d Bernardi L, Gothelf AS, Hazell RG, Jorgensen KA. *J. Org. Chem* 2003;68:2583–2591. [PubMed: 12662026] e Akiyama T, Itoh J, Yokota K, Fuchibe K. *Angew. Chem., Int. Ed* 2004;43:1566–1568. f Cozzi PG, Rivalta E. *Angew. Chem., Int. Ed* 2005;44:ASAP
5. For representative examples of metal-catalyzed asymmetric Mannich reactions involving glyoxylate-derived imines, see: a Ferraris D, Young B, Dudding T, Lectka T. *J. Am. Chem. Soc* 1998;120:4548–4549. b Fujii A, Hagiwara E, Sodeoka M. *J. Am. Chem. Soc* 1999;121:5450–5458. c Trost BM, Terrell LR. *J. Am. Chem. Soc* 2003;125:338–339. [PubMed: 12517138] d Kobayashi S, Matsubara R, Nakamura Y, Kitagawa H, Sugiura M. *J. Am. Chem. Soc* 2003;125:2507–2515. [PubMed: 12603138] e Uraguchi D, Terada M. *J. Am. Chem. Soc* 2004;126:5356–5357. [PubMed: 15113196] f Ooi T, Kameda M, Fujii J, Maruoka K. *Org. Lett* 2004;6:2397–2399. [PubMed: 15228288]
6. For examples of Mannich reactions promoted by chiral organic catalysts, see: a List B, Pojarliev P, Biller WT, Martin HJ. *J. Am. Chem. Soc* 2002;124:827–833. [PubMed: 11817958] b Notz W, Tanaka F, Barba CF. *Acc. Chem. Res* 2004;37:580–591. [PubMed: 15311957] c Cordova A. *Chem. Eur. J* 2004;10:1987–1997.
7. a Kobayashi S, Kobayashi J, Ishiani H, Ueno M. *Chem. Eur. J* 2002;8:4185–4190. b Matsunaga S, Yoshida T, Morimoto H, Kumagai N, Shibasaki M. *J. Am. Chem. Soc* 2004;126:8777–8785. [PubMed: 15250731]
8. For an example of a non-catalytic enantioselective synthesis of a β -alkynyl- β -amino ester, see: Saito S, Hatanaka K, Yamamoto H. *Org. Lett* 2000;2:1891–1894. [PubMed: 10891184]
9. For other Ag-catalyzed asymmetric reactions promoted in the presence of phosphine 1, see: a Josephsohn NS, Snapper ML, Hoveyda AH. *J. Am. Chem. Soc* 2003;125:4018–4019. [PubMed: 12670201] b Josephsohn NS, Snapper ML, Hoveyda AH. *J. Am. Chem. Soc* 2004;126:3734–3735. [PubMed: 15038725]
10. a Porter JR, Traverse JF, Hoveyda AH, Snapper ML. *J. Am. Chem. Soc* 2001;123:10409–10410. [PubMed: 11604001] b Akullian LC, Snapper ML, Hoveyda AH. *Angew. Chem., Int. Ed* 2003;42:4244–4247. c Traverse JF, Hoveyda AH, Snapper ML. *Org. Lett* 2003;5:3273–3275. [PubMed: 12943405]
11. For example, see: a Zablocki JA, Rico JG, Garland RB, Rogers TE, Williams K, Schretzman LA, Rao SA, Bovy PR, Tjoeng FS, Lindmark RJ, Toth MV, Zupc ME, McMakins DE, Adams SP, Miyano M, Markos CS, Milton MN, Paulson S, Herin M, Jacmin P, Nicholson NS, Panzer-Knodle SG, Haas NF, Page JD, Szalony JA, Taite BB, Salyers AK, King LW, Campion JG, Feigen L. *J. Med. Chem* 1995;38:2378–2394. [PubMed: 7608903] b Cossy J, Schmitt A, Cinquin C, Buisson D, Belotti D. *J. Med. Chem. Lett* 1997;7:1699–1700.

Scheme 1. Ag-catalyzed enantioselective synthesis of β -amino ketones

Scheme 2.

One-pot removal of N-activating group and subsequent functionalization

Table 1Ag-catalyzed enantioselective Mannich reaction of silylketene acetal **3** to alkynyl imines **2a—h**

entry	R		yield (%) ^a	ee (%) ^b
1	SiMe ₃	a	84	92
2	C ₆ H ₅	b	91	94
3 ^c	<i>p</i> -OMeC ₆ H ₄	c	61	92
4 ^c	1-naphth	d	87	92
5	<i>n</i> -C ₆ H ₁₃	e	71	86
6		f	67	84
7		g	89	90
8		h	91	90

^aIsolated yields after silica gel chromatography; all reactions proceeded to >98% conv., based on 400 MHz ¹H NMR analysis^bDetermined by chiral HPLC analysis; see the Supporting Information for details.^cReaction times are 36 h.