Published in final edited form as:

Org Lett. 2007 November 8; 9(23): 4849–4852. doi:10.1021/ol702185y.

syn Additions to 4 α -Epoxy pyranosides: Synthesis of L-Idopyranosides

Gang Cheng, Renhua Fan, Jesús M. Hernández-Torres, Fabien P. Boulineau, and Alexander Wei

Department of Chemistry, Purdue University, 560 Oval Drive, West Lafayette, IN 47907-2084

Abstract

The *syn*-selective addition of organozinc compounds to 4 α -epoxy pyranosides (4 α -EPs), generated from methyl β -D-glucoside or aminoglucoside, provides an efficient route to pyranosides with α -L-*ido* configurations, including L-iduronic acid, neosamine B, and higher monosaccharide derivatives.

Unsaturated pyranosides such as glycals are used extensively in carbohydrate and natural products synthesis.¹ One widespread application involves the formation of epoxyglycals by stereoselective addition of dimethyldioxirane (DMDO), which can be employed as glycosyl donors in the synthesis of *O*- and *C*-glycosides.² 4-Deoxypentenoses (4-DPs) bear a strong structural homology to glycals, but retain a stereogenic anomeric center. The reactivity profile of 4-DPs is similar to that of glycals but is somewhat more selective with respect to stereocontrol. 4-DPs can be epoxidized by DMDO in a highly facioselective manner,^{3,4} and 4 β -epoxy pyranosides (4 β -EPs) can undergo *anti*-selective (S_N2) ring opening to generate novel pyranosides with an L-*altro* configuration.³ 4 α -epoxy pyranosides (4 α -EPs), which can also be obtained with high stereoselectivity,⁴ should be equally valuable as precursors to L-pyranosides. Here we demonstrate the utility of organozinc reagents for *syn*-selective ring opening of 4 α -EPs into pyranosides with an L-*ido* configuration.

The efficient formation of α -C-glycosides via *syn*-selective ring openings of α -epoxyglycals has been achieved with mild organometallic reagents having a significant Lewis acid character, namely those derived from aluminum,^{5,6} boron,⁵ titanium,⁷ zirconium,⁸ or zinc.⁹ *Syn* delivery of the nucleophile is assumed to proceed by coordination and simultaneous activation of the epoxide by the metal center. While many of these reagents are likely to be useful for *syn*-selective addition to 4-EPs,¹⁰ organozinc halides (RZnX) are most appealing with respect to atom efficiency and functional group compatibility, and can be generated by several different methods.^{9,11}

4 α -EPs **6** and **7** were derived in good yields from β -methyl-D-glucoside **1** and β -methyl-D-glucosaminoside **2** respectively, using an oxidation–decarboxylative elimination sequence.³ ¹⁰ Phthalimide **4** was transformed into azido derivative **5** by diazo transfer onto the intermediate free amine.¹² DMDO oxidations of 4-DPs **3–5** were carried out at –55 °C to

produce 4 α -EPs **6–8** with 10:1 α : β stereoselectivity and in quantitative yield (Scheme 1).⁴ The 4 α -EPs could be stored at $-20\text{ }^{\circ}\text{C}$ for several months without concern for decomposition.

The most reliable *syn* additions were produced by generating RZnX species via the transmetallation of organolithium or Grignard reagents (see Supporting Information). Systematic examination of reaction conditions revealed that the presence of excess ZnBr₂ significantly improved the efficiency of addition (Table 1); adducts were produced in high yields using 2 equiv of RZnBr, whereas stronger Lewis acids such as Zn(OTf)₂ or BF₃·Et₂O were often incompatible with the epoxide. Furthermore, allyltrimethylsilane and other electron-rich olefins tended to react poorly with 4 α -EPs in the presence of Lewis acids, in contrast to their efficacy as nucleophiles in C-glycoside synthesis. These studies suggest that ZnBr₂ promotes *syn* addition by forming an active complex with RZnBr (see below), rather than by direct activation of the epoxide. The activation of organometallic reagents by Lewis acids has been well documented over the years.¹³

The addition of RZnX species to 4 α -EPs **6**, **7** and **8** proceeded with high selectivity (*syn:anti* > 20:1) for a variety of *sp*²- and *sp*-carbon nucleophiles, producing the corresponding L-idopyranoside derivatives in good to high yields (Table 2). For example, 4 α -EP **6** was converted into L-idopyranoside **9a** in 84% isolated yield, accompanied by a small amount of *anti* adduct and a product derived from the minor 4 β -EP isomer (<5% combined yield).¹⁴ 4 α -EP **6** also reacted efficiently with activated *sp*³-carbon nucleophiles such as allylzinc bromide (entry 7) but was less receptive to additions with (*Z*)-alkenylzinc agents (entries 10, 13). These derivatives could be obtained with higher overall yields by partial hydrogenation of the corresponding alkynyl derivatives.

Several of the *syn* adducts in Table 2 were examined as intermediates for preparing L-idopyranosides with natural or unnatural configurations. In particular, we were interested to develop novel routes to L-iduronic acid, a vital component of heparin and heparan sulfate,¹⁵ neosamine B, a 2,6-diaminopyranoside with L-*ido* configuration found in several aminoglycoside antibiotics,¹⁶ and higher monosaccharides with structural analogy to sialic acid and other biologically active congeners.^{17,18} Syntheses of these sugars typically involve manipulation of acyclic intermediates, either for epimerization of the C5 stereocenter in the cases of L-iduronic acid and neosamine B,^{19,20} or for chain extension at the reducing end in the case of higher monosaccharides.^{17,21} In comparison, 4-EPs enable the stereoselective installation of carbon fragments at C5 with retention of the pyranose ring and anomeric configuration.

L-iduronic acid methyl glycoside **12** was prepared in 73% yield from 2-furyl adduct **9a** by ozonolysis with Me₂S workup (Scheme 2). Neosamine B methyl glycoside **14** was derived in 3 steps from **11e** (the C5 vinyl adduct of 4 α -EP derivative **8**) by ozonolysis, reductive amination, and hydrogenation, and fully characterized as peracetate **15**.

Reactions involving unactivated alkylzinc halides and dialkylzincs did not proceed as smoothly under comparable reaction conditions. For example, phenethylzinc bromide and diphenethylzinc gave low to fair yields of expected L-idopyranoside **16** accompanied by significant amounts of C5 phenethoxy adducts **17** (Scheme 3), an indication of the oxidative sensitivity of alkylzinc species.²² Rigorous removal of oxygen from the reaction mixture did not improve the ratio of C- to O-addition products, implying that alkylzincs could be oxidized by the 4-EPs or an unidentified byproduct of the DMDO reaction (Scheme 1).²³

L-Idopyranosides with C5 alkyl substituents can be prepared with considerable structural diversity and in synthetically useful yields via the *syn* addition of alkenyl groups. We illustrate this with the stereoselective syntheses of several octopyranoside derivatives, all with L-*ido* configuration but stereochemically divergent at C6 and C7. Compounds **20** and **21** were prepared respectively from *trans*-allylic ether **9h** and *cis*-allylic ether **18** (via reduction of propargylic ether **9i**) by catalytic osmylation, with diastereomeric ratios of 5:1 and 6:1 favoring the desired product (Scheme 4). The configurations of octopyranosides **20** and **21** were tentatively assigned as D-*threo*-L-*ido* and L-*erythro*-L-*ido* with the stereochemical relationship of C5 and C6 being *erythro* in each case, according to the empirical rule developed by Kishi and coworkers.²⁴ The stereochemical assignments were confirmed by nuclear Overhauser enhancement (NOE) measurements (Figure 1). The osmylation of α,β -unsaturated lactone **9m**, which was expected to proceed with selectivity opposite to that of **18**, yielded a single diastereomer which was assigned as D-*erythro*-L-*ido* by NOE analysis of diacetate **22**. Compound **9m** could also be oxidized stereoselectively into epoxide **19** by hypochlorite oxidation²⁵ and converted cleanly by phenylselenol reduction²⁶ into 7-deoxyoctopyranoside **23**, whose C6 configuration was assigned to be L-*glycero* by NOE measurements.

In conclusion, ZnBr₂-mediated *syn* additions to 4 α -epoxy pyranosides provides a general and efficient route to natural and unnatural L-idopyranosides.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

This work was supported by the National Institutes of Health (GM-06982). The authors also gratefully acknowledge support from the Purdue Cancer Center and the Purdue interdepartmental NMR facility, and Mr. Zhihong Huang (Purdue Univ.) for helpful discussions on organozinc chemistry.

References

1. Ferrier, R.J.; Hoberg, J.O. *Adv Carbohydr Chem Biochem*. Horton, D., editor. 58. Academic Press; New York: 2003. p. 55-119.
2. (a) Halcomb RL, Danishefsky SJ. *J Am Chem Soc* 1989;111:6661–6666. (b) Cheshev P, Marra A, Dondoni A. *Carbohydr Res* 2006;341:2714–2716. [PubMed: 17014831]
3. Boulineau FP, Wei A. *Org Lett* 2002;4:2281–2283. [PubMed: 12074687]
4. Cheng G, Boulineau FP, Liew ST, Shi Q, Wenthold PG, Wei A. *Org Lett* 2006;9:4545–4548. [PubMed: 16986946]
5. (a) Rainier JD, Cox JM. *Org Lett* 2000;2:2707–2709. [PubMed: 10990433] (b) Allwein SP, Cox JM, Howard BE, Johnson WB, Rainier JD. *Tetrahedron* 2002;58:1997–2009.
6. Bailey JM, Craig D, Gallagher PT. *Synlett* 1999:132–134.
7. Parrish JD, Little RD. *Org Lett* 2002;4:1439–1442. [PubMed: 11975598]
8. Wipf P, Pierce JG, Zhuang N. *Org Lett* 2005;7:483–485. [PubMed: 15673270]
9. (a) Leeuwenburgh MA, Timmers CM, van der Marel GA, van Boom JH, Mallet JM, Sinaÿ PG. *Tetrahedron Lett* 1997;38:6251–6254. (b) Steinhuebel DP, Fleming JJ, Du Bois J. *Org Lett* 2002;4:293–295. [PubMed: 11796073] (c) Xue S, Han KZ, He L, Guo QX. *Synlett* 2003:870–872.
10. Boulineau FP, Wei A. *J Org Chem* 2004;69:3391–3399. [PubMed: 15132547]
11. (a) Knochel P, Almena JJ, Jones P. *Tetrahedron* 1998;54:8275–8319. (b) Ikegami R, Koresawa A, Shibata T, Takagi K. *J Org Chem* 2003;68:2195–2199. [PubMed: 12636380] (c) Huo S. *Org Lett* 2003;5:423–425. [PubMed: 12583734] (d) Kneisel FF, Dochnahl M, Knochel P. *Angew Chem Int Ed* 2004;43:1017–1021. (e) Huang Z, Negishi E. *Org Lett* 2006;8:3675–3678. [PubMed: 16898789]
12. Alper PB, Hung SC, Wong CH. *Tetrahedron Lett* 1996;37:6029–6033.

13. (a) Negishi, E-i. *Pur Appl Chem* 1981;53:2333–2356. (b) Olah GA. *Angew Chem Int Ed* 1993;32:767–922. (c) Negishi, E-i. *Bull Chem Soc Japan* 2007;80:233–257.
14. Addition to the 4 β -EP isomer is presumed also to proceed with high *syn* selectivity.
15. Conrad, HE. *Heparin-Binding Proteins*. Academic Press; San Diego: 1998. (b) Linhardt RJ. *J Med Chem* 2003;46:2551–2564. [PubMed: 12801218]
16. Umezawa S. *Adv Carbohydr Chem Biochem* 1974;30:111–182. [PubMed: 4143427]
17. Danishefsky SJ, DeNinno MP. *Angew Chem Int Ed* 1987;26:15–23.
18. Unger, FM. *Adv Carbohydr Chem Biochem*. Horton, D., editor. 38. Academic Press; New York: 1982. p. 323–388. (b) Stenutz R, Weintraub A, Widmalm G. *FEMS Microbiol Rev* 2006;30:382–403. [PubMed: 16594963]
19. (a) Blanc-Muesser M, Defaye J. *Synthesis* 1977:568–569. (b) Jacquinet JC, Petitou M, Duchaussoy P, Lederman I, Choay J, Torri G, Sinaÿ P. *Carbohydr Res* 1984;130:221–241. (c) Ojeda R, de Paz JL, Martín-Lomas M, Lassaletta JM. *Synlett* 1999;8:1316–1318. (d) Takahashi H, Hitomi Y, Iwai Y, Ikegami S. *J Am Chem Soc* 2000;122:2995–3000.
20. Usui T, Takagi Y, Tsuchiya T, Umezawa S. *Carbohydr Res* 1984;130:165–177.
21. Györgydeák, Z.; Pelyvás, I. *Monosaccharide Sugars: Chemical synthesis by chain elongation, degradation, and epimerization*. Academic Press; San Diego: 1998. (b) Krülle T, Holst O, Brade H, Schmidt RR. *Carbohydr Res* 1993;247:145–158.
22. Katritzky AR, Luo Z. *Heterocycles* 2001;55:1467–1474.
23. In some instances the deoxygenated reactant (4-DP 3) could be recovered in low yield.
24. Cha JK, Christ WJ, Kishi Y. *Tetrahedron* 1984;40:2247–55.
25. Zhu L, Kedenburg PJ, Xian M, Wang GP. *Tetrahedron Lett* 2005;46:811–813.
26. Ding F, Jennings PM. *Org Lett* 2005;7:2321–2324. [PubMed: 15932188]

Figure 1. Stereochemical assignments of bicyclic lactones **20–23**. Conformations are drawn to emphasize NOE interactions.

Scheme 1. Synthesis of 4α-epoxy pyranosides ^a

^a Selected abbreviations: DMDO = dimethyldioxirane; DMFDNpA = dimethylformamide dineopentyl acetal; en = ethylenediamine; Phth = phthalimide; TEMPO = tetramethylpiperidine oxide; TfN₃ = triflyl azide.

Scheme 2.
Synthesis of L-iduronic acid and neosamine B methyl glycosides

Scheme 3.

Addition of mono- and dialkylzinc species to 4α-epoxypyranoside **6**

^a Epoxide **6** added as a CH₂Cl₂ solution at −78 °C to PhCH₂CH₂ZnBr (4 equiv) and ZnBr₂ (3 equiv) in THF, slowly warmed to −30 °C, then stirred for 12 h at −30 °C. ^b Epoxide **6** added at −78 °C to (PhCH₂CH₂)₂Zn (4 equiv) in THF, slowly warmed to −30 °C, then stirred for 4 h at 0 °C.

Scheme 4. Stereoselective synthesis of octopyranosides with *L-ido* configuration ^a

^a Selected abbreviations: BAIB = bisacetoxyiodobenzene; CSA = camphorsulfonic acid; MP = methoxypropene; NMO = morpholine *N*-oxide; TBS = *t*-butyldimethylsilyl; TEMPO = tetramethylpiperidinoxy radical.

Table 1

Selected conditions for the *syn* addition of 2-furylzinc halides to 4 α -epoxypyranoside **7** in the presence of ZnX₂^a

entry	furylZnX, equiv	ZnX ₂ (equiv)	yield ^b
1	2	ZnCl ₂ (2)	41%
2	2	ZnCl ₂ (5)	60%
3	4	ZnCl ₂ (2)	64%
4	10	ZnCl ₂ (2)	76%
5	2	ZnBr ₂ (0)	56%
6	2	ZnBr ₂ (2)	67%
7	2	ZnBr₂ (5)	83%

^aPreparation of 2-furylzinc halide in THF: (i) Furan, *n*-BuLi, THF, -78 °C, 30 min; (ii) ZnX₂, THF, 0 °C, 30 min. Epoxide **7** was added as a CH₂Cl₂ solution at -78 °C, warmed to 0 °C, then stirred for 2 h while warming to rt.

^bIsolated yields.

Table 2

Syn

additions to 4 α -epoxypyranosides **6–8**^a

6: X = OBn 7: X = NPhth 8: X = N ₃				
9: X = OBn 10: X = NPhth 11: X = N ₃				
entry	RZnX	prep method ^b	product	yield ^c
1		A	9a	84%
2		A	10a 9b	83% 74%
3			10b 9c	83% 69%
4		B	10c 9d	82% 80%
5		B	10d 9e	83% 69%
6		A	10e 11e 9f	74% 58% 77%
7		B	10f 9g	74% 73%
8		C	9h	60%
9		A	9i	79%
10		C	9j	37% ^d
11		C	9k	(77%) ^e 66%
12		C	9l	77%
13		D	9m	23% (73%) ^f

^a See Supporting Information for reaction conditions.^b A: (i) RH (4 eq), *n*-BuLi (2 equiv), THF, −78 °C; (ii) ZnBr₂ (7 equiv), THF, 0 °C. B: RMgBr (2 eq), ZnBr₂ (7 equiv), 0 °C. C: (i) RBr or RI (2 equiv), *t*-BuLi (4 equiv), Et₂O, −78 °C; (ii) ZnBr₂ (7 equiv). D: RI (2 equiv), Zn (activated powder), THF, 45 °C.

^cIsolated yields.

^d**9f** formed as a byproduct (40% yield).

^eYield in 2 steps via **9f**.

^fYield in 4 steps via **9i**.