Published in final edited form as:

Mol Pharm. 2007 ; 4(6): 873–882. doi:10.1021/mp0601190.

Disposition of Flavonoids via Enteric Recycling: Determination of the UDP-Glucuronosyltransferase (UGT) Isoforms Responsible for the Metabolism of Flavonoids in Intact Caco-2 TC7 Cells Using siRNA

Xing Liu¹, Vincent H. Tam², and Ming Hu^{1*}

¹*Department of Pharmacological and Pharmaceutical Sciences, College of Pharmacy, University of Houston, Houston, TX 77030*

²*Department of Clinical Sciences and Administration, College of Pharmacy, University of Houston, Houston, TX 77030*

Abstract

Our recent study indicates that microsomal glucuronidation rates are not predictive of the cellular glucuronide excretion rates and whole cell systems are needed to accurately determine the metabolic rates. This study aims to determine the contribution of UGT isoforms responsible for the metabolism of flavonoids in intact Caco-2 cells and cell lysates using siRNA. The results showed that UGT1A6 activities (as measured by p-nitrophenol glucuronidation) and expression were typically decreased 60–80% by siRNA treatment. Using siRNA-mediated silencing, we also showed that in intact cells, siRNA treatment substantially decreased the excretion of apigenin glucuronide at low and high concentrations (>35%, $p < 0.05$), although it only moderately decreased the excretion of genistein glucuronide at a high concentration (29%). The results also indicated that well expressed UGT isoforms in the Caco-2 cells, UGT1A1, UGT1A3, UGT1A6 and UGT2B7 were capable of metabolizing apigenin faster than genistein and that UGT1A6 silencing did not substantially increase the expression of genistein-metabolizing UGT isoforms. We also determined the contribution of UGT1A6 to the apigenin and genistein metabolisms as a function of concentration and the results indicated that metabolism of apigenin and genistein was saturable and siRNA treatment greatly reduced the rate of metabolism of apigenin but not that of genistein. In conclusion, we show for the first time that siRNA can be used effectively to determine which UGT isoform contributes to the metabolism of its substrate in intact cells. The results also indicate that UGT1A6 is a major contributor to glucuronidation of apigenin but not genistein in intact Caco-2 cells and in cell lysates.

INTRODUCTION

Apigenin (Fig.1) is a nonmutagenic bioflavonoid present in leafy plants and vegetables with significant chemopreventive activity against UV-radiation. Current literatures indicate that it may reduce DNA oxidative damage, inhibit the growth of human leukemia cells, induce cancer cells to differentiate, block growth signal transduction, and cause apoptosis in cancer cells ^{1–5}. However, the bioavailability of apigenin is poor ⁶, which may be a serious impediment to

¹Address correspondence to: Ming Hu, Ph.D., Department of Pharmacological and Pharmaceutical Sciences, College of Pharmacy, University of Houston, 1441 Moursund St. Houston, TX 77030, Tel: (713)-795-8320, Fax: (713)-795-8305, E-mail: mhu@uh.edu.

its clinical development since observed IC₅₀ or EC₅₀ values (in μM range) are significantly higher than its in vivo concentration (nM) achievable from dietary intake.

Genistein (Fig.1) is a constitutional isomer of apigenin, and it plays a major role as a weak estrogenic photochemical, and is a known estrogen modulator in animal models. It has anti-carcinogenic and anti-cancer properties against a variety of cancers including breast, colon, and prostate cancers⁷. Genistein exerts its pleiotropic effects on cancer cells by affecting cell survival and growth. However, genistein is also poorly bioavailable and most isoflavones in plasma are present as conjugated forms^{8, 9}. This is a serious concern because in vivo plasma concentrations of aglycones (unconjugated isoflavone) are in the range of 0.01 to 0.4 μM ^{8–10}, significantly less than the IC₅₀ or EC₅₀ values of 5 to 50 μM commonly reported for its in vitro anticancer effects^{6, 11}.

To address these concerns, our previous works had aimed at finding the causes of this bioavailability problem. Our studies have showed that the low bioavailability of apigenin and genistein was not the result of poor absorption but of extensive metabolism^{6, 11–15}. However, we have limited understanding of the enzyme isoforms responsible for the glucuronidation of apigenin and genistein and the contribution of each UGT isoform to their metabolism. Few published reports had systemically determined the isoforms responsible for flavonoid glucuronidation^{16, 17}, but none has been done using intact cells. In expressed isoform studies, it was found that 1A1, 1A4, 1A6, 1A7, 1A9, and 1A10 were capable of metabolizing genistein¹⁶. Limited studies of the UDP-glucuronosyltransferase (UGTs) or UGT isoforms responsible for apigenin metabolism suggested that it was metabolized rapidly in HepG2 cells via UDP-glucuronosyltransferase 1A1 (UGT1A1)¹⁸. However, UGT1A1 was only moderately expressed in the Caco-2 TC7 cells, whereas UGT1A9 was poorly expressed and UGT1A7 and UGT1A10 were not expressed in the Caco-2 TC7 cells¹⁹. Furthermore, investigators have not actually determined the contribution of a specific UGT isoform in an intact cell system. Study using intact cell system is highly valuable since our recent study indicates that microsomal derived kinetic parameters were not predictive of the intestinal or biliary excretions of glucuronides in rats²⁰. Therefore, the purpose of this study is to determine the contribution of relevant UGT isoforms in the metabolism of apigenin and genistein Caco-2 TC7 cells, and to develop an alternative approach to elucidate the contribution of a specific UGT isoform in intact cells.

MATERIALS AND METHODS

Materials

Cloned Caco-2 TC7 cells were a kind gift from Dr. Monique Rousset of INSERM U178 (Villejuif, France). Genistein and apigenin were purchased from Indofine Chemicals (Somerville, NJ). P-Nitrophenol and p-nitrophenol-glucuronide (PNP-g) were purchased from Sigma-Aldrich (St Louis, MO). A siRNA mixture or SMARTpool against UGT1A6 (Cat# M-020196-00-0050), 4 individual duplex siRNA, and the negative control-pool (Cat# D-001206-13-20) were products of DHARMACON Inc. (Dallas, TX). Lipofectamine 2000 and NOVEX pre-cast gels were from Invitrogen (CA). Expressed human UGTs 1A1, 1A3, 1A4, 1A6, 2B7, 2B15, 2B17 were purchased from BD Biosciences (Woburn, MA). All other materials (typically analytical grade or better) were used as received.

Cell Culture

The culture conditions for growing Caco-2 cells have been described previously^{15, 21, 22}. In the current experiments, we used cells that were grown for a short period of time using the same growth media (Dulbecco's modified Eagle's medium supplemented with 10% fetal bovine serum). Caco-2 TC7 cells were fed every other day.

siRNA Design

Based on the rules suggested by Elbashir et al.²³, the antisense strand of siRNA was targeted against anAA(N)₁₉ sequence at least 100 nucleotides downstream of the start codon. The GC content of the duplexes was kept within the 40–70% range. In addition to these rules, a BLAST search is conducted against GenBank to minimize the possibility of silencing another gene. A *SMART*pool that contained an equal molar mixture of four short siRNAs was generated by Dharmacon. The four pairs of siRNA have the sequences of:

GUACAGGAAUAACAUGAUUU (sense, pair 1)
 5'-P-AAUCAUGUUAUCCUG UACUU (antisense, pair 1),
 GAUAUGACUUUGUGCUUGAUU (sense, pair 2)
 5'-P-UCAAGCACAAAGUCAUAUCUU (antisense, pair 2),
 GAUCCUGGCUGAGUAUUUGUU (sense, pair 3)
 5'-P'-CAAAUACUCAGCCAGGAUCUU (antisense, pair 3)
 GAACCGUUACCAAUCAUUUUU (sense, pair 4)
 5'-P-AAAUGAUUGGUAACGGUUCUU (antisense, pair 4).

Transient transfection

Caco-2 TC7 cells were seeded at 5×10^4 cells/cm², and maintained in Dulbecco's modified Eagles medium (DMEM) containing 10% v/v fetal bovine serum (FBS, Hyclone), but without antibiotics. Twelve hours post-plating, the cells were doubly transfected (6 hours each) with serum-free media containing Lipofectamine along with either 50 ng total siRNA mixture or *SMART*pool, equal molar of an individual siRNA duplex, or the negative control-pool, whereas the control cells were treated with normal growth medium with Lipofectamine. At 48 hours post transfection, cells were harvested and cellular lysates were made and used for analyzing UGT1A6 enzyme activities in vitro and for blotting UGT1A6 protein levels as described later.

Caco-2 Cell Lysate

Caco-2 cell monolayers (3.3 cm² each) were harvested and put in 0.5 ml of 50 mM potassium phosphate (pH 7.4 buffer), and sonicated in Aquasonic 150D sonicator (VWR Scientific, Bristol, CT) for 30 min at the maximum power (135 average watts) in an ice-cold water bath. The resulting cell lysate was then used in the UGT activity assay.

Lysate Protein Concentration

Protein concentration of cell lysate was determined using the Bio-Rad protein assay kit (Hercules, CA), using bovine serum albumin as standards.

Measurement of UGT Activities Using Cell Lysate

The incubation procedures for measuring UDP-glucuronosyltransferase (UGTs) activities using cell lysate are as follows: 1) mix 114.3 μ l of cell lysate (final concentration ~0.5 mg of protein/ml), magnesium chloride (0.88 mM), saccharolactone (4.4 mM), alamethicin (0.022 mg/ml); different concentrations of substrates in a 50 mM potassium phosphate buffer (pH 7.4); and uridine diphosphoglucuronic acid (UDPGA) (3.5 mM, add last) to a final volume of 170 μ l; 2) incubate the mixture at 37°C for 45, 90min and 480 min for p-nitrophenol, apigenin and genistein, respectively; and 3) stop the reaction by the addition of 50 μ l of 94% acetonitrile/6% glacial acetic acid containing 100 μ M testosterone as an internal standard. In determining the kinetic parameters, apigenin concentrations were varied from 0.5 μ M to 100 μ M, and genistein from 0.5 μ M to 200 μ M.

Following this reaction scheme, the reaction progressed linearly for as long as the substrate is less than 50% exhausted or up to 12 hours, whichever is shorter. When measuring glucuronidation at different concentrations, we followed the time and % substrate exhaustion limit so we can accurately determine the rate of glucuronidation.

Measurement of UGT Activities Using Expressed Human UGTs

The incubation procedures for measuring UDP-glucuronosyltransferase (UGTs) activities using expressed Supersomes are similar to what was used for measuring UGT activities in cell lysate except the concentration of Supersomes in the final reaction mixture was about 0.026 mg/ml (for apigenin) to 0.052 mg/ml (for genistein) since the microsomes have much higher concentration than the cell lysate. The reaction time was 30 min for apigenin and 60 min for genistein. Under these reaction time and microsomal protein concentration, the maximal % substrate loss was 21% for apigenin and 26% for genistein respectively, well within the linear range of no more than 50% substrate loss.

Screening UGT Supersomes for Apigenin, Genistein and p-Nitrophenol Glucuronidation Activity

Glucuronidation activity by expressed UGT1A1, 1A3, 1A4, 1A6, 2B7, 2B15 and 2B17 were measured using the same assay conditions as described in “Measurement of UGT Activities Using Cell Lysate” except that the final protein concentration was 0.05 mg/ml.

Western Blot Analyses of UGT1A6 Protein Levels

Caco-2 cell lysate (12 µg protein) was separated on an SDS-PAGE (12% Tris-Glycine Gel), and then the separated proteins were transferred onto a nitrocellulose membrane by a standard protocol. After blocking for nonspecific binding using Tris buffer solution containing 5% nonfat milk and 1% Tween 80, the nitrocellulose membrane was incubated with WB-UGT1A6 (BD Biosciences, Wuburn, MA) antibody at 1:2000 dilution for 1 hr (at room temperature). After washing three times to remove nonspecifically bound antibody, the membrane was incubated with a horse radish peroxidase-conjugated secondary antibody followed by detection with enhanced chemiluminescence using SuperSignal West Femto reagents from Pierce (Rockford, IL). Western blotting of glyceraldehyde-3-phosphate dehydrogenase (GAPDH) was used as the loading control. Densities of the western blot bands were quantified using the density measurement software supplied by FluoroChem.

Excretion of Apigenin and Genistein Glucuronides by Intact Caco-2 Cells

The Caco-2 cell monolayers were washed three times with 37°C, pH 7.4 Hank's balanced salt solution (HBSS). The monolayers were incubated with the buffer for 1 h and the incubation medium was then aspirated. Afterward, the solution containing the compound of interest was loaded and samples (400 µl) were taken at the different incubation time, followed by the addition of 400 µl fresh buffer to keep the constant volume. To each sample, 50 µl of 6% glacial acetic acid in acetonitrile solution containing 100 µM testosterone was added as the internal standard. Afterward, the mixture was centrifuged at 13,000 rpm for 15 min, and the supernatant was analyzed by HPLC (Table 1).

Sample Analysis

The conditions for HPLC analysis of isoflavones were modified based on a previously published method (Chen et al., 2003). The HPLC conditions for analyzing p-nitrophenol, apigenin, genistein (see Fig.1 for structures), and their conjugates are as follows: system, Hewlett Packard 1090 with diarray detector and HP Chemstation (Wilmington, DE); column, Synergi Polar-RP (Phenomenex, Torrence, CA), 4 µm, 150 × 4.60 mm; mobile phase A (MBA), 100% acetonitrile; mobile phase B (MPB), 0.04% (w/v) phosphoric acid plus 0.06% (v/v)

triethylamine (pH 2.8). For p-nitrophenol and its glucuronide, the gradient is: 0 to 2 min, 2% MPA; 2 to 15 min 2%–20% MPA; 15 to 20 min 20%–33% MPA; 20 to 21 min 45.5% MPA. For apigenin and its glucuronide, the gradient is: 1 to 3 min 20% MPA; 3 to 23 min 20–48.5% MPA. For genistein and its glucuronide, the gradient is: 0 to 3 min 2% MPA; 3 to 35 min 2–45.5% MPA. Each injection is followed by a 5 min equilibrium time before the next injection. The retention times of P-Nitrophenol, apigenin, genistein and their conjugates are listed in Table 1. The limit of detection of this method is 0.25 μ M or 50pmol in the 200 μ l sample, the average accuracy was 92.7% for genistein and 91.1% for apigenin standards (8 total), using standard curves forced through the origin. Typical CV% is less than 5% between injections. Representative HPLC profiles of p-nitrophenol, genistein, and apigenin as well as their glucuronides are shown in Fig. 2.

Quantitation of Glucuronides

We have previously described how to calculate the concentration of metabolites¹⁵. Briefly, the metabolites are obtained from rat intestinal perfusate, which contains both glucuronides and parent flavonoid aglycones. After the perfusate is extracted with methylene chloride, the resulting aqueous phase contains only glucuronide. We then make serial dilution of phase II conjugate metabolites. Each solution of metabolites is divided into two halves. In one half, we add glucuronidase in water and the other water only. We allow the mixture to incubate in water bath for about 4–8 hrs (time varied) to allow the sample with glucuronidase to hydrolyze completely. We then measure the peak areas of original glucuronides and that of the released aglycones. We found a linear correlation between the released aglycone and the glucuronides and the correlation coefficient is then used to normalize the metabolite peak we measured in the experiments. In this way, we can quantify the metabolites using standard curve for the parent compound as was done previously¹⁵.

Data Analysis

The apparent kinetic parameters of K_m and V_{max} were estimated. For apigenin, as suggested by the Eadie-Hofstee analysis, 3 saturable (Michaelis-Menten based) models with and without autoactivation were fit to the observations between the rate of reaction and substrate concentrations, using the ADAPT II program (University of Southern California). The general model structure is shown in equation (1). To determine the best-fit model, the model candidates were discriminated using the Akaike's information criterion (AIC)²⁴, and the rule of parsimony was applied.

$$\text{Reaction rate} = \frac{[V_{\max-0} + V_{\max-d}(1 - e^{-CR})] \times C}{K_m + C} \quad (1)$$

where

$V_{\max-0}$	- intrinsic maximum enzyme reaction rate
$V_{\max-d}$	- maximum induction of enzyme activity
R	- rate of enzyme activity induction
C	- concentration of substrate
K_m	- concentration of substrate to achieve 50% of ($V_{\max-0} + V_{\max-d}$)

For genistein, the reaction kinetic data showed biphasic kinetics (in which two isoforms with different kinetic behaviors were responsible for the glucuronidation). Therefore, formation rates (V) of isoflavone glucuronides at various substrate concentrations (C) were fitted to the following equation: (2)

$$\text{Reaction rate} = \frac{V_{\max 1} \times C}{K_{m1} + C} + \frac{V_{\max 2} \times C}{K_{m2} + C} \quad (2)$$

where

$V_{\max 1}$	- maximum enzyme reaction rate of one UGT isoform
$V_{\max 2}$	- maximum enzyme reaction rate of another UGT isoform
K_{m1}	- concentration of substrate to achieve 50% of $V_{\max 1}$
K_{m2}	- concentration of substrate to achieve 50% of $V_{\max 2}$

Statistical Analysis

One-way ANOVA or an unpaired Student's *t* test (Microsoft Excel) was used to analyze the data. The prior level of significance was set at 5%, or $p < 0.05$.

RESULTS

Inhibition of UGT1A6 Expression in Caco-2 Cells using siRNA

Based on the expression pattern of UGT1As in the Caco-2 cells and activities of various UGT1As reported in the literature¹⁹, we expect UGT1A6 to be an important isoform for flavonoid glucuronidation. The results showed that UGT1A6 activities, as measured by p-nitrophenol (a prototypic UGT1A6 substrate) glucuronidation rates, decreased significantly in siRNA-treated Caco-2 cells ($p < 0.05$) (Fig.3A). This decrease in activities was corroborated by a decrease in UGT1A6 protein level in a Western blots, as signified by a 59% decrease in the band density ratio (UGT1A6/GAPDH) from 0.062 to 0.026 (Fig.3B). The same experiment was replicated several times, and UGT1A6 protein levels typically decreased by 60 to 80%. Furthermore, when the 4 siRNA duplexes were tested individually at the same molar concentration, results indicated that UGT1A6 activities and protein levels were decreased by a similar extent, which was also reflected by a maximum of 45% in the band density ratio (UGT1A6/GAPDH) from 0.097 to 0.053 (Fig.4 A, B). Based on these results, and the fact that a mixture of siRNA was easier to use, and provided more effective and consistent gene suppression, the rest of the experiments were performed with the siRNA mixture or simply siRNA.

Effect of siRNA on the Cellular Excretion of Apigenin and Genistein Glucuronides

To determine the contribution of each UGT isoform to apigenin glucuronide excretion in intact Caco-2 cells, we used siRNA since isoform specific chemical inhibitors were not available and UGT reaction rates were always the rate-controlling step in apigenin glucuronide excretion¹⁴. The results indicated that siRNA treatment in intact Caco-2 cells affected apical excretion of glucuronides in intact Caco-2 cells. At a low loading concentration of 5 μ M, siRNA treatment significantly decreased ($p < 0.05$) excretion of apigenin glucuronide from 0.0386 nmol/min/mg protein to 0.0286 nmol/min/mg protein (or 35%). On the other hand, the same siRNA treatment did not affect excretion of genistein glucuronide (0.0038 vs. 0.00348 nmol/min/mg protein or 8%) (Fig.5). In contrast, siRNA treatment appeared to have a more substantial effect on the excretion of both flavonoid glucuronides at a higher concentration (20 μ M). The excretion rates decreased from 0.149 to 0.085 nmol/min/mg protein for apigenin glucuronide (79% decrease, $p < 0.05$) and from 0.00801 to 0.00621 nmol/min/mg protein for genistein glucuronide (29% decrease, $p < 0.05$), respectively (Fig.5).

Glucuronidation Activities by Human UGT Isoforms towards Apigenin, Genistein and p-Nitrophenol

We then determined which UGT isoforms may contribute to the metabolism of apigenin and genistein by measuring the glucuronidation activities of expressed human UGTs 1A1, 1A3, 1A4, 1A6, 2B4, 2B7, 2B15, and 2B17. These four UGT1As were selected because they are well expressed by Caco-2 TC7 cells whereas other UGT1As such as UGT1A7, UGT1A8,

UGT1A9 and UGT1A10 were either poorly expressed (UGT1A8 and UGT1A9) or not expressed (UGT1A7 and UGT1A10) (Jeong et al, 2005). UGT2Bs were chosen since their expression levels in Caco-2 TC7 cells were unknown. The results indicated that human UGTs 1A1, 1A3, 1A6, 2B7, 2B15 glucuronidated apigenin at different rates (Fig. 6A) with UGT1A1 having the highest activity (set as 100%), followed by UGT1A6 (60% of UGT1A1), UGT1A3 (55%), UGT2B7 (15%), UGT2B15 (9%). Whereas, these same UGT isoforms also glucuronidated genistein, but at a slower rates than apigenin. Once again, UGT1A1 glucuronidated the fastest followed by UGT1A3 (35% of UGT1A1), UGT2B7 (16%), UGT1A6 (15%), UGT2B17 (4%) (Fig. 6B). In addition, we found that p-nitrophenol (at low concentration of 50 μ M) is a nearly specific substrate for UGT1A6 isoform (Fig. 6C), since only UGT2B7 had a minor contribution. This result is consistent with a previous finding that showed p-nitrophenol as a specific UGT1A6 substrate²⁵.

Effect of UGT1A6 Silencing on the Metabolism of Apigenin and Genistein by Caco-2 Cell Lysate

To determine the effects of UGT1A6 silence (by siRNA) on the metabolism of apigenin and genistein, the conjugation activities were measured at low, medium and high concentrations in Caco-2 cell lysates with or without siRNA treatment. The results indicated that in siRNA treated cells, glucuronidation rates of apigenin decreased significantly at all tested concentrations (57%, $p < 0.05$), whereas that of genistein was only modestly decreased (32%, $p < 0.05$) at a high concentration (50 μ M) (Table 2).

Kinetic Analysis of Apigenin and Genistein Glucuronidation by Caco-2 Cell Lysates

We determined the glucuronidation rates as a function of concentration because changes in rates of flavonoid glucuronidation as the result of siRNA treatment are dependent on the concentration of flavonoids (Table 2). Results indicated that the rates of glucuronidation increased with concentration and reached a plateau as the concentration of apigenin increased (Fig. 7A). Eadie-Hofstee plot of the same data (a hooked curve) was consistent with autoactivation kinetics²⁶ (Fig. 7B). Therefore, the data were fit using an atypical Michaelis-Menten model with autoactivation, and the best-fit model was found to be an autoactivation model with three parameters (Table 3). The atypical K_m values were 13.14 μ M for control and 5.32 μ M for siRNA treated cells, respectively. The V_{max} values were determined to be 0.574 nmol/min/mg for control, and 0.191 nmol/min/mg for siRNA treated cells, respectively.

Genistein glucuronidation as a function of concentration was also determined in Caco-2 cells without or with UGT1A6 gene silencing (Fig. 8). The Eadie-Hofstee plot indicated that the reaction kinetics followed a biphasic pattern regardless of siRNA treatment (Fig. 8B). Furthermore, the plot showed very sharp angles, suggesting two isoforms with significantly different K_m and/or V_{max} values. We first fitted these data using equation (2) with the ADAPTI software, but the fitting did not consistently generate meaningful kinetic parameters, most likely due to technical issues associated with model identifiability. Since there were minor differences in glucuronidation of genistein in siRNA treated cells and in control cells, we did not obtain the kinetic parameters to describe genistein glucuronidation.

DISCUSSION

The present study is a continuation of our effort to dissect the mechanisms involved in the intestinal disposition of flavonoids via the enteric recycling mechanism. It seeks to use siRNA to determine the contribution of a single UGT isoform in the metabolism of a flavonoid because siRNA appeared to be one of the few viable approaches to complete this task. The results of this study clearly indicate that siRNA is effective in defining the contribution of a particular UGT isoform inside intact cells and in cell lysates. The results of cell lysate studies in siRNA

treated cells are consistent with the fact that multiple human UGTs in both UGT1A and UGT2B subfamilies are involved in the metabolism of apigenin and genistein in the Caco-2 cells, and in the human intestinal and liver cells as well. Based on the fact that UGT1A6 is well expressed in Caco-2 cells but not particularly active against genistein, it was not surprising to see that siRNA-mediated UGT1A6 silencing in the Caco-2 cells decreased glucuronidation of apigenin but not genistein. It was somewhat unexpected, however, to observe that a 60–80% decrease in cellular UGT1A6 activity resulted in a similar decrease in cellular excretion of apigenin conjugates, since we have expected a more moderate effect as an earlier study using mature Caco-2 cells showed that efflux transporter was the rate-limiting step¹⁴.

It is difficult to separate the contribution of a single UGT isoform in the metabolism of a UGT substrate for the following reasons. First of all, UGT substrates such as flavonoids and raloxifene are often metabolized by multiple UGT isoforms (Fig.6)^{16–18}. Second, expression patterns of UGT isoforms are organ and tissue specific, and have high inter-subject variability^{27, 28}. Third, there is a lack of isoform specific substrate, or UGT isoform-specific chemical inhibitor or inhibitory antibody. Therefore, alternative approach is needed to define the contribution of a single UGT isoform to the metabolism of a substrate in functional cells.

This is the first report that details the use of siRNA as a plausible and effective tool to determine the contribution of a UGT isoform to the cellular metabolism of its substrate, which is a flavonoid in this case. Whereas the use of siRNA is not without its challenge (e.g., possible compensatorily increased expression of alternative UGT isoforms), the results suggest that expressions of several important UGT isoforms (UGT1A1, UGT1A5 and UGT2B7) did not increase substantially to compensate for UGT1A6 silencing in our study. This is because genistein metabolism rates at higher concentration would have increased if expression levels of highly active UGT1A1, UGT1A5 and UGT2B7 (Fig. 6B) were increased substantially to compensate for the of UGT1A6 silencing. Therefore, our data suggest that siRNA treatment has sufficient selectivity and did not cause substantial compensatory increases in the expressions of other (at least four) UGT isoforms in the Caco-2 cells.

It is fairly remarkable that siRNA-mediated UGT1A6 silencing decreased apigenin glucuronidation rates without affecting genistein glucuronidation in Caco-2 cell lysates (Table 2) since these two compounds are constitutional isomers that have very similar structures (Fig. 1). Based on the PCR expression factor of 2.6 UGT1A6, 1.3 UGT1A3 and 1 UGT1A1¹⁹, we had predicted that a 50% decrease in apigenin glucuronidation in cell lysates was likely (Fig. 7) since UGT1A1 was 40% more effective and UGT1A3 were equally effective as UGT1A6 in the glucuronidation of apigenin (Fig.6). The results showed a 60% decrease in metabolism rates by siRNA at high apigenin concentration (Fig. 7A), slightly exceeding our expectation. In addition, we expect less than 25% difference in genistein metabolism in cell lysates as the result of UGT1A6 silence, and the results (32% decrease) were mostly consistent with our expectation (Fig.8A). Taken together, these results suggest that siRNA is an effective method to determine the contribution of a dominant UGT isoform in the glucuronidation of drugs and chemicals in cell lysates.

We also expect siRNA silencing to significantly impact the kinetics of apigenin reaction but not that of genistein in the cell lysates based on the fact the UGT1A6 is a major isoform for metabolism of apigenin but not of genistein (Fig.6). As expected, there was a large difference in the rates of metabolism as a consequence of siRNA silencing (Fig.7A, Table 3). V_{max} values of apigenin glucuronidation decreased 67% as the result of siRNA treatment (Table 3). This decrease was comparable to a maximum reduction of 60–80% in silencing efficiency as measured by Western blots and p-nitrophenol glucuronidation (Fig.3 and Fig.4). In contrast, siRNA silencing did not significantly affect the rates of genistein glucuronidation at the tested concentrations (Fig.8A).

Whereas we have expected that siRNA treatment will be highly useful for the determination of the contribution of UGT isoform to the metabolism of its substrate in cell lysates, we could not predict what might happen to cellular excretion of phase II conjugates in intact Caco-2 cells. This is because our studies in Caco-2 cells suggest that cellular UGT activities are often not the rate-limiting step in the cellular excretion of phase II conjugates^{12, 14}. Moreover, recent studies in the rat models clearly indicate that microsomal (and therefore other subcellular preparations) glucuronidation rates are not predictive of cellular glucuronide excretion rates²⁰. Therefore, we were not surprised by the fact that siRNA silencing had a smaller effect on the cellular excretion of glucuronides of apigenin (35%) and genistein (8%), which were obtained using a flavonoid concentration of 5 μ M (Fig.5). We were somewhat surprised that the cellular excretion of glucuronides was more affected at a higher flavonoid concentration of 20 μ M since excretion of glucuronides of apigenin and genistein decreased 79% and 29%, respectively (Fig.5). These decreases in excretion were similar to or exceeded decreases in cellular lysate metabolism, which were not expected. We attributed this large than expected decreases in cellular excretion to inhibition of BCRP by high concentrations of flavonoids^{29, 30} since BCRP has been shown recently to be important for the excretion of glucuronides^{31, 32}, but further investigation into this mechanism is necessary.

Taken together, the effect of siRNA treatment on intact cellular excretion points to the significance of a coupled action between flavonoid conjugating enzyme and efflux transporters in that functions of efflux transporters can significantly impact the cellular excretion of phase II conjugates. As a consequence of this coupling effect, the cells can at least partially compensate for the deficiency in enzyme (UGT1A6 in this case) activities to maintain the metabolic barrier of the intestine against flavonoids. Specifically, at 5 μ M concentration, a 50% decrease in apigenin glucuronide formation (Table 2) in cell lysate resulted in only 32% decrease in its cellular excretion. This ability to compensate and therefore maintain barrier properties would help explain why flavonoids have poor bioavailabilities in humans.

In conclusion, this study demonstrates for the first time that siRNA can be used effectively to determine the contribution of a single UGT isoform in the metabolism of a substrate in intact cells and in cell lysates. The results show that UGT1A6 is mainly responsible for the cellular metabolism of apigenin, but not genistein in Caco-2 TC7 cells. This study also provides direct evidence at the cellular level that multiple UGT isoforms are involved in the metabolism of flavonoids since silencing of UGT1A6 did not completely abolish the metabolism of apigenin or genistein. Lastly, a coupled mechanism between efflux transporters and UGT enzyme isoforms may serve as a viable mechanism to compensate for the deficiency in enzyme function.

ABBREVIATIONS

RNAi, RNA interference
siRNA, small interfering RNA
UGT1A6, UDP-glucuronosyltransferases 1A6
PNP-G, p-nitrophenol glucuronide
UDPGA, uridine diphosphoglucuronic acid
HBSS, Hanks' balanced salt solution
DMEM, Dulbecco's modified Eagle's medium
FBS, fetal bovine serum
MPA, Mobile phase A
MPB, Mobile phase B
TBS, TRIS buffer solution
GAPDH, Glyceraldehyde-3-phosphate dehydrogenase

REFERENCES

1. Farah M, Parhar K, Moussavi M, Eivemark S, Salh B. 5,6-Dichloro-ribifuranosylbenzimidazole- and apigenin-induced sensitization of colon cancer cells to TNF-alpha- mediated apoptosis. *Am J Physiol Gastrointest Liver Physiol* 2003;285(5):G919–G928. [PubMed: 12842827]
2. Gupta S, Afaq F, Mukhtar H. Involvement of nuclear factor-kappa B, Bax and Bcl-2 in induction of cell cycle arrest and apoptosis by apigenin in human prostate carcinoma cells. *Oncogene* 2002;21(23):3727–3738. [PubMed: 12032841]
3. Hessenauer A, Montenarh M, Gotz C. Inhibition of CK2 activity provokes different responses in hormone-sensitive and hormone-refractory prostate cancer cells. *Int J Oncol* 2003;22(6):1263–1270. [PubMed: 12738992]
4. Kobayashi T, Nakata T, Kuzumaki T. Effect of flavonoids on cell cycle progression in prostate cancer cells. *Cancer Lett* 2002;176(1):17–23. [PubMed: 11790449]
5. Owen RW, Haubner R, Mier W, Giacosa A, Hull WE, Spiegelhalder B, Bartsch H. Isolation, structure elucidation and antioxidant potential of the major phenolic and flavonoid compounds in brined olive drupes. *Food Chem Toxicol* 2003;41(5):703–717. [PubMed: 12659724]
6. Birt DF, Hendrich S, Wang W. Dietary agents in cancer prevention: flavonoids and isoflavonoids. *Pharmacol Ther* 2001;90(2–3):157–177. [PubMed: 11578656]
7. Piersen CE. Phytoestrogens in botanical dietary supplements: implications for cancer. *Integr Cancer Ther* 2003;2(2):120–138. [PubMed: 15035899]
8. King RA, Bursill DB. Plasma and urinary kinetics of the isoflavones daidzein and genistein after a single soy meal in humans. *Am J Clin Nutr* 1998;67(5):867–872. [PubMed: 9583843]
9. Setchell KD, Brown NM, Desai P, Zimmer-Nechemias L, Wolfe BE, Brashear WT, Kirschner AS, Cassidy A, Heubi JE. Bioavailability of pure isoflavones in healthy humans and analysis of commercial soy isoflavone supplements. *J Nutr* 2001;131(4 Suppl):1362S–1375S. [PubMed: 11285356]
10. Busby MG, Jeffcoat AR, Bloedon LT, Koch MA, Black T, Dix KJ, Heizer WD, Thomas BF, Hill JM, Crowell JA, Zeisel SH. Clinical characteristics and pharmacokinetics of purified soy isoflavones: single-dose administration to healthy men. *Am J Clin Nutr* 2002;75(1):126–136. [PubMed: 11756070]
11. Kurzer MS, Xu X. Dietary phytoestrogens. *Annu Rev Nutr* 1997;17:353–381. [PubMed: 9240932]
12. Chen J, Lin H, Hu M. Absorption and metabolism of genistein and its five isoflavone analogs in the human intestinal Caco-2 model. *Cancer Chemother Pharmacol* 2005;55(2):159–169. [PubMed: 15455178]
13. Chen J, Lin H, Hu M. Metabolism of flavonoids via enteric recycling: role of intestinal disposition. *J Pharmacol Exp Ther* 2003;304(3):1228–1235. [PubMed: 12604700]
14. Hu M, Chen J, Lin H. Metabolism of flavonoids via enteric recycling: mechanistic studies of disposition of apigenin in the Caco-2 cell culture model. *J Pharmacol Exp Ther* 2003;307(1):314–321. [PubMed: 12893842]
15. Liu Y, Hu M. Absorption and metabolism of flavonoids in the caco-2 cell culture model and a perused rat intestinal model. *Drug Metab Dispos* 2002;30(4):370–377. [PubMed: 11901089]
16. Doerge DR, Chang HC, Churchwell MI, Holder CL. Analysis of soy isoflavone conjugation in vitro and in human blood using liquid chromatography-mass spectrometry. *Drug Metab Dispos* 2000;28(3):298–307. [PubMed: 10681374]
17. Galijatovic A, Otake Y, Walle UK, Walle T. Induction of UDP-glucuronosyltransferase UGT1A1 by the flavonoid chrysin in Caco-2 cells--potential role in carcinogen bioinactivation. *Pharm Res* 2001;18(3):374–379. [PubMed: 11442279]
18. Walle UK, Walle T. Induction of human UDP-glucuronosyltransferase UGT1A1 by flavonoids-structural requirements. *Drug Metab Dispos* 2002;30(5):564–569. [PubMed: 11950788]
19. Jeong EJ, Liu Y, Lin H, Hu M. Species- and disposition model-dependent metabolism of raloxifene in gut and liver: role of ugt1a10. *Drug Metab Dispos* 2005;33(6):785–794. [PubMed: 15769887]
20. Wang SW, Chen J, Jia X, Tam VH, Hu M. Disposition of Flavonoids via Enteric Recycling: Structural Effects and Lack of Correlations between In Vitro and In Situ Metabolic Properties. *Drug Metab Dispos*. 2006

21. Hu M, Chen J, Zheng L, Dantzig AH, Stratford RE Jr. Uptake characteristics of loracarbef and cephalexin in the Caco-2 cell culture model: effects of the proton gradient and possible presence of a distinctive second component. *J Pharm Sci* 1996;85(7):767–772. [PubMed: 8819004]
22. Hu M, Chen J, Zhu Y, Dantzig AH, Stratford RE Jr, Kuhfeld MT. Mechanism and kinetics of transcellular transport of a new beta-lactam antibiotic loracarbef across an intestinal epithelial membrane model system (Caco-2). *Pharm Res* 1994;10:1405–1413. [PubMed: 7855043]
23. Elbashir SM, Harborth J, Weber K, Tuschl T. Analysis of gene function in somatic mammalian cells using small interfering RNAs. *Methods* 2002;26(2):199–213. [PubMed: 12054897]
24. Yamaoka K, Nakagawa T, Uno T. Application of Akaike's information criterion (AIC) in the evaluation of linear pharmacokinetic equations. *J Pharmacokinet Biopharm* 1978;6(2):165–175. [PubMed: 671222]
25. Hanioka N, Jinno H, Tanaka-Kagawa T, Nishimura T, Ando M. Determination of UDP-glucuronosyltransferase UGT1A6 activity in human and rat liver microsomes by HPLC with UV detection. *J Pharm Biomed Anal* 2001;25(1):65–75. [PubMed: 11274859]
26. Hutzler JM, Tracy TS. Atypical kinetic profiles in drug metabolism reactions. *Drug Metab Dispos* 2002;30(4):355–362. [PubMed: 11901086]
27. Burchell B. Genetic variation of human UDP-glucuronosyltransferase: implications in disease and drug glucuronidation. *Am J Pharmacogenomics* 2003;3(1):37–52. [PubMed: 12562215]
28. Zheng Z, Fang JL, Lazarus P. Glucuronidation: an important mechanism for detoxification of benzo [a]pyrene metabolites in aerodigestive tract tissues. *Drug Metab Dispos* 2002;30(4):397–403. [PubMed: 11901093]
29. Zhang S, Yang X, Morris ME. Combined effects of multiple flavonoids on breast cancer resistance protein (ABCG2)-mediated transport. *Pharm Res* 2004;21(7):1263–1273. [PubMed: 15290869]
30. Zhang S, Yang X, Morris ME. Flavonoids are inhibitors of breast cancer resistance protein (ABCG2)-mediated transport. *Mol Pharmacol* 2004;65(5):1208–1216. [PubMed: 15102949]
31. Zamek-Gliszczynski MJ, Hoffmaster KA, Humphreys JE, Tian X, Nezasa KI, Brouwer KL. Differential involvement of Mrp2 (Abcc2) and Bcrp (Abcg2) in biliary excretion of 4-methylumbelliferyl glucuronide and sulfate in the rat. *J Pharmacol Exp Ther*. 2006
32. Zamek-Gliszczynski MJ, Nezasa KI, Tian X, Kalvass JC, Patel NJ, Raub TJ, Brouwer KL. The important role of Bcrp (Abcg2) in the biliary excretion of sulfate and glucuronide metabolites of acetaminophen, 4-methylumbelliferone, and harmol in mice. *Mol Pharmacol*. 2006

ACKNOWLEDGMENT

This work was supported by National Institutes of Health grant CA87779. Our work started when XL and MH were at Washington State University, where we received outstanding technical contribution from Ms. Huimin Lin and Dr. Jun Chen. The work was completed at University of Houston, College of Pharmacy.

Fig. 1.
Chemical Structures of Apigenin, Genistein, and p-Nitrophenol.

Fig. 2. Representative HPLC profiles of p-nitrophenol (P), p-nitrophenol-glucuronide (Pg), apigenin (A), apigenin-glucuronide (Ag), genistein (G), genistein-glucuronide (Gg), and testosterone (IS). The HPLC conditions were described in the “Method” section. The retention time of each aglycone and its glucuronide is listed in Table. 1.

Fig. 3A **Fig. 3B** **Fig. 3.**

Effects of siRNA Treatment on Cellular UGT1A6 Activities and Expression. In Fig.3A, p-nitrophenol (50 μ M) glucuronidation rates were measured in Caco-2 cells treated with Lipofectamine alone (control), negative control (nonsense) siRNA pool (Non-control) or siRNA targeted to human UGT1A6. Fig.3B showed protein levels of UGT1A6 (top) and GAPDH (bottom) in Caco-2 cells transfected with control (buffer alone) (lanes 1–3), 50 nM siRNA-UGT1A6 (lanes 4–6) and Non-target control (lanes 7–9) using Western blot. The numerical values below the bands were average of three densitometer readings and associated standard deviation of the means. The cells were double-transfected within 24 hrs and grown for 72 hrs before experiments. Each data point represents the average of three determinations and the error bar represents the standard deviation of the mean.

Fig. 4A **Fig. 4B** **Fig. 4.**

Effect of Different siRNA Sequences on Cellular Glucuronidation Rates of p-Nitrophenol (50 μ M) in Caco-2 Cell Lysates. All siRNAs were targeted to human UGT1A6 but had different sequences and each was used at equal molar amount for 72 hrs. Each bar represents the average of three determinations (normalized arbitrary values) and the error bar represents the standard deviation of the mean (Fig.4A). Fig.4B shows the Western blot for protein levels of UGT1A6 (top) and GAPDH (bottom) in Caco-2 cells transfected with control (buffer alone) (lanes 1, 2), 50 nM siRNA-D1 (lanes 3, 4), siRNA-D2 (lanes 5, 6), siRNA-D3 (lanes 7, 8), siRNA-D4 (lanes 9, 10). The band density ratio below the bands is the average of the two lanes.

Fig. 5A Apigenin **Fig. 5B Genistein**

Fig. 5. Effects of siRNA Treatment on the Cellular Excretion of Apigenin (Fig.5A) and Genistein (Fig.5B) Glucuronides from the Caco-2 Cells. Cell monolayers were incubated with two different concentrations of the flavonoids (5 and 20 μ M) and metabolites produced were followed as a function of time, with or without siRNA treatment using protocol described in Materials and Methods. Each data point represents the average of three determinations and the error bar represents the standard deviation of the mean.

Fig. 6A **Fig. 6B** **Fig. 6C**

Fig. 6. Metabolism of Apigenin (A), Genistein (B), and p-Nitrophenol (C) by Expressed Human UGTs. The experiments were performed using 10 μ M of flavonoids and 50 μ M of p-nitrophenol for 4 hours using procedures described in “Method” section.

Fig. 7A **Fig. 7B**

Fig. 7. Metabolism of Apigenin in Caco-2 Cell Lysates without (solid line) or with siRNA Treatment (dash line) ($n = 3$). Rates of metabolism were determined from 0.56 to 100 μM , and reaction time was for 90 min (Fig. 7A). Each data point represents the average of three determinations and the error bar represents the standard deviation of the mean. Fig. 7B shows the Eadie-Hofstee plots of the same data without (solid square) or with siRNA treatment (empty circle).

Fig. 8A **Fig. 8B**

Fig. 8. Metabolism of Genistein in Caco-2 Cell Lysates without (solid line) or with siRNA Treatment (dash line) ($n = 3$). Rates of metabolism were determined from 0.5 to 200 μM , and reaction time was 480 min (Fig. 8A). Each data point represents the average of three determinations and the error bar represents the standard deviation of the mean. Fig. 8B shows the Eadie-Hofstee plots of the same data without (solid square) or with siRNA treatment (empty circle).

Table 1

Retention time of PNP, apigenin, genistein and their glucuronides. The HPLC conditions were described in the method section.

Compounds	Wavelength (nm)	Retention time (min)
PNP (P)	254	21.9
PNP-glucuronides (Pg)	264	14.3
Apigenin (A)	340	17.5
Apigenin-glucuronides (Ag)	340	9.7
Genistein (G)	254	30.0
Genistein-glucuronides (Gg)	242	21.2
Testosterone(internal standard, IS)	254	Variable, but always the last peak

Table 2

Glucuronidation of apigenin in Caco-2 cell lysate without (control) or with siRNA treatment. The siRNA was targeted to human UGT1A6 and the cells were treated twice with siRNA and used at 72 hrs after the first treatment (see “Methods”; for details).

Concentration of apigenin	Glucuronidation rates (pmol/min/mg)±SD	
	Control	siRNA Treated
0.5 µM	2.86±0.33	Below Detection*
5.0 µM	57.8±6.5	30.2±2.1 *
20 µM	297.9±44.5	128.5±9.8 *
Concentration of genistein		
0.5 µM	1.45±0.21	2.00±0.25
10 µM	12.5±2.2	9.60±1.53
50 µM	45.8±1.5	31.4±3.4*

Each data point represents the average of three determinations. The star symbol indicates there were statistically significant differences between rates of metabolism in Caco-2 cell lysate with UGT1A6 gene silencing versus control ($P<0.05$).

Table 3
Kinetic parameters of apigenin glucuronidation in Caco-2 cell lysates prepared from control and siRNA treated cells using three different kinetic models.

Model		1	2	3
Description		Classic Michaelis-Menten	Autoactivation without residual activity	Autoactivation with residual activity
Number of parameters		2	3	4
Best-fit parameter values±CV				
Control	$V_{\max,0}$ (nmol/min/mg)	0.711±0.067	0 (fixed)	0.030±1.11
	$V_{\max,d}$ (nmol/min/mg)	0 (fixed)	0.574±0.03	0.542±0.068
	K_m (μM)	33.7±0.2	13.1±4.2	12.6±0.4
	R	0 (fixed)	0.091±0.238	0.0833±0.292
	r^2	0.99	0.999	0.999
	AIC	-55.0	-86.3	-85.1
siRNA-treated	$V_{\max,0}$ (nmol/min/mg)	0.235±0.074	0 (fixed)	0.8E-08
	$V_{\max,d}$ (nmol/min/mg)	0 (fixed)	0.191±0.011	0.191±0.08
	K_m (μM)	23.3±0.2	5.31±1.03	5.32±1.86
	R	0 (fixed)	0.0788±0.46	0.0788±0.63
	r^2	0.979	0.994	0.994
	AIC	-73.3	-89.3	-87.3