Published in final edited form as:

Tetrahedron Lett. 2008 September 8; 49(37): 5363–5365. doi:10.1016/j.tetlet.2008.06.123.

Ring expansion of substituted norbornadienes for the synthesis of mono- and disubstituted 2-azabicyclo[3.2.1]octadienes

Nova Emelda and Stephen C. Bergmeier^{*}

Department of Chemistry & Biochemistry, Clippinger Laboratories, Ohio University, Athens, Ohio, 45701, USA

Abstract

We have studied the conversion of substituted norbornadienes into a substituted 2-azabicyclo[3.2.1]octadiene system via reaction with toluenesulfonyl azide. We have found that both mono- and disubstituted norbornadienes will undergo the cycloaddition/rearrangement sequence to provide the bicyclooctadiene ring system as a single regioisomer. The 2-azabicyclo[3.2.1]octane ring system can be prepared from the unsaturated 2-azabicyclo[3.2.1]octadiene ring system.

The 2-azabicyclo[3.2.1]octane ring system **1** (Figure 1) is found in a number of natural products including nordipertenoid alkaloids such as methyllycaconitine^{1–3} as well as himandrine.⁴ In addition a number of nonnatural biologically active ring systems contain the 2-azabicyclo[3.2.1]octane ring system.^{5,6} The development of new methods for the synthesis of this ring system is important both for natural product synthesis as well as the preparation of simpler analogues of these natural products. An attractive option for the synthesis of substituted derivatives of **1** would be the functionalization of the readily available N-phenylsulfonyl 2-azabicyclo[3.2.1]octadiene **2a**. N-phenylsulfonyl 2-azabicyclo[3.2.1]octadiene was first reported in 1965.⁷ This unique compound was prepared by the reaction of phenylsulfonyl azide with norbornadiene (**3a**). While the yields are only moderate the reaction is readily scalable and easy to carry out. It has been reported that the reaction proceeds through an initial dipolar cycloaddition of the azide followed by loss of nitrogen to form the fused-ring aziridine. This then undergoes a ring opening reaction to a bicyclo[3.1.0]hexene imine. This highly strained bicyclic ring system then undergoes a Cope rearrangement to form the observed 2-azabicyclo[3.2.1]octadiene **2a**.⁸

Two options exist for the conversion of **2** into the saturated and substituted ring system **1**. The first is the chemical modification of the parent bicyclo[3.2.1]octadiene ring system. Few examples of such modifications have been reported. These modifications are largely limited to the reduction of one or both double bonds,^{9–11} dihydroxylation of the 6,7-olefin¹² and cycloaddition reactions with the 6,7-olefin.^{13,14}

Given the limited methods for the functionalization of **2a**, an alternate option for the synthesis of substituted systems such as **1** is the use of substituted norbornadienes **3b** in the addition/rearrangement reaction. The product of this reaction, a substituted 2-azabicyclo[3.2.3]octadiene **2b** could then be further modified via reduction, dihydroxylation, or cycloadditions

© 2008 Elsevier Ltd. All rights reserved.

Corresponding author. Tel.: +1-740-517-8462; fax: +1-740-593-01148; e-mail: bergmeis@ohio.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

to provide highly substituted derivatives of **1**. This strategy has several advantages, including the relative ease with which some substituted norbornadienes can be prepared and the ability to further substitute the products of such an addition/rearrangement process. This would then produce derivatives of **1** with a much wider range of substitution and functionality. A key unknown is the regioselectivity of the cycloaddition/rearrangement process with a substituted norbornadiene.

Given a unique substituted norbornadiene **4** (Figure 2), the reaction with a sulfonyl azide could produce four possible products when $R^1 \neq R^2$ (or two when $R^1 = R^2$). Reaction on the opposite side of the substitution would provide regioisomeric products **5** and **6**, while reaction on the same side would provide regioisomers **7** and **8**. One report has briefly examined this reaction.¹⁵ Treatment of a compound where $R^1 = R^2 = \text{CN}$ or $R^1 = R^2 = \text{CO}_2\text{Me}$ provide largely a single product **5**, where the addition of the phenylsulfonyl azide occurred distal to the substitution. We wished to determine if other functional groups, where $R^1 = R^2$, would provide the same level of regiocontrol. More interestingly, would be an examination of the reaction of **4** where $R^1 \neq R^2$. Such compounds would likely provide only **4** and **5** but *a priori* it is not obvious which of these two regioisomers will be formed.

In order to begin to address these questions several monosubstituted norbornadienes were prepared as outlined in Scheme 1. The Diels Alder cycloaddition of ethyl propiolate with cyclopentadiene provided monosubstituted norbornadiene **4a** in 83% yield.¹⁶ While the Diels Alder method provides an excellent route for the synthesis of norbornadienes with electron withdrawing substituents, an alternate approach was needed for the synthesis of non-electron withdrawing substituted norbornadienes. Alcohol **4b** can be readily prepared via metallation and reaction of norbornadiene with formaldehyde.¹⁷ The resulting alcohol was acylated or silylated to provide monosubstituted norbornadienes **4c**,¹⁸ **4d**, and **4e**.

With a group of monosubstituted norbornadienes in hand, their reaction with tosyl azide was investigated. Based on previous studies,¹⁵ it was expected that addition would occur distal to the substitution. In concert with the original reaction conditions and in contrast to the report by Umano and co-workers,¹⁵ all reactions were carried out at room temperature.¹⁹

The reaction of **4a** with tosyl azide resulted in only a 6% yield of an 85:15 mixture of **5a** and **6a** were obtained. This reaction provided multiple products. The reaction of the acetate provided a significantly better yield of the product in a 94:6 mixture of **5c**:**6c**. Changing the ester to a silyl ether (**4d**) lowered the yield somewhat (50% to 30%) but dramatically improved the product ratio yielding only the 7-substituted product **5d**. Suspecting that this moderate change in regioisomer ratio might be due to the size of the group on the oxygen, the pivalate ester **4e** was prepared. The yield returned to the moderate yield observed with the acetate and retained the improved ratio of regioisomers. Clearly the substitution of these monosubstituted norbornadienes is of paramount importance. Electron withdrawing substitution provides little to no product while the substituted hydroxymethyl derivatives provided a moderate yield of a single regioisomer.

In order to investigate regioselectivity in the reaction of disubstituted norbornadienes, a series of disubstituted norbornadienes were prepared via the Diels Alder reaction of cyclopentadiene with substituted alkynes (Scheme 2). The reaction of diethyl acetylene dicarboxylate with cyclopentadiene provided norbornadiene **4f** in 97% yield. Reaction of the diacetate of butyne diol with cyclopentadiene required microwave heating of the reaction to 220 °C in DMF and provided **4g**²⁰ in 50% yield.

In order to prepare norbornadienes where $R^1 \neq R^2$, cyclopentadiene was heated with several substituted propiolate esters in a sealed tube at 160 °C. Norbornadiene **4h**²¹ was obtained in an almost quantitative yield. Compounds **4i** and **4j** were prepared in moderate yield, however

they could not be completely purified due to contamination with ~40% of an unidentified non-polar impurity that could not be readily removed.

The ring expansion of symmetrically disubstituted norbornadiene **4f** was examined first. Following the reaction conditions that worked well for norbornadiene itself as well as the monosubstituted norbornadienes provided only a 4:1 mixture of aziridine **10** and rearrangement product **5f**. Heating the reaction to 80 °C for 18 hours provides the same mixture of aziridine and rearrangement product. Increasing the heat to 110 °C (refluxing toluene) shows a steadily increasing amount of the rearrangement product. After 3 days in refluxing toluene only the rearrangement product was observed.

With a viable procedure in hand a range of disubstituted norbornadienes were examined. The diester **4f** provided **5f** in 94% isolated yield. The yield obtained in refluxing toluene is similar to that obtained by Umano et al in refluxing 1,2-dichlorobenzene.¹⁵ However the use of toluene provides a somewhat more convenient procedure than using 1,2-dichlorobenzene as the solvent. The diacetate **4g** provided **5g** in 50% yield.

Several nonsymmetrical derivatives were next examined. Our prediction was that the larger of the two groups R¹ or R² would end up on the same side of the bicyclic ring as the N-Ts group. A single product, **5h**, was obtained upon treatment of **4h** with tosyl azide. The regiochemistry was determined through NOESY experiments that showed a crosspeak between H1 and H1' on the phenyl ring. This is consistent with the reactions of the monosubstituted norbornadienes **4a** – **4e** in which the larger group was on the same side of the ring as the N-Ts group. The reaction of unsymmetrically disubstituted norbornadienes **4i** and **4j** provided identical regiochemical results. While the yields of **5i** and **5j** were not as good as **5h**, these were clean reactions providing only a single regioisomer. The regiochemistry of **5i** and **5j** was also assigned by the presence of a similar crosspeak in a NOESY spectrum, between the bridgehead H1 and a methylene proton of the alkyl group at C7.

We also wished to examine the reaction of the benzofused norbornadiene **11**²³ with toluenesulfonyl azide. Reaction of **11** with tosyl azide provided only the aziridine **12**²⁴ in 77% yield. All attempts to convert **12** to the desired **13** gave either no reaction or complete decomposition. These reaction conditions include heating **12** to over 250 °C, or the addition of a number of Lewis acids with or without heating. In trying to determine the reason for the lack of reaction of **12**, Umano and coworkers determined that endo aziridines (e.g. endo-**10**) do not undergo the rearrangement reaction to provide the [3.2.1]bicyclic ring system.¹⁵ We carried out a NOESY experiment with **12** in order to determine the relative stereochemistry of the aziridine. We observed no crosspeaks between the aziridine protons and the bridgehead protons, indicating a likely exo-aziridine. Consequently, it may be that the rigidity of the system compounded with the loss of aromaticity in the rearrangement reaction preclude the rearrangement of this system.

In conclusion, we have found that mono- and disubstituted norbornadienes undergo an addition/rearrangement reaction with tosyl azide. The reaction proceeds with excellent levels of regiocontrol in modest to excellent yield. This reaction provides an excellent route for the synthesis of substituted 2-azabicyclo[3.2.1]octadienes. These substituted 2-azabicyclo[3.2.1]octadienes can be readily converted to the reduced bicyclooctane ring system found in a number of natural products and pharmacologically active molecules.

Acknowledgments

We would like to thank The National Agency of Drug & Food Control, Republic of Indonesia for a graduate fellowship (NE). We would like to thank the National Institutes on Drug Abuse at the NIH (DA13939) for partial support of this work and Ohio University for support of the BioMolecular Innovation and Technology (BMIT) project.

References

1. Ward JM, Cockcroft VB, Lunt GG, Smillie FS, Wonnacott S. FEBS Lett 1990;270:45. [PubMed: 2226787]
2. Goodall KJ, Barker D, Brimble MA. Synlett 2005:1809.
3. Manske RH. Can. J. Res 1938;16B:57.
4. O'Connor PD, Mander LN, McLachlan MMW. Org. Lett 2004;6:703. [PubMed: 14986954]
5. Gowrisankar S, Lee HS, Lee KY, Lee J-E, Kim JN. Tetrahedron Lett 2007;48:8619.
6. Noguchi K, Takeda M, Nurimoto S. Chem. Pharm. Bull 1977;25:890. [PubMed: 264181]
7. Oehlschlager AC, Zalkow LH. J. Org. Chem 1965;30:4205.
8. Huda E, Martin HD, Mayer B, Somnitz KH, Steigel A, Haddad H, Distefano G, Modelli A. Chem. Ber 1991;124:2879.
9. Barraclough P, Bilgic S, Pedley JB, Rogers AJ, Young DW. Tetrahedron 1979;35:99.
10. Barraclough P, Bilgic S, Young DW. Tetrahedron 1979;35:91.
11. Rhee H, Yoon DO, Jung ME. Nucleosides, Nucleotides and Nucleic Acids 2000;19:619.
12. Reed DD, Bergmeier SC. J. Org. Chem 2007;72:1024. [PubMed: 17253828]
13. Inoue H, Tokisato K, Umamo K. Bull. Chem. Soc. Japan 1982;55:1661.
14. Umamo K, Mizone S, Tokisato K, Inoue H. Tetrahedron Lett 1981;22:73.
15. Umamo K, Taniguchi H, Inoue H, Imoto E. Tetrahedron Lett 1979:247.
16. Wang YG, M XW, Huang X. Synthesis 2007:28.
17. Yip C, Handerson S, Tranmer GK, Tam W. J. Org. Chem 2001;66:276. [PubMed: 11429911]
18. Tanyeli C, Celikel G, Akhmedov IM. Tetrahedron-Asymmetry 2001;12:2305.
19. Tosyl azide (100 mol%) and a monosubstituted norbornadiene **4** (100 mol%) were dissolved in toluene (0.5 M). After stirring for 3 days the reaction was concentrated and chromatographed (10% EtOAc in hexanes) to provide the product.
20. Muchow G, Brunel JM, Maffei M, Buono G. J. Org. Chem 1995;60:852.
21. Bauml E, Mayr H. J. Org. Chem 1983;48:2600.
22. Tosyl azide (100 mol%) and a disubstituted norbornadiene **4** (200 mol%) were dissolved in toluene (0.5 M). The reaction was stirred at rt for 24 h then heated to reflux for 72 h. The reaction was concentrated and chromatographed to provide the product and recovered starting norbornadiene.
23. Brooks PR, Caron S, Coe JW, Ng KK, Singer RA, Vazquez E, Vetelino MG, Watson HH, Whitenour DC, Wirtz MC. Synthesis 2004:1755.
24. Martin MM, Koster RA. J. Org. Chem 1968;33:3428.

Figure 1.
2-Azabicyclo[3.2.1]octane ring system.

Figure 2. Possible products derived from the reaction of a substituted norbornadiene with a sulfonyl azide.

Figure 3.
Observed NOESY crosspeak for product **5h**.

Scheme 1.

(a) CH_2Cl_2 , reflux, 18 h, 83%. (b) CH_2Cl_2 , AcCl, pyridine, DMAP, 98%. (c) CH_2Cl_2 , $\text{Me}_2\text{tBuSiCl}$, Et_3N , DMAP, 98%. (d) CH_2Cl_2 , pivaloyl chloride, Et_3N , DMAP, 98%.

Scheme 2.

4f, $\text{R}^1 = \text{COOEt}$, $\text{R}^2 = \text{COOEt}$, toluene, reflux, 18 h, 97%. **4g**, $\text{R}^1 = \text{CH}_2\text{OAc}$, $\text{R}^2 = \text{CH}_2\text{OAc}$, DMF, microwave, 220 °C, 2 h, 50%. **4h**, $\text{R}^1 = \text{Ph}$, $\text{R}^2 = \text{COOMe}$, toluene, sealed tube, 160 °C, 42 h, 99%. **4i**, $\text{R}^1 = n\text{C}_5\text{H}_{11}$, $\text{R}^2 = \text{COOEt}$, toluene, sealed tube, 160 °C, 60 h, 56%*. **4j**, $\text{R}^1 = \text{CH}_2\text{OPh}$, $\text{R}^2 = \text{COOMe}$, toluene, sealed tube, 160 °C, 60 h, 56%*.

Scheme 3.
Time and temperatures studies on **4f**

Scheme 4.

Attempted reaction of benzonorbornadiene with tosyl azide

Table 1Reaction of monosubstituted norbornadienes^{a, b}

R¹	Yield (%)	Ratio 5:6
4a , CO ₂ Et	6	85:15 5a:6a
4c , CH ₂ OAc	50	94:6, 5c:6c
4d , CH ₂ OTBS	30	Only 5d
4e , CH ₂ OPiv	50	Only 5e

^a When R² = H, R² is not shown for clarity.^b The regiochemistry of products **5a** – **5e** was determined by the observation of coupling between H5 (δ 2.62 – 3.01) and H6 (δ 5.93 – 6.09) as determined via COSY experiments.

Table 2Reactions of disubstituted norbornadienes.²²

Starting material	R ¹	R ²	% Yield, ^a Product
4f	CO ₂ Et	CO ₂ Et	94, 5f
4g	CH ₂ OAc	CH ₂ OAc	50, 5g
4h	CO ₂ Me	Ph	98, 5h
4i	CO ₂ Et	nC ₅ H ₁₁	40, 5i
4j	CO ₂ Me	CH ₂ OPh	33, 5j

^aYields are relative to tosyl azide.