CORRESPONDENCE

Array comparative genomic hybridisation analysis of boys with X-linked hypopituitarism identifies a 3.9 Mb duplicated critical region at Xq27 containing SOX3

In our previous article,¹ we used array-comparative genomic hybridisation (CGH) analysis to identify Xq26–q27 duplications in families with X-linked hypopituitarism (XH). The array-CGH assay was validated using affected male genomic DNA from two previously characterised XH families that carry Xq26–q27 duplications^{2,3} (fig 1B,C) and by interphase fluorescence in situ hybridisation (FISH) analysis (fig 2). Array-CGH analysis of three novel XH families, A, B and C (fig 3), indicated that each of these contained a different Xq26–q27 duplication (fig 1D–G).

Recently, we repeated these array-CGH experiments using a more extensive X chromosome array containing over 2000 bacterial artificial chromosome (BAC) clones.⁴ As expected, duplications were identified in males from the two previously characterised XH families.^{2,3} However, we have been unable to detect Xq26–q27 duplications in affected males from families A, B and C. Furthermore,

repeated quantitative real-time PCR experiments performed by an independent collaborator in a blinded assay detected *SOX3* duplication in affected males from control families,^{2,3} but not in affected males from families A, B and C. On the basis of these new data, we now cannot replicate the observation of duplications in families A, B and C families that was published in our paper.

Nicola M Solomon, Shelley A Ross, Susan M Forrest, Paul Q Thomas

Murdoch Childrens Research Institute, Royal Children's Hospital, Melbourne, Australia

Thomas Morgan, Joseph L Belsky

Yale University School of Medicine, Department of Genetics, New Haven, USA

Frans A Hol

Department of Human Genetics, University Medical Center Nijmegen, The Netherlands

Pamela S Karnes

Mayo Clinic, University of Medical Genetics, Rochester, Minnesota, USA

Nancy J Hopwood

University of Michigan Medical Center, Mississippi, USA

Susan E Myers

Saint Louis University, Missouri, USA

Anjanette S Tan

Division of Endocrinology, Diabetes, and Metabolism, University of Missouri-Columbia, Columbia, Missouri, USA

Garry L Warne

Department of Endocrinology, Royal Children's Hospital, Melbourne, Australia

doi: 10.1136/jmg.2007.049049

Received 7 January 2007
Accepted 9 January 2007

Competing interests: None declared.

References

- 1 **Solomon NM**, Ross SA, Morgan T, Belsky JL, Hol FA, Karnes PS, Hopwood NJ, Myers SE, Tan AS, Warne GL, Forrest SM, Thomas PQ. Array comparative genomic hybridisation analysis of males with X-linked hypopituitarism identifies a 3.9 Mb duplicated critical region at Xq27 containing *SOX3*. *J Med Genet* 2004;**41**:669–78.
- 2 **Lagerström-Fermér M**, Sundvall M, Johnsen E, Warne GL, Forrest SM, Zajac JD, Rickards A, Ravine D, Landegren U, Pettersson U. X-linked recessive panhypopituitarism associated with a regional duplication in Xq25–q26. *Am J Hum Genet* 1997;**60**:910–16.
- 3 **Hol FA**, Schepens MT, van Beersum SE, Affer M, Vezzoni P, Hamel BC, Karnes PS, Mariman EC, Zucchi I. Identification and characterisation of an Xq26–q27 duplication in a family with Spina-Bifida and panhypopituitarism suggests the involvement of two distinct genes. *Genomics* 2000;**69**:174–81.
- 4 **Bauters M**, Van Esch H, Marynen P, Froyen G. X chromosome array-CGH for the identification of novel X-linked mental retardation genes. *Eur J Med Genet* 2005;**48**:263–75.