Published in final edited form as:

Theochem. 2008 August 30; 863(1-3): 39–43. doi:10.1016/j.theochem.2008.05.014.

Conformation dependence of pKa's of the chromophores from the purple asFP595 and yellow zFP538 fluorescent proteins

A.V. Nemukhin^{a,b}, I.A. Topol^c, B.L. Grigorenko^a, A.P. Savitsky^d, and J.R. Collins^c

^aDepartment of Chemistry, M.V. Lomonosov Moscow State University, Leninskie Gory, 1/3, Moscow, 119991, Russian Federation

^bN.M. Emanuel Institute of Biochemical Physics, Russian Academy of Sciences, ul. Kosygina, 4, Moscow, 119334, Russian Federation

^cAdvanced Biomedical Computing Center, Advanced Technology Program, SAIC-Frederick Inc., NCI-Frederick, Frederick, MD, 21702-1201, USA

^dA.N. Bach Institute of Biochemistry, Russian Academy of Sciences, Leninsky prospekt, 33, Moscow, 119071, Russian Federation

Abstract

Two members of the green fluorescent protein family, the purple asFP595 and yellow zFP538 proteins, are perspective fluorescent markers for use in multicolor imaging and resonance energy-transfer applications. We report the results of quantum based calculations of the solution pKa values for selected protonation sites of the denatured asFP595 and zFP538 chromophores in the trans- and cis-conformations in order to add in the interpretation of photophysical properties of these proteins. The pKa values were determined from the thermodynamic cycle based on B3LYP/6-311++G(2df, 2p) calculations of the gas phase free energies of the molecules and the B3LYP/6-311++G(d,p) calculations of solvation energies. The results show that the pKa's of the protonation sites of the chromophore from asFP595 noticeably depend on the isomer conformation (cis- or trans-), while those of zFP538 are much less sensitive to isomerization.

Keywords

biological chromophores; pKa; cis-trans isomerization; fluorescent proteins

Introduction

The properties of chromophores from the extended family of the green fluorescent protein (GFP) are being intensively investigated because of great importance of GFP-like biomarkers in living cells [1]. In this work we focus on two members of the family: the purple chromoprotein from the sea anemone *Anemonia sulcata*, asFP595 [2], and the yellow chromoprotein from button polyp *Zoanthus*, zFP538 [3], which are perspective fluorescent

© 2008 Elsevier B.V. All rights reserved.

Fax: +74959390283; Tel: +74949391096; E-mail: anem@lcc.chem.msu.ru.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

markers for use, in particular, in multicolor imaging and resonance energy-transfer applications.

The fluorescent protein asFP595, which is the Ala143Gly mutant of the wild type chromoprotein asCP, is famous for its unique photo-physical properties. The protein asFP595 is initially nonfluorescent, but in response to intense green light irradiation at 568 nm, it becomes brightly fluorescent (kindles) with emission at 595 nm. Irradiation of the emitting protein form with blue light quenches fluorescence. The photoswitching properties of this kindling fluorescent protein may be also useful for information storage in macromolecules or for creating triggerable markers in living cells. A large amount of experimental data [4-9] indicate that the mechanism of kindling may be related to the trans- to cis- photo-isomerization of the chromophore. The most relevant direct observations are due to Andresen et al. [7] who presented the crystal structures of the Ala143Ser mutant of asCP. In the dark state of these crystals, the chromophore was in the trans-form, but after irradiation the chromophore could be visualized in the cis-configuration.

Unlike asFP595, the protein zFP538 is not known as a photoswitching species. Absorption and fluorescence of zFP538 containing a three-ring chromophore [3] correspond to the wavelengths intermediate between fluorescence of green proteins and absorption of red proteins. This property allows one to use zFP538 for the resonance energy transfer processes important in analytical applications.

The chromophores of asFP595 and zFP538 may be considered as derivatives of 1 the chromophore from the parent protein GFP, 4-hydroxybenzylidene-imidazolinone (HBI), shown in Fig.1 in the anionic form. Two most important protonation sites of the molecule refer to oxygen of the phenolic ring (O_{phen}) and the imidazoline nitrogen atom (N_{imid}).

The molecules of chromophores from asFP595 (Fig.2) and zFP538 (Fig.3) possess extra π -electron conjugation accounting for absorption and emission shifts to longer wavelengths compared to GFP. The structures of the chromophores from asFP595 and zFP538 are basically known from the corresponding crystals, however, some uncertainties still remain. In particular, an alternative interpretations of X-ray experiments resulted either in the carbonyl group (as shown in the upper panel of Fig.1) [5] or the iminic group (lower panel of Fig.2) [6,7] in the asFP595 chromophore.

Many details of the fluorescent properties of these proteins remain unknown in spite of substantial efforts to understand the chemistry and physics of chromophore excitations. Specifically, the role of trans-cis isomerization of the asFP595 chromophore inside the protein is still under debate [7,10,11,12], although the working hypothesis assumes this trans- to cis-transformation. If such isomerization stimulated by photoexcitation actually occurs in the protein, it should proceed with an active involvement of the neighboring amino acid side chains. In particular, protons from these residues may be attached to electronegative sites of the chromophore modulating its spectral properties. Therefore it is important to compare proton affinities and respective pKa values of different sites of the chromophore in various conformations. The trans- and cis- forms of the chromophore are the first candidates for studies of conformational dependence of pKa's.

Previously, the protonation sites at the phenolic oxygen atom and the imidazoline nitrogen atom of the model chromophore (Fig.1) of the parent GFP have been experimentally [13] and theoretically [14-18] studied for their relative acidity (pKa). El Yazal et al. [17] calculated the theoretical pKa values for all possible protonation sites of the chromophore in denatured GFP chromophores. Scharnagl and Raupp-Kossmann [18] also reported theoretical pKa's for GFP from hybrid quantum-classical calculations with an average error of 0.8 units. In particular, the authors computed the constant pKa=8.3 for neutral/anion equilibria in excellent agreement

with the experimental data ($pK_a=8.2$) on solution-phase acid-base dissociation constants of the model GFP chromophore [13]. No comparison of pK_a 's for the original cis- and hypothetical trans- forms of the GFP chromophore were performed.

The extended chromophores (Figs.2,3) of the primary interest of this work contain additional protonation sites and one might expect certain changes in the pK_a values compared to those of GFP. The experimental data for asFP595 and zFP538 are rare: (1) The denatured asFP595 chromophore in aqueous solution was studied in the work of Yampolsky et al. [19], but no experimentally based estimates of pK_a 's were reported. (2) The seeming pK_a constants of the zFP538 chromophore in solution were evaluated from the absorption and fluorescent spectra giving the values $pK_1 \approx 5.6-5.7$ and $pK_2 \approx 7.4-7.8$ [20]. From theoretical side, the only estimate of the zwitterionic imidazolinone NH proton of the asFP595 chromophore in aqueous solution by using the density functional theory calculations which resulted in the pK values 4.7 and 9.1 for the cis- and trans- forms, respectively, was described in Ref.[11].

In order to provide better understanding the photophysical phenomena with the fluorescent proteins asFP595 and zFP538 we performed accurate quantum chemistry calculations of the solution pK_a values for selected protonation sites of the asFP595 and zFP538 chromophores in the trans- and cis-conformations. We follow the strategy combining the correlated quantum chemistry technique and continuum solvation models through the thermodynamic cycle, which is capable to determine the pK_a values accurate within half of a pK_a unit [21,22].

Calculation details

Our calculations use the thermodynamic cycle shown in Scheme 1, leading to the well-known formula

$$pK_a = (2.303RT)^{-1} \{ \Delta G_g + \Delta G_s(A^-) - \Delta G_s(AH) + \Delta G_s(H^+) \} \quad (1)$$

In Scheme 1 and formula (1) ΔG_g and ΔG_s are free energies of deprotonation in the gas phase and solvent, respectively. $\Delta G_s(A^-)$, $\Delta G_s(AH)$ and $\Delta G_s(H^+)$ are solvation free energies of anion A^- , protonated compound AH, and proton H^+ , respectively. Optimization of equilibrium geometry parameters and calculations of harmonic frequencies for the gas-phase species were performed in the density functional theory B3LYP/6-31++G(2d,2p) approximation by using the Gaussian-03 program [23]. Corrections to the free energy changes for the gas-phase part of the reaction in Scheme 1 were obtained in the single point calculations in the B3LYP/6-311++G(2df,2p) approximation. Solvation energies $\Delta G_s(A^-)$ and $\Delta G_s(AH)$ were estimated at the B3LYP/6-311++G(d,p) level by using the Jaguar program [24].

The proton solvation free the energy $\Delta G_s(H^+)$ last term in Eq.(1) is not known with high precision. Based on previous calculations of pK_a values of substituted imidazoles [21], the value -262.5 kcal/mol [22] was for a long time considered as the best estimate. However, recently Palascak and Schields [25] suggested an even lower value - 264.0 kcal/mol. In this work we used both estimates and found that although the absolute pK_a 's may differ by up to one pK_a unit, the results are qualitatively independent of $\Delta G_s(H^+)$. Below we cite only the values calculated using the value -264.0 kcal/mol.

Results

The results of calculations for all model species are summarized in Table 1.

Model GFP chromophore (Fig.1)

In the protein, the cis-conformation of the GFP chromophore occurs and no evidences of the trans-form have been reported so far. The cis-trans isomerization of a model GFP chromophore, 4-hydroxybenzylidene-1,2-dimethylimidazolinone (HBDI), was studied experimentally in aqueous solution [26].

Our computed pKa value on the phenolic oxygen site in the cis-conformation (8.5) agrees with the experimental (8.2 [13]) and previous theoretical (8.2 [17], 8.3 [18]) results within the generally accepted uncertainties of pKa calculations. Here we notice the obtained changes in pKa's due to cis-trans isomerization, $1.3 \div 1.7$ pKa units for the phenolic oxygen and the imino nitrogen of the imidazolinone ring.

zFP538 chromophore (Fig.3)

In calculations for the zFP538 chromophore we first added protons to the different sites of anionic species; the corresponding pKa's are indicated in Table 1 in the rows 'Anionic zFP538'. Remarkably, the computed constants are practically the same for both (cis- and trans-) conformers. Next set of calculations was carried out for the possible neutral chromophore with the protonated O_{phen}. These values are shown in the rows 'Neutral zFP538' in Table 1. The nitrogen site of the imidazolinone ring turns out to be the most sensitive to conformational changes (-0.2 vs 2.3). One of the goals of the present modeling of the zFP538 chromophore is to provide interpretation of the experimental measurements of the apparent constants of the chromophore in solution (pK1 $\approx$ 5.6-5.7 and pK2 $\approx$ 7.4-7.8 [20]). The computed pKa's for the nitrogen sites (4.4-4.5 and 7.0-7.4) correlate well with these experimental estimates.

asFP595 chromophore

As mentioned in Introduction, Quillin et al. [5] reported the chromophore structure for the asFP595 protein in which the conjugated-system in the GFP-like moiety is extended by the chain with the carbonyl group (upper panel in Fig.2). However, Wilmann et al. [6] and Andresen et al. [7] placed the imino group instead of the carbonyl oxygen (lower panel in Fig. 2). In all crystal structures corresponding to the dark state of the protein, the chromophore appears in the trans-conformation [5-7]. It hard to assume formation of the asFP595 chromophore with the protonated O_{phen} site in the protein; therefore, only anionic species are considered in this work. The results of calculations of pKa's shown in Table 1 demonstrate larger sensitivity of the protonation constants of the asFP595 chromophore compared to those of the zFP538 chromophore.

Discussion and conclusions

The results of calculations demonstrate that the pKa's of the protonation sites of the chromophore from the asFP595 chromoprotein noticeably depend on the isomer conformation (cis- or trans-). The most striking difference of more than 3 pK units is obtained for the imidasole nitrogen of the asFP595(O) chromophore: 1.29 for the cis-form versus 4.63 for the trans-form. Such change in the local structure of the chromophore upon isomerization may be related, among other factors, to the kindling fluorescent properties of this protein, the phenomenon which is often attributed to the occurrence of the photo-induced trans-cis isomerization of the chromophore.

Fig.4 illustrates a fragment of the asFP595(NH) protein with the trans-form of the chromophore as obtained in quantum mechanical - molecular mechanical (QM/MM) calculations [10]. Clearly, formation of the zwitterionic form of the chromophore may occur in the protein matrix when protons from the nearby amino acid side chains are transferred to nitrogen atoms N_{Imid} (or N_{Chain} in case of the asFP595(NH) form). The present calculations show that the proton

affinity of these particular chromophore atoms strongly depends on the isomer conformation affecting the location of the proton along the corresponding hydrogen bond. Even recognizing that the pKa values inside the protein may noticeably differ from those in solution the tendencies in both condense phases are believed to be similar. Study of the mechanism of kindling in asFP595 including the trans-cis chromophore isomerization by using the QM/MM approach is a subject of current research in our laboratories.

Remarkably, calculations for the chromophore from zFP538 do not show noticeable differences in solution pKa's for the trans- and cis-forms. Only if the phenolic oxygen is protonated the properties of the imidazole nitrogen are affected (-0.2 for the cis-form vs 2.3 for the trans-form). As mentioned in the Introduction, no evidence has been reported that the trans-cis isomerization affects the fluorescent properties of this protein. The only experimental data for the seeming pKa constants evaluated directly from the absorption and fluorescent spectra of the zFP538 chromophore in solution are as follows: pK1 $\approx$ 5.6-5.7 and pK2 $\approx$ 7.4-7.8 [20]. Apparently, the computed pKa's for the nitrogen sites (4.4-4.5 and 7.0-7.4) correlate well with these experimental estimates.

Acknowledgement

This work was partly supported by the Science and Innovation Federal Russian Agency (project # 02.522.11.2002), by the Program for Molecular and Cell Biology from Russian Academy of Sciences, and by the grant from the Russian Foundation for Basic Research (project #07-03-00059). We thank the staff and administration of the Advanced Biomedical Computing Center for their support of this project. This project has been funded in whole or in part with federal funds from the National Cancer Institute, National Institutes of Health, under contract N01-CO-12400. The content of this publication does not necessarily reflect the views or policies of the Department of Health and Human Services, nor does mention of trade names, commercial products, or organization imply endorsement by the U.S. Government.

References

- [1]. Zimmer M. Chem. Rev 2002;102:759. [PubMed: 11890756]
- [2]. Lukyanov KA, Fradkov AF, Gurskaya NG, Matz MV, Labas YA, Savitsky AP, Markelov ML, Zarausky AG, Zhao X, Fang Y, Tan W, Lukyanov SA. J. Biol. Chem 2000;275:25879. [PubMed: 10852900]
- [3]. Remington SJ, Wachter RM, Yarbrough DK, Branchaud B, Anderson DC, Kallio K, Lukyanov KA. Biochemistry 2005;44:202. [PubMed: 15628861]
- [4]. Chudakov DM, Feofanov AV, Mudrik NN, Lukyanov S, Lukyanov KA. J. Biol. Chem 2003;278:7215. [PubMed: 12496281]
- [5]. Quillin ML, Anstrom DM, Shu X, O'Leary S, Kallio K, Chudakov DM, Remington SJ. Biochemistry 2005;44:5774. [PubMed: 15823036]
- [6]. Wilmann PG, Petersen J, Devenish RJ, Prescott M, Rossjohn J. J. Biol. Chem 2005;280:2401. [PubMed: 15542608]
- [7]. Andresen M, Wahl MC, Stiel AC, Gräter F, Schäfer LV, Trowitzsch S, Weber G, Eggeling C, Grubmüller H, Hell SW, Jakobs S. Proc. Natl. Acad. Sci. USA 2005;102:13070. [PubMed: 16135569]
- [8]. Schüttrigkeit TA, von Feilitzsch T, Kompa C, Lukyanov KA, Savitsky AP, Voityuk AA, Michel-Beyerle M-E. Chem. Phys 2006;323:149.
- [9]. Henderson JN, Remington SJ. Physiology 2006;21:162. [PubMed: 16714474]
- [10]. Grigorenko BL, Savitsky AP, Topol IA, Burt SK, Nemukhin AV. J.Phys.Chem.B 2006;110:18735.
- [11]. Schäfer LV, Groenhof G, Klingen AR, Ullmann GM, Boggio-Pasqua M, Robb MA, Grubmüller H. Angew. Chemie Int. Ed 2007;46:530.
- [12]. Schäfer LV, Groenhof G, Boggio-Pasqua M, Robb MA, Grubmüller H. PLOS Comput. Biol 2008;4:e1000034. [PubMed: 18369426]
- [13]. Bell AF, He X, Wachter RM, Tonge PJ. Biochemistry 2000;39:4423. [PubMed: 10757992]

- [14]. Voityuk AA, Michel-Beyerle M-E, Rösch N. Chem.Phys.Lett 1997;272:162.
- [15]. Voityuk AA, Michel-Beyerle M-E, Rösch N. Chem. Phys 1998;231:13.
- [16]. Scharnagl C, Raupp-Kossmann RA, Fischer SF. Biophys. J 1999;77:1839. [PubMed: 10512807]
- [17]. El Yazal J, Prendergast FG, Shaw DA, Pang Y-P. J. Am. Chem. Soc 2000;122:11411.
- [18]. Scharnagl C, Raupp-Kossmann RA. J. Phys. Chem. B 2004;108:477.
- [19]. Yampolsky IV, Remington SJ, Martynov VI, Potapov VK, Lukyanov S, Lukyanov KA. Biochemistry 2005;44:5788. [PubMed: 15823037]
- [20]. Zubova, N. PhD thesis. Department of Chemistry, M.V. Lomonosov Moscow State University; 2006.
- [21]. Topol IA, Tawa GJ, Burt SK, Rashin AA. J. Phys. Chem. A 1997;101:10075.
- [22]. Tawa GJ, Topol IA, Burt SK, Caldwell RA, Rashin AA. J. Chem.Phys 1998;109:4852.
- [23]. Frisch, MJ.; Trucks, GW.; Schlegel, HB.; Scuseria, GE.; Robb, MA.; Cheeseman, JR.; Montgomery, JA., Jr.; Vreven, T.; Kudin, KN.; Burant, JC.; Millam, JM.; Iyengar, SS.; Tomasi, J.; Barone, V.; Mennucci, B.; Cossi, M.; Scalmani, G.; Rega, N.; Petersson, GA.; Nakatsuji, H.; Hada, M.; Ehara, M.; Toyota, K.; Fukuda, R.; Hasegawa, J.; Ishida, M.; Nakajima, T.; Honda, Y.; Kitao, O.; Nakai, H.; Klene, M.; Li, X.; Knox, JE.; Hratchian, HP.; Cross, JB.; Bakken, V.; Adamo, C.; Jaramillo, J.; Gomperts, R.; Stratmann, RE.; Yazyev, O.; Austin, AJ.; Cammi, R.; Pomelli, C.; Ochterski, JW.; Ayala, PY.; Morokuma, K.; Voth, GA.; Salvador, P.; Dannenberg, JJ.; Zakrzewski, VG.; Dapprich, S.; Daniels, AD.; Strain, MC.; Farkas, O.; Malick, DK.; Rabuck, AD.; Raghavachari, K.; Foresman, JB.; Ortiz, JV.; Cui, Q.; Baboul, AG.; Clifford, S.; Cioslowski, J.; Stefanov, BB.; Liu, G.; Liashenko, A.; Piskorz, P.; Komaromi, I.; Martin, RL.; Fox, DJ.; Keith, T.; Al-Laham, MA.; Peng, CY.; Nanayakkara, A.; Challacombe, M.; Gill, PMW.; Johnson, B.; Chen, W.; Wong, MW.; Gonzalez, C.; Pople, JA. Gaussian 03, Revision C.02. Gaussian, Inc.; Wallingford CT: 2004.
- [24]. Jaguar. Schrodinger, LLC; Portland, OR: 19912005.
- [25]. Palascak MW, Schields GC. J. Phys. Chem A 2004;108:3692.
- [26]. He X, Bell AF, Tonge PJ. FEBS Letters 2003;549:35. [PubMed: 12914920]

Figure 1. Structural formula of the GFP chromophore, 4-hydroxybenzylidene-imidazolinone, in the *cis*- (top) and *trans*- (bottom) conformations. Two important protonation sites are designated as O_{Phen} and N_{Imid} .

asFP595(O) **asFP595(NH)** **Figure 2.**

Two possible structures of the chromophore of the asFP595 protein, asFP595(O) (top) and asFP595(NH) (bottom), in the cis-anionic form. Here and below carbon atoms are designated by green, oxygen by red and nitrogen by blue.

Figure 3.
The chromophore of the zFP538 protein in the cis-anionic form.

Figure 4. Geometry configuration of the asFP595(NH) chromophore and the nearby amino acid side chains of Glu215 and Cys62 inside the protein matrix. Two possible regions of proton transfers either to the N_{Imid} or N_{Chain} protonation sites of the chromophore are indicated by black ovals.

Scheme 1.

Thermodynamic cycle for proton abstraction reaction in the gas phase (g) and in solvent (s).

Table 1

Computed pKa values for the selected protonation sites of the chromophores shown in Figs.1-3

Species	O _{Phen}	N _{Imid}	N _{Ring}	N _{Chain}
GFP, cis	8.53	5.30	---	---
GFP, trans	7.27	3.58	---	---
Anionic zFP538, cis	9.06	4.52	7.02	---
Anionic zFP538, trans	9.09	4.38	7.39	---
Neutral zFP538, cis	---	-0.18	4.13	---
Neutral zFP538, trans	---	2.33	4.21	---
asFP595(O), cis	6.96	1.29	---	---
asFP595(O), trans	8.38	4.63	---	---
asFP595(NH), cis	8.14	3.22	---	7.57
asFP595(NH), trans	7.69	4.74	---	6.77