must be provided at the most basic primary care level, and need continues to far outstrip the availability of services.

Sex Transm Infect 2007;**83**:503–505.

doi: 10.1136/sti.2007.027748

Authors' affiliations

A Boule, University of Cape Town, South Africa
N Ford, Médecins Sans Frontières, South Africa

Correspondence to: A Boule, University of Cape Town, PO Box 13203 Mowbray, South Africa; andrew.boule@uct.ac.za

Accepted 2 October 2007

Competing interests: None declared.

REFERENCES

- 1 **Gilks C**, AbouZhar C, Turmen T. HAART in Haiti—evidence needed. *Bull World Health Organ* 2001;**79**:1154–5.
- 2 **Harries AD**, Nyangulu DS, Hargreaves NJ, *et al*. Preventing antiretroviral anarchy in sub-Saharan Africa. *Lancet* 2001;**358**:410–4.
- 3 **Marseille E**, Hofmann PB, Kahn JG. HIV prevention before HAART in sub-Saharan Africa. *Lancet* 2002;**359**:1851–6.
- 4 **Loewenson R**, McCoy D. Access to antiretroviral treatment in Africa. *BMJ* 2004;**328**:241–2.
- 5 **McCoy D**, Chopra M, Loewenson R, *et al*. Expanding access to antiretroviral therapy in sub-Saharan Africa: avoiding the pitfalls and dangers, capitalizing on the opportunities. *Am J Public Health* 2005;**95**:18–22.
- 6 **Farmer P**, Leandre F, Mukherjee J, *et al*. Community-based treatment of advanced HIV disease: introducing DOT-HAART (directly observed therapy with highly active antiretroviral therapy). *Bull World Health Organ* 2001;**79**:1145–51.
- 7 **Hogan DR**, Salomon JA. Prevention and treatment of human immunodeficiency virus/acquired immunodeficiency syndrome in resource-limited settings. *Bull World Health Organ* 2005;**83**:135–43.
- 8 **Cleary SM**, McIntyre D, Boule AM. The cost-effectiveness of antiretroviral treatment in Khayelitsha, South Africa—a primary data analysis. *Cost Eff Resour Alloc* 2006;**4**:20.
- 9 **Braitstein P**, Brinkhof MW, Dabis F, *et al*. Mortality of HIV-1-infected patients in the first year of antiretroviral therapy: comparison between low-income and high-income countries. *Lancet* 2006;**367**:817–24.
- 10 **Stringer JS**, Zulu I, Levy J, *et al*. Rapid scale-up of antiretroviral therapy at primary care sites in Zambia: feasibility and early outcomes. *JAMA* 2006;**296**:782–93.
- 11 **Libamba E**, Makombe SD, Harries AD, *et al*. Malawi's contribution to "3 by 5": achievements and challenges. *Bull World Health Organ* 2007;**85**:156–60.
- 12 **Van Cutsem G**, Hilderbrand H, Coetzee D, *et al*. Clinical outcomes and emerging challenges after 5 years of ART in Khayelitsha, South Africa. *14th Conference on Retroviruses and Opportunistic Infections* 2007.
- 13 **Etard JF**, Ndiaye I, Thierry-Mieg M, *et al*. Mortality and causes of death in adults receiving highly active antiretroviral therapy in Senegal: a 7-year cohort study. *AIDS* 2006;**20**:1181–9.
- 14 **Coetzee D**, Hildebrand K, Boule A, *et al*. Outcomes after two years of providing antiretroviral treatment in Khayelitsha, South Africa. *AIDS* 2004;**18**:887–95.
- 15 **Lawn SD**, Myer L, Harling G, *et al*. Determinants of mortality and nondeath losses from an antiretroviral treatment service in South Africa: implications for program evaluation. *Clin Infect Dis* 2006;**43**:770–6.
- 16 **Martinson NA**, Karstaedt A, Venter WF, *et al*. Causes of death in hospitalized adults with a presumptive diagnosis of tuberculosis: an autopsy study. *AIDS* 2007;**21**:2043–50.
- 17 **Yu JK**, Chen SC, Wang KY, *et al*. True outcomes for patients on antiretroviral therapy who are "lost to follow-up" in Malawi. *Bull World Health Organ* 2007;**85**:550–4.
- 18 **Egger M**, Boule A, Schechter M, *et al*. Antiretroviral therapy in resource-poor settings: scaling up inequalities? *Int J Epidemiol* 2005;**34**:509–12.
- 19 **Fairall L**, Bachmann M, Louwagie G, *et al*. Effectiveness of antiretroviral treatment in a South African program: cohort study. *Arch Intern Med*. In press.
- 20 **Marins JR**, Jamal LF, Chen SY, *et al*. Dramatic improvement in survival among adult Brazilian AIDS patients. *AIDS* 2003;**17**:1675–82.
- 21 **Teixeira PR**, Vitoria MA, Barcarolo J. Antiretroviral treatment in resource-poor settings: the Brazilian experience. *AIDS*, 2004;**18**(Suppl 3), S5–S7.
- 22 **Laubsher R**, Bradshaw D, Bourne D, *et al*. Rapid surveillance of AIDS mortality in South Africa (437). *3rd South African AIDS Conference* 2007.
- 23 **Braitstein P**, Boule A, Nash D, *et al*. Gender and the use of potent antiretroviral treatment in resource-constrained settings: Findings from a multi-centre collaboration. *J Women's Health*. In press.
- 24 **Orrell C**, Harling G, Lawn S, *et al*. Conservation of first line antiretroviral treatment regimen where therapeutic options are limited. *Antivir Ther*. In press.
- 25 **Coetzee D**, Boule A, Hildebrand K, *et al*. Promoting adherence to antiretroviral therapy: the experience from a primary care setting in Khayelitsha, South Africa. *AIDS*, 2004;**18**(Suppl 3), S27–S31.
- 26 **Egger M**. Outcomes of ART in resource-limited and industrialized countries. *14th Conference on Retroviruses and Opportunistic Infections* 2007.
- 27 **Mills EJ**, Nachega JB, Buchan I, *et al*. Adherence to antiretroviral therapy in sub-Saharan Africa and North America: a meta-analysis. *JAMA* 2006;**296**:679–90.
- 28 **Bussmann H**, Novitsky V, Wester W, *et al*. HIV-1 subtype C drug-resistance background among ARV-naïve adults in Botswana. *Antivir Chem Chemother* 2005;**16**:103–15.
- 29 **Bessong PO**, Mphahlele J, Choge IA, *et al*. Resistance mutational analysis of HIV type 1 subtype C among rural South African drug-naïve patients prior to large-scale availability of antiretrovirals. *AIDS Res Hum Retrovir* 2006;**22**:1306–12.
- 30 **Kamya MR**, Mayanja-Kizza H, Kambugu A, *et al*. Predictors of long-term viral failure among Ugandan children and adults treated with antiretroviral therapy. *J Acquir Immune Defic Syndr* 2007;**46**:187–93.
- 31 **Kennedy C**, O'Reilly K, Medley A, *et al*. The impact of HIV treatment on risk behaviour in developing countries: a systematic review. *AIDS Care* 2007;**19**:707–20.
- 32 **Berkman A**. Confronting global AIDS: prevention and treatment. *Am J Public Health* 2001;**91**:1348–9.
- 33 **Médecins Sans Frontières**. Help wanted. *Confronting the health care worker crisis to expand access to HIV/AIDS treatment: MSF experience in southern Africa*. Johannesburg: Médecins Sans Frontières, 2007.
- 34 **Parkhurst JO**, Lush L. The political environment of HIV: lessons from a comparison of Uganda and South Africa. *Soc Sci Med* 2004;**59**:1913–24.
- 35 **Harries AD**, Schouten EJ, Libamba E. Scaling up antiretroviral treatment in resource-poor settings. *Lancet* 2006;**367**:1870–2.
- 36 **Boule A**, Orrell C, Kaplan R, *et al*. Substitutions due to antiretroviral toxicity or contraindication in the first 3 years of antiretroviral therapy in a large South African cohort. *Antivir Ther* 2007;**12**:753–60.
- 37 **Calmy A**, Hirschel B, Cooper DA, *et al*. Clinical update: adverse effects of antiretroviral therapy. *Lancet* 2007;**370**:12–4.
- 38 **Ford N**, Wilson D, Costa CG, *et al*. Sustaining access to antiretroviral therapy in the less-developed world: lessons from Brazil and Thailand. *AIDS*, 2007;**21**(Suppl 4), S21–S29.
- 39 **Havlic DV**, Hammer SM. Patents versus patients? Antiretroviral therapy in India. *N Engl J Med* 2005;**353**:749–51.
- 40 **Calmy A**, Ford N, Hirschel B, *et al*. HIV viral load monitoring in resource-limited regions: optional or necessary? *Clin Infect Dis* 2007;**44**:128–34.

Vaginal practices

Vaginal practices, microbicides and HIV: what do we need to know?

A Martin Hilber, M F Chersich, J H H M van de Wijgert, H Rees, M Temmerman

The goal of a safer vaginal environment could be reached by identifying harmful vaginal practices and an effective microbicide, thereby increasing options for HIV prevention

The global burden of HIV, its increasing feminisation, and chronic difficulties with development of options

for HIV prevention all argue for an intensified re-examination of factors influencing the efficiency of heterosexual

HIV transmission. This includes vaginal practices and products used by large numbers of women worldwide to tighten, dry, warm and clean their vagina. Women's efforts to change their genital environment can undermine each component of innate defences against pathogens.¹ In particular, vaginal practices have been linked with loss of lactobacilli and disruption of the vaginal epithelium.^{2–4} These practices may therefore be an important mediator in acquisition of STI, including HIV, or worsen pre-existing infections. Despite this, surprisingly little is known about the effects of specific vaginal practices on HIV transmission dynamics.

In past decades, both cross-sectional and longitudinal studies have found an association between intravaginal cleans-

Figure 1 Hypothesised causal pathway of vaginal practices and HIV. Results of the GSVP study indicated that practice types vary and are mitigated by practice characteristics, some of which have been detailed above. Intravaginal cleansing and insertion of substances are hypothesised as being the most potentially harmful and thus the most likely to be associated with increased risk of HIV acquisition.

ing and adverse reproductive outcomes, including pelvic inflammatory disease, ectopic pregnancy and bacterial vaginosis (BV).¹⁻⁶ BV could be an intermediary factor between vaginal practices and HIV infection (fig 1). Though much uncertainty remains about the pathophysiology of BV, its accompanying inflammatory milieu, characterised by pro-inflammatory cytokines and immune cell changes, is likely to enhance HIV transmission.⁷ Also, immunological changes with BV can stimulate HIV expression, raising HIV levels in the genital tract virus and likely the infectivity of women.⁷ Acquisition of HSV-2 may also be higher among women with BV.⁸

Evidence on whether intravaginal cleansing or insertion of substances increases risk for HIV acquisition is conflicting.¹ From the nonoxynol-9 trials, it is clear that some substances inserted into the vagina can cause epithelial disruption and are likely to facilitate HIV acquisition.⁴ It is possible that many commercial and traditional products have similar effects. Largely, these products remain to be identified. Limited prospective evidence is inconsistent, with two studies finding increased risk for HIV infection with the practice of using fingers to clean inside the vagina⁹⁻¹⁰ while a study in Zimbabwe and Uganda did not.¹¹ The latter study found an increased

risk for HIV acquisition in women who inserted substances to dry and tighten the vagina in preparation for sex. The authors hypothesised that these practices deserved greater research attention as they are potentially more abrasive and harmful.¹¹ Conversely, a few studies have suggested that some intravaginal cleansing practices may have beneficial health effects.¹²⁻¹³

Given the above biological rationale and limitations of available evidence, further prospective evaluation of the effects of vaginal practices is needed. With cross-sectional studies, it is difficult to distinguish the temporality of associations between HIV and specific practices.

Table 1 Classification of vaginal practices in the WHO Gender, Sexuality and Vaginal Practices Study

Practice	Definition
1. External washing	Cleaning of the external area around the vagina and genitalia using a product or substance with or without water normally using your hand. Products used vary from soap and water, to traditional and chemical detergent-like substances.
2. External application	Placing or rubbing various substances or products to the external genitalia—that is the labia, clitoris, vulva. Included is the “steaming” or “smoking” of the vagina, by sitting above a source of heat (fire, coals, hot rocks) on which water, herbs or oils are placed to create steam or smoke.
3. Anatomical modification (“cutting” and “pulling”)	Surgical procedures used for modifying the vagina, or restoration of the hymen; includes female genital circumcision, incision with insertion of substance into the lesion (scarification process, tattoos of the vulva or labia); excludes episiotomies or operations to repair a protruding uterus. In some countries, elongation or pulling of the labia minora is practised from early childhood. Sometimes, special creams or powders are applied during the pulling process.
4. Intravaginal cleansing (“washing”)	Internal cleansing or washing inside the vagina includes wiping the internal genitalia with fingers and other substances (eg, cotton, cloths, paper,) for the purpose of removing fluids. It also includes douching, which is the pressurised shooting or pumping of water or solution (including douching gel) into the vagina.
5. Intravaginal insertion	Pushing or placing something inside the vagina (including powders, creams, herbs, tablets, sticks, stones, leaves, cotton, paper, tampons, tissue, etc) regardless of the duration it is left inside.
6. Oral ingestion	Ingesting (drinking, swallowing) of substances perceived to affect the vagina and uterus. This includes the ingestion of substances/medicines to dry or lubricate the vagina. Substances may be dissolved in water or other liquids.

The WHO GSVP Study Group conducted extensive qualitative and prevalence research on vaginal practices in Asia and Africa. The GSVP Classifications summarised above are a result of this work. The consolidated classifications cover all practices that have been documented in the scientific literature to date on vaginal and genital practices and can be used for future studies.

HIV infection itself can lead to increased prevalence of bacterial vaginosis or exacerbate other reproductive-tract infections, resulting in increased vaginal discharge, which in turn motivates vaginal practices. Reverse causation may thus be a reasonable explanation for cross-sectional associations.¹⁰

Standardised and more detailed measures of vaginal practices would also improve the quality of this research. In studies of this topic to date, categorisation of practices varies considerably, hindering cross-study comparison and assessment of effects of individual practices. Recently, however, a vaginal practices classification and measurement framework has been developed in the WHO Multi-Country Study on Gender, Sexuality and Vaginal Practices (GSVP). Six categories of vaginal practices have been defined (table 1). This framework emanated from extensive qualitative investigation and subsequent testing in a GSVP household survey of vaginal practices in four countries.¹⁴ In addition to using a standardised classification system, the ability of future studies to detect effects of these practices would be improved by collecting detailed data on products used, frequency of practices, timing of practices with coitus and whether these practices vary with nature of relationship and condom use. Social norms about lubrication during sex and corresponding vaginal practices have important implications for acceptability of microbicides that alter vaginal lubrication. In microbicide trials, it is important that the candidate and placebo product have similar effects on vaginal lubrication. This would avoid eliciting differential vaginal practices in each group, some of which potentially increase susceptibility to HIV. Also, women who remove vaginal lubrication or cleanse the vagina around the time of sex may unintentionally remove or dilute the microbicide product. Of perhaps greater concern, substances inserted in the vagina may chemically interact with the microbicide, with potentially harmful by-products of such reactions. With the multitude of products used in vaginal practices, a myriad of chemical reactions between such products and microbicides is possible. As an investigator at a microbicides trial site in South Africa noted: "Cultural rituals pose challenges to microbicides clinical trials as women combine the gel with other hoped for barriers to HIV including Dettol (a disinfectant), herbs, snuff, and lemon juice".¹⁵

An approach needs to be developed to investigating potential reactions between microbicides and products women insert vaginally, analogous to the need for

investigating potential interactions between ingested or intravenous drugs. Detailed, repeated documentation of vaginal practices in microbicide trials would facilitate investigation of such effects. In sum, potential effects of vaginal practices on microbicide acceptability, efficacy and safety warrant further consideration.

Most clearly, vaginal practices and their motivations reveal much about the gendered nature of the HIV epidemic, and its underlying drivers. Particularly in resource-constrained settings, sex may be a primary means of achieving economic security, either directly through sex work or through maintaining economically essential relations with sexual partners or husbands. Optimising sexual pleasure for men by, for example, responding to societal norms about vaginal lubrication or tightness increases a woman's ability to leverage such economic sustenance. A large variety of products are used in each vaginal practice, the sale of which often supports a sizeable informal and formal industry. Women may spend a considerable portion of their limited resources on such products, hoping to obtain a return from their investment by achieving the vaginal state men desire. In settings with unequal gender power and economic disparities, women will continue to have limited ability to negotiate protected sex and few alternatives to adopting practices which explicitly aim to satisfy men's sexual desires. Other reasons have also been reported for engaging in vaginal practices, and these vary widely in different settings, including genital hygiene, pregnancy prevention, and prevention or self-treatment of vaginal infections.¹⁴⁻¹⁶ It is important to note that some evidence suggests women's vaginal practices are modifiable. Though studies have reported women are reluctant to alter their practices, in a randomised trial, women who received a counselling intervention were 1.6 times more likely to stop douching compared with women in the control group.¹⁷

Conceptually, in recent years the role of the cervix in HIV infection has gained prominence, with a diminished emphasis on the vagina as an entry site. Investigation of cervical barriers is consistent with these trends, and supported by animal studies and biological plausibility.¹⁸ However, recent data showing poor efficacy of the diaphragm in preventing HIV acquisition¹⁹ suggest the cervical-centric approach to HIV prevention among women is now being questioned, and that vaginal health is also paramount. Documenting why women engage in vaginal practices, the mechanisms by which such practices might

increase HIV acquisition and how modifiable these practices are, will be essential for the development of interventions to improve vaginal health. While efforts continue to reduce women's underlying gendered vulnerability to HIV, the goal of a safer vaginal environment could be reached by identifying harmful vaginal practices and an effective microbicide, thereby increasing options for HIV prevention.

Sex Transm Infect 2007;**83**:505–508.

doi: 10.1136/sti.2007.028597

.....

Authors' affiliations

A Martin Hilber, Institute of Social and Preventive Medicine, University of Bern, Switzerland

M F Chersich, International Centre for Reproductive Health, Kenya

J H H M van de Wijgert, Academic Medical Center, Center for Poverty-Related Communicable Diseases, Amsterdam, The Netherlands

H Rees, Reproductive Health and HIV Research Unit, University of Witwatersrand, South Africa

M Temmerman, Department of Obstetrics and Gynaecology, Ghent University, Belgium

Correspondence to: Ms A Martin Hilber, Institute of Social and Preventive Medicine, University of Bern, Finkenhubelweg 11, CH-3012 Bern, Switzerland; amartinhilber@ispm.unibe.ch

Accepted 2 October 2007

Competing interests: None declared.

REFERENCES

- 1 Myer L, Kuhn L, Stein ZA, *et al.* Intravaginal practices, bacterial vaginosis, and women's susceptibility to HIV infection: epidemiological evidence and biological mechanisms. *Lancet Infect Dis* 2005;**5**:786–94.
- 2 Kilmarx PH, Limpakarnjanarat K, Supawitkul S, *et al.* Mucosal disruption due to use of a widely-distributed commercial vaginal product: potential to facilitate HIV transmission. *AIDS* 1998;**12**:767–73.
- 3 McClelland RS, Ndinya-Achola JO, Baeten JM. Is vaginal washing associated with increased risk of HIV-1 acquisition? *AIDS* 2006;**20**:1347–48.
- 4 Wilkinson D, Ramjee G, Tholandi M, *et al.* Nonoxonyl-9 for preventing vaginal acquisition of HIV infection by women from men. In: *Cochrane Database Syst Rev* 2002:CD003936.
- 5 Hassan WM, Lavreys L, Chohan V, *et al.* Associations between intravaginal practices and bacterial vaginosis in Kenyan female sex workers without symptoms of vaginal infections. *Sex Transm Dis* 2007;**34**:384–8.
- 6 Simpson T, Merchant J, Grimley DM, *et al.* Vaginal douching among adolescent and young women: more challenges than progress. *J Pediatr Adolesc Gynecol* 2004;**17**:249–55.
- 7 St John E, Mares D, Spear GT. Bacterial vaginosis and host immunity. *Curr HIV/AIDS Rep* 2007;**4**:22–8.
- 8 Chernes TL, Meyn LA, Krohn MA, *et al.* Association between acquisition of herpes simplex virus type 2 in women and bacterial vaginosis. *Clin Infect Dis* 2003;**37**:319–25.
- 9 McClelland RS, Ndinya-Achola JO, Baeten JM. Re: distinguishing the temporal association between women's intravaginal practices and risk of human immunodeficiency virus infection: a prospective study of South African women.

- Am J Epidemiol*, 2007;**165**:474–5; author reply 475–6.
- 10 **Myer L**, Denny L, de Souza M, *et al.* Distinguishing the temporal association between women's intravaginal practices and risk of human immunodeficiency virus infection: a prospective study of South African women. *Am J Epidemiol* 2006;**163**:552–60.
 - 11 **van de Wijgert J**, Morrison C, Cornelisse P, *et al.* Bacterial vaginosis and vaginal yeast, but not vaginal cleansing, increase HIV-1 acquisition in African women. *JAIDS*. In press.
 - 12 **Gresenguet G**, Kreiss JK, Chapko MK, *et al.* HIV infection and vaginal douching in central Africa. *AIDS* 1997;**11**:101–6.
 - 13 **La Ruche G**, Messou N, Ali-Napo L, *et al.* Vaginal douching: association with lower genital tract infections in African pregnant women. *Sex Transm Dis* 1999;**26**:191–6.
 - 14 **WHO**. *Multi-country study on gender, sexuality and vaginal practices*, 2007. Available from: <http://www.who.int/reproductive-health/fgm/harmfulpractices.htm> (accessed 24 Oct 2007).
 - 15 **Smith C**. The closures of trials: perspectives from the ground. *The Microbicide Quarterly* 2007;**5**:14–19. http://www.microbicide.org/microbicideinfo/reference/TMQ_Jan-Mar2007.FINAL.pdf (accessed 30 Sep 2007).
 - 16 **Braunstein S**, van de Wijgert J. Preferences and practices related to vaginal lubrication: implications for microbicide acceptability and clinical testing. *J Womens Health (Larchmt)* 2005;**14**:424–33.
 - 17 **Grimley DM**, Oh MK, Desmond RA, *et al.* An intervention to reduce vaginal douching among adolescent and young adult women: a randomized, controlled trial. *Sex Transm Dis* 2005;**32**:752–8.
 - 18 **Moench TR**, Chipato T, Padian NS. Preventing disease by protecting the cervix: the unexplored promise of internal vaginal barrier devices. *AIDS* 2001;**15**:1595–602.
 - 19 **Padian NS**, van der Straten A, Ramjee G, *et al.* Diaphragm and lubricant gel for prevention of HIV acquisition in southern African women: a randomised controlled trial. *Lancet* 2007;**370**:251–61.

Global strategy

WHO global strategy for the prevention and control of sexually transmitted infections: time for action

D A Lewis, A S Latif, F Ndowa

STI health professionals should use every opportunity to influence those able to initiate change to improve global STI control and prevention activities

Worldwide, sexually transmitted infections (STIs) continue to be a major cause of morbidity and mortality. Global estimates suggest that more than 340 million new cases of syphilis, gonorrhoea, chlamydial infection and trichomoniasis occurred throughout the world in 1999.¹ Congenital syphilis, prevention of which is relatively easy and cost-effective, may still be responsible for as many as 14% of neonatal deaths.² Up to 10% of those women who are untreated, or inadequately treated, for chlamydial and gonococcal infections may become infertile as a consequence.² On a global scale, up to 4000 newborn babies each year may become blind because of gonococcal and chlamydial ophthalmia neonatorum.²

There is evidence that STIs may enhance both the transmission and acquisition of HIV infection, and that improved control of STIs may slow down HIV transmission.³ The prevention and control of STIs is not an easy task. Epidemiological patterns of STIs vary geographically and are influenced by cultural, political, economical and social forces. Many affected by STIs are in marginalised vulnerable groups. The asymptomatic nature of some STIs remains a challenge to healthcare providers in areas of the world where laboratory screening tests are unaffordable.

The World Health Organization's "Global Strategy for the prevention and control of sexually transmitted infections: 2006–2015" was presented to Member States at the 59th World Health Assembly in May 2006. The World Health Assembly endorsed the Global Strategy and urged member states to adopt and draw on it in order to ensure that national efforts to achieve the United Nations' Millennium Development Goals include plans and actions for the prevention and control of STIs.²

The objectives of the Global Strategy are (a) to increase the commitment of both national governments and international development partners to STI prevention and control activities, (b) to promote both reallocation of existing resources and mobilisation of funds to support these activities, (c) to ensure that initiatives, policies and laws related to in-country provision of STI care are non-stigmatising and gender-sensitive, and (d) to facilitate better networking between all relevant partners and institutions in order to scale up and sustain interventions for STI prevention and control.

The Global Strategy has technical and advocacy components.² The technical component includes promotion of healthy sexual behaviour, delivery of good-quality STI care (Box 1), reliable supply of

essential commodities and medicines, strengthening of support components and review of policies, laws and regulations regarding STI control. The advocacy component addresses the need for an effective STI advocacy campaign to raise awareness and mobilise resources at the national and international level.

WHO, with other international partners and governments, is in the process of elaborating an Action Plan for the implementation of the Global Strategy structured around the technical and advocacy components of the Strategy. It sets out priorities (Box 2), actions, a time frame and performance indicators, for implementing the Strategy at global and regional levels. A draft of the Action Plan was reviewed at a stakeholders' consultation in Geneva in June this year. This meeting brought together government representatives, bilateral donor agencies, representatives from Member States in the WHO Regions, laboratory experts and other scientists. Each WHO Region will select countries of focus where scaling up the priority interventions may be supported more intensively.

In terms of support components, the lack of effective STI surveillance systems in many countries presents a major challenge. Surveillance of clinical syndromes is easier to establish in resource-poor countries than more costly and technically demanding microbiological surveillance systems. The success of STI syndromic management does, however, rely on up-to-date local knowledge of the pathological agents causing the syndromes (aetiological surveillance) and the antimicrobial susceptibility profile of bacteria such as *Neisseria gonorrhoeae*. In order to undertake microbiological surveillance studies in resource poor areas of the world, investment will be required to strengthen regional laboratories. As part of surveillance initiatives, it will be necessary to define minimum essential sets of STI syndromes and standardised laboratory investigations, in order to allow comparisons between countries and regions.