Published in final edited form as:

Schizophr Res. 2008 October ; 105(1-3): 197–200. doi:10.1016/j.schres.2008.06.017.

Twin pregnancy and the risk of schizophrenia

K Kleinhaus^a, S Harlap^{a,b}, MC Perrin^a, O Manor^c, R Calderon-Margalit^c, Y Friedlander^c, and D Malaspina^a

^aDepartment of Psychiatry, New York University School of Medicine, New York, NY, 10016

^bDepartment of Epidemiology, Mailman School of Public Health, Columbia University, New York, NY, 10032

^cHebrew University-Hadassah Braun School of Public Health and Social Medicine, Jerusalem, Israel

Abstract

Background—Twins are exposed to intrauterine environments that differ significantly from those of singletons. These diverse environments might alter the risk for schizophrenia in twins and make it difficult to generalize from findings in twins when studying the risk of schizophrenia in the general population. Previous studies report contradictory findings on the risk for schizophrenia in twins.

Methods—We studied the incidence of schizophrenia spectrum disorders, ascertained from Israel's national Psychiatric Registry, in a cohort of 2,124 twins and 87,955 singletons. These offspring were followed from their birth in 1964–76 in the Jerusalem Perinatal study. Cox proportional hazards methods were used to compare outcomes over 28–41 years, adjusting for ages of parents.

Results—Twins showed a relative risk [RR] of .84 relative to singletons, with a 95% confidence interval [CI] of (.51–1.4). RRs and CIs for males and females were .68 [.34–1.4] and 1.1 [.55–2.2] respectively. Twins in male-male, female-female or opposite-sex sets showed no significant variation in RRs; furthermore, first- or second-born twins did not differ significantly from each other. Siblings of twins had the same risk of schizophrenia as siblings of singletons.

Conclusion—Twins have the same risk for schizophrenia as the general population.

Keywords

schizophrenia; twins; paternal age; maternal age

1.1 Introduction

Comparisons of concordance of conditions in monozygotic versus dizygotic twins have been the basis for estimates of heritability and gene-environment interactions in schizophrenia¹. The intrauterine environment of twins, however, differs significantly from that of singletons in ways that might be related to the future risk of disease. Twins are exposed to higher levels of steroid hormones such as estrogen², are more likely to be born prematurely, and to weigh less at birth even when born at term³. Twins are at increased risk for breech presentation, hypoxia and birth injuries⁴. These obstetric complications have been associated with risk of

Correspondence: Dr. K. Kleinhaus, Department of Psychiatry, NYU Medical Center, 550 First Avenue, Millhauser Labs, Room HN323, New York, NY 10016, Email: Kleink03@nyumc.org, Phone: 212-263-7626.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

schizophrenia⁵. Twins often have different postnatal experiences; parents must provide more resources, and twins may be treated differently in social situations.

The current literature disagrees on whether the incidence of schizophrenia in twins differs from that in the general population, and whether this incidence varies across sets of different zygosity. One study found a 28% increase in the rate of first admissions for schizophrenia in twins compared to the general population, as well as an increased rate in twins for first admission for any psychiatric disorder⁶. Another study found no difference in the incidence of psychotic disorders in twins⁷. Regarding siblings of twins, a recent report suggested that the rate of schizophrenia in siblings of dizygotic twins was 35% greater than the rate in siblings of singletons, while the rate in siblings of monozygotic twins was not increased. The authors concluded that their results accorded with a hypothetical linkage between genes influencing the rate of dizygotic twinning and those predisposing to schizophrenia⁸. We used existing data from a prospective cohort study to test whether the incidence of schizophrenia is higher in twins than in singletons.

2.1 Methods

We used data from the Jerusalem Perinatal Study, a population-based cohort derived from all 92,408 births in 1964–76 to mothers resident in Western Jerusalem. The cohort includes linkages within nuclear families and a 29–41 year follow-up. Core information from the notification of birth was supplemented with other data from multiple sources. The methods and characteristics of the population have been described in previous publications⁹, as have the characteristics of fathers of different ages¹⁰. The cohort was linked with Israel's population registry and National Psychiatric Registry; the latter contains a record of all admissions to psychiatric wards and day-facilities and includes the dates of admission and discharge and the discharge diagnosis for each episode, assigned by a board-certified psychiatrist. The diagnoses are coded with the International Statistical Classification of Diseases 10th Revision [ICD-10]. The diagnoses of psychotic disorders have been validated¹¹.

2.2 Data Analysis

SAS 9.1 (SAS Institute Inc, Cary, North Carolina) was used to analyze the data. We assigned a diagnosis of schizophrenia to any person with at least one hospital admission for which the discharge diagnosis was coded with ICD-10 code F20-29, hereafter termed "schizophrenia". The date of onset was taken as the first episode in the psychiatric registry, regardless of the diagnosis at that time. After comparing demographic and other characteristics of singleton and multiple deliveries, we developed Cox proportional hazards models to compare the incidence of schizophrenia in twins versus singletons, running time in days from date of birth until date of diagnosis of schizophrenia or death; survivors were censored on December 31, 2004. Schizophrenia was coded as 1 (if present) or 0. Maternal and paternal ages were included in the models because both have been related to twinning¹² and schizophrenia¹³. As has previously been done in this cohort¹³, paternal age was treated as a continuous variable, expressed as deviations from the mean age (30) with unknowns (0.8%) assigned to the mean; while maternal age was modeled using two dichotomies 30–34, 35+, versus <30. Other variables considered were mothers' and fathers' places of birth (Israel, other West Asia, North Africa, Europe), social class (based on occupation of father) sex, birth order (1,2,3–4,5+), year of birth, and education of parents (0–8 years, 9+ years). None of these other variables were included in the final models, as they were not related to both schizophrenia and twin status at $p < .05$ or did not alter the crude hazard ratio (HR) for twins by at least 10%.

3.1 Results

Of the 92,408 offspring in the cohort (91,479 born alive), 90,079 (98%) were traced in Israel's Population Registry. Linkage with Israel's Psychiatric Registry detected 860 cases of schizophrenia, of which 16 were twin offspring and 844 were singletons. Of the twins with schizophrenia, all were in discordant sets and both twins in each set were born alive.

In proportional hazards analysis the unadjusted incidence of schizophrenia was higher in males than in females (RR=1.6, 95% confidence limits=1.4–1.8), offspring of older fathers (1.03, 1.02–1.04 per 10 years increase in father's age) and older mothers aged 30–34 (1.3, 1.1–1.6) or 35+ (1.6, 1.3–1.9) versus offspring of mothers aged <30 (reference group). Table 1 shows that the relative risk of schizophrenia in twins was the same as in singletons, and that adjustment for maternal or paternal age did not change the risk. The risk for opposite-sex twins compared to singletons was about half of that in same-sex twins, although the confidence limits included one. The adjusted RR was .46 with a 95% confidence interval of .15–1.4. When the risk for twins from male-male sets was compared to the risk for male singletons, the RR was 0.86 (0.41–1.8). When female singletons were designated as the reference group, the RR for schizophrenia in twins from female-female sets was 1.2, 0.55–2.7. These relative risks are not significantly different from those in Table 1 where the entire cohort was used as the reference group. The birth order of co-twins did not affect the risk for schizophrenia, nor did discordant birth weights (data not shown). There were 2,658 traced singleton siblings of twins, and these also showed no significantly changed risk of schizophrenia as compared to the rest of the population (data not shown).

4.1 Discussion

In this cohort the incidence of schizophrenia in twins did not differ from that in singletons, nor did it vary across types of twin sets, by the sex of the twins, by birth order within the set, or with discordance in birth weights. These results corroborate those reported by Kendler et al in their retrospective study of data from Sweden's registry of twins⁷. In contrast, Klaning et al's 1996 analysis of data from Denmark, using the same retrospective methods as the Swedish study, found a 28% increase in risk for schizophrenia in twins as compared to singletons, a higher rate ratio in female twins than in male twins, and in twins from opposite-sex pairs than in twins from same-sex pairs⁶. We also observed a higher relative risk in female twins than in male twins, although with overlapping confidence intervals the results were likely to be due to chance. The relative risk for opposite-sex twins in our study was roughly half of the risk in male-male or female-female sets, although here too the confidence intervals overlapped.

The Danish group proposed a possible association between the genetic tendency to conceive dizygotic twins and a genetic vulnerability for schizophrenia {Klaning, 2002 #38}. Our results lend no support for this hypothesis, as the siblings of twins had the same incidence of schizophrenia as the siblings of singletons. It is not evident why the findings in Denmark diverge from our study and the study from Sweden.

This analysis of prospectively collected data shows that the risk for schizophrenia in singletons and twins are comparable. The findings support assumptions made by the field when estimating risk factors for schizophrenia using data on twins, and also help resolve the discrepancy in the literature.

Acknowledgements

Supported by NIH Grants: # 2R01 CA080971 (S.Harlap) and # 1R01 MH059114 (D. Malaspina) and the National Alliance for Research on Schizophrenia and Depression (DM, SH, MP).

References

1. Sullivan PF, Kendler KS, Neale MC. Schizophrenia as a complex trait: evidence from a meta-analysis of twin studies. *Arch Gen Psychiatry* 2003 Dec;60(12):1187–1192. [PubMed: 14662550]
2. Swerdlow AJ, De Stavola BL, Swanwick MA, Maconochie NE. Risks of breast and testicular cancers in young adult twins in England and Wales: evidence on prenatal and genetic aetiology. *Lancet* 1997 Dec 13;350(9093):1723–1728. [PubMed: 9413462]
3. Herruzo AJ, Martinez L, Biel E, Robles R, Rosales MA, Miranda JA. Perinatal morbidity and mortality in twin pregnancies. *Int J Gynaecol Obstet* 1991 Sep;36(1):17–22. [PubMed: 1683296]
4. Bjelic-Radisic V, Pristauz G, Haas J, et al. Neonatal outcome of second twins depending on presentation and mode of delivery. *Twin Res Hum Genet* 2007 Jun;10(3):521–527. [PubMed: 17564511]
5. Byrne M, Agerbo E, Bennedsen B, Eaton WW, Mortensen PB. Obstetric conditions and risk of first admission with schizophrenia: a Danish national register based study. *Schizophr Res* 2007 Dec;97(1–3):51–59. [PubMed: 17764905]
6. Klaning U, Mortensen PB, Kyvik KO. Increased occurrence of schizophrenia and other psychiatric illnesses among twins. *Br J Psychiatry* 1996 Jun;168(6):688–692. [PubMed: 8773810]
7. Kendler KS, Pedersen NL, Farahmand BY, Persson PG. The treated incidence of psychotic and affective illness in twins compared with population expectation: a study in the Swedish Twin and Psychiatric Registries. *Psychol Med* 1996 Nov;26(6):1135–1144. [PubMed: 8931159]
8. Klaning U, Pedersen CB, Mortensen PB, Kyvik KO, Skytthe A. A possible association between the genetic predisposition for dizygotic twinning and schizophrenia. *Schizophr Res* 2002 Nov 1;58(1):31–35. [PubMed: 12363387]
9. Harlap S, Davies AM, Deutsch L, et al. The Jerusalem Perinatal Study cohort, 1964–2005: methods and a review of the main results. *Paediatr Perinat Epidemiol* 2007 May;21(3):256–273. [PubMed: 17439536]
10. Harlap S, Paltiel O, Deutsch L, et al. Paternal age and preeclampsia. *Epidemiology* 2002 Nov;13(6):660–667. [PubMed: 12410007]
11. Weiser M, Kanyas K, Malaspina D, et al. Sensitivity of ICD-10 diagnosis of psychotic disorders in the Israeli National Hospitalization Registry compared with RDC diagnoses based on SADS-L. *Compr Psychiatry* 2005 Jan-Feb;46(1):38–42. [PubMed: 15714193]
12. Tough S, Benzie K, Fraser-Lee N, Newburn-Cook C. Factors influencing childbearing decisions and knowledge of perinatal risks among Canadian men and women. *Matern Child Health J* 2007 Mar;11(2):189–198. [PubMed: 17237994]
13. Malaspina D, Harlap S, Fennig S, et al. Advancing paternal age and the risk of schizophrenia. *Arch Gen Psychiatry* 2001 Apr;58(4):361–367. [PubMed: 11296097]

Table 1
Number of people with and without schizophrenia, relative risk (RR) and confidence limits (CL)

		Schizophrenia		Unadjusted Analysis		Analysis adjusted for mothers' and fathers' ages		
		+	-	RR	95% CL	RR	95% CL	P
Males	Singletons	844	87,111	1	---	1	---	---
	Twins	16	2,108	.87	(.53, 1.4)	.84	(.51, 1.4)	.5
Females	Singletons	528	44,768	1	---	1	---	---
	Twins	8	1,084	.70	(.35, 1.4)	.68	(.34, 1.4)	.3
Twin Sets	Singletons	316	42,332	1	---	1	---	---
	Twins	8	1,024	1.1	(.57, 2.3)	1.1	(.55, 2.2)	.8
Twin Sets	Singletons	844	87,111	1	---	1	---	---
	Male-Male	7	764	1.1	(.51, 2.3)	1.1	(.5, 2.2)	.9
	Female-Female	6	704	.96	(.43, 2.1)	.94	(.42, 2.1)	.9
	Opposite-sex	3	697	.5	(.16, 1.5)	.46	(.15, 1.4)	.2