Published in final edited form as:

J Neuropsychol. 2008 September ; 2(Pt 2): 509–514. doi:10.1348/174866407X269767.

Serial reaction time performance following right parietal lobe damage

M.E Berryhill, Y Mazuz, and I.R Olson

Abstract

The serial reaction time task (SRT) is used to assess implicit sequence learning. Neuroimaging studies implicate parietal involvement, however, the necessity of this area is unclear. We tested six unilateral right parietal patients and compared their performance to matched controls. Both groups showed similar levels of learning and explicit awareness. Two patients with the largest lesions extending into either frontal or cerebellar regions showed no learning. These data suggest implicit sequence learning can occur despite damage to the right parietal lobe.

Introduction

In the serial response time (SRT) task, subjects make speeded key press responses to visual cues. Unbeknownst to subjects, a specific sequence is embedded and repeated. Reaction times typically become faster with repetition, only to slow down when a novel stimulus order is introduced. Because many subjects fail to notice the sequence, performance on the SRT is often considered a measure of implicit learning (Nissen & Bullemer, 1987).

The SRT task is frequently used to investigate how the brain implicitly learns sequences. Neuroimaging studies using the SRT paradigm consistently report either bilateral parietal activations (Keele, Ivry, Mayr, Hazeltine, & Heuer, 2003; Poldrack, Sabb, Foerde, Tom, Asarnow, Bookheimer, & Knowlton, 2005; van der Graaf, Maguire, Leenders, & de Jong, 2006) or right parietal activations (Bischoff-Grethe, Martin, Mao & Berns, 2001; Olson, Rao, Moore, Wang, Detre, & Aguirre, 2006). Prior neuropsychological studies of performance on the SRT task have not examined the effects of damage to this region, however. Whether or not it plays an essential role in sequence learning is unknown. To evaluate this, we administered the SRT task to a group of patients with right parietal lobe damage.

Methods

Participants

The patient population consisted of 6 patients with unilateral right parietal lobe damage, aged 40–75 years (mean 58.6 ± 13.9 , 1 male), with an average of 12.7 years of education. All patients were right handed, high functioning, free from neglect and visual field cuts. Neglect was assessed using several tasks: Albert's line cancellation, clock face drawing, greyscales and stimulus extinction. Patient lesions can be viewed in Figure 1; patient descriptions can be found in Table 1.

Patients' performance was compared to that of 9 age ($M = 55.7$ years, range 43–71 years, 4 males) and education ($M = 14.8$) matched controls. Controls were right handed and free of neurological disorders. All subjects were paid \$15 per hour for their participation. Written consent was given according to an Institutional Review Board approval from the University of Pennsylvania.

Task

The stimuli consisted of four horizontally-arrayed squares on a medium gray background. Each square became active when filled with its own distinct color (green, blue, red, and yellow, from left to right). Subjects were instructed to respond by pressing the spatially corresponding key as quickly and as accurately as possible. A new square activated immediately following a key press. Subjects used the index and middle fingers of each hand on the *D*, *F*, *J*, and *K* keys of a computer keyboard. One patient (592) used the four fingers of her right hand on the *F*, *G*, *H*, and *J* keys due to hemiparesis of the left hand.

The invariant sequence was 8-items long consisting of two key presses from each finger. This sequence length was chosen in order to be slightly easier for the patients than commonly-used longer sequences, and to show learning more quickly. The 8-item invariant sequence repeated 80 times and was followed by a test phase, which controlled for earlier reaction time improvements attributable to general features of the task by presenting a pseudorandom order of stimuli. The test phase consisted of 120 pseudorandomly ordered button presses. The final time bin was pseudorandom to ensure that no three-step sub-sequence from the repeated sequence reoccurred. The experiment was performed without pause and lasted approximately 15 minutes.

After completing the task, we tested explicit awareness by asking subjects if they noticed the embedded sequence. Subjects were then asked to type the remembered pattern into the computer.

Analysis

The dependent measure was key press reaction time (RT). Incorrect trials and outliers were excluded on a per-subject basis ($RT > \text{mean} + 2 \times \text{standard deviation}$). Trials were binned into five epochs: 1st – 4th consisted of the repeated-sequence trials (20 repetitions, or 160 key presses), and the 5th consisted of the 120 pseudorandom stimuli. Mean RTs were calculated and analyzed using a repeated-measures ANOVA with SPSS 11.0.

Results

Accuracy was high in both groups (controls, $M = 0.97 \pm 0.02$; patients, $M = 0.95 \pm 0.04$). Learning was compared using a two-factor repeated-measures ANOVA on group (patient, control) and epoch (1st – 5th); results are depicted in Figure 2. The patients were on average, 348 ms slower than the controls ($F_{1, 13} = 8.50$, $P < 0.01$), a finding commonly observed in lesions populations. Of interest, both groups showed RT improvement over the training epochs ($F_{4, 52} = 9.10$, $P < 0.001$) and there was no significant interaction between group and epoch ($F_{4, 52} < 1$, n.s.).

To assess implicit learning in more detail, we examined the change in RT between the 4th epoch and the 5th epoch. This measure captures the change in performance when switching from a predictable to an unpredictable pattern. This t-test was significant for control ($t_8 = 4.37$, $P = .002$) and patient ($t_5 = 2.94$, $P = .03$) groups.

Last, a comparison of the difference scores (epoch 5 - epoch 4) for each group found no difference in the magnitude of learning between patients and controls ($t_{13} = 0.04$, $P = 0.97$).

Looking at individual performance, only one patient, patient (#560), showed fairly flat performance over the four sequence epochs. However he evidenced learning in two other ways: his performance slowed during the pseudorandom epoch, and he noticed the presence of a pattern during the test of conscious memory (see below) that followed the training session.

It is possible that with a larger sample of patients, we would have observed relatively poorer SRT learning in the patient group. To estimate our chances of finding this, we conducted a power analysis suggested that approximately 70 patients would be needed in order to detect a small effect size ($= .1$) with high power ($\beta = .8$). This finding indicates that there may be small differences in the SRT learning abilities of patients and controls that become evident when a large population is tested, but that such differences are not observed at the individual or small-group level.

Test of Conscious Memory

Two of the 9 controls (22%) and 2 of 6 patients (33%) claimed to notice a repeating pattern during the task. One control, unable to reproduce the pattern by typing, verbally described the first five steps in terms of the colors that appeared. One patient successfully re-typed the first five key presses of the pattern. No other participant produced more than three steps of the learned sequence. These findings suggest that (a) patients and controls did not differ in access to explicit awareness; and (b) for most subjects, learning was implicit.

Discussion

Sequence learning deficits on the SRT task have been observed in several patient populations. Not surprisingly, damage to motor regions such as the cerebellum and basal ganglia cause sequence learning deficits (Doyon et al., 1998; Gomez-Belderrain, Garcia-Monco, Rubio, & Pascual-Leone, 1998; Molinari et al., 1997; Willingham & Koroshetz, 1993). Further analyses have indicated that the cerebellum and basal ganglia may be critical for motor performance and/or the formation of higher order sequences, rather than for the formation of learning per se (Seidler et al., 2002).

Prefrontal damage, to the right, but not left hemisphere, causes impaired sequence learning (Gomez Beldarrain, Grafman, Ruiz de Velasco, Pascual-Leone, & Garcia-Monco, 2004; Keele et al., 2003). Accordingly, TMS to prefrontal cortex disrupts sequence learning for spatial, but not non-spatial sequences (Robertson, Tormos, Maeda, & Pascual-Leone, 2001). In contrast, bilateral medial temporal lobe damage does not dramatically affect sequence learning (Curran, 1997), and as seen here, neither does right parietal lobe damage.

Although damage to the right parietal lobe does not affect sequence learning, it is possible that damage to the left parietal lobe will impair sequence learning. Left parietal lobe activity is associated with motor attention and intention (Rushworth, Krams & Passingham, 2001). The underlying computations required for monitoring and planning actions may support those required for implicitly learning action sequences.

Acknowledgements

We would like to thank Anjan Chatterjee and Marianna Stark for allowing us to use their patient database, and David Drowos for assistance with testing controls. This research was supported by an NIH grant to I. Olson (RO1 MH071615-01).

References

- Bischoff-Grethe A, Martin M, Mao H, Berns GS. The context of uncertainty modulates the subcortical response to predictability. *Journal of Cognitive Neuroscience* 2001;13:986–993. [PubMed: 11595100]
- Curran T. Higher order associative learning in amnesia: Evidence from the serial reaction time task. *Journal of Cognitive Neuroscience* 1997;9:522–533.
- Doyon J, Laforce RJ, Bouchard G, Gaudreau D, Roy J, Poirier M, et al. Role of the striatum, cerebellum, and frontal lobes in the automatization of a repeated visuomotor sequence of movements. *Neuropsychologia* 1998;36:625–641. [PubMed: 9723934]

- Gomez-Belderrain M, Garcia-Monco JC, Rubio B, Pascual-Leone A. Effect of focal cerebellar lesions on procedural learning on the serial reaction time task. *Experimental Brain Research* 1998;120:25–30.
- Keele S, Ivry R, Mayr U, Hazeltine E, Heuer H. The cognitive and neural architecture of sequence representation. *Psychological review* 2003;110(2):316–339. [PubMed: 12747526]
- Molinari M, Leggio MG, Solida A, Ciorra R, Misciagna S, Silveri M, et al. Cerebellum and procedural learning: Evidence from focal cerebellar lesions. *Brain and Cognition* 1997;120:1753–1762.
- Nissen M, Bullemer P. Attentional requirements of learning: Evidence from performance measures. *Cognitive Psychology* 1987;19:1–32.
- Olson IR, Rao H, Moore KS, Wang J, Detre JA, Aguirre GK. Using perfusion fMRI to measure continuous changes in neural activity with learning. *Brain & Cognition* 2006;60:262–271. [PubMed: 16423439]
- Poldrack RA, Sabb FW, Foerdes K, Tom SM, Asarnow RF, Bookheimer SY, Knowlton BJ. The neural correlates of motor skill automaticity. *Journal of Neuroscience* 2005;25:5356–5364. [PubMed: 15930384]
- Robertson EM, Tormos JM, Maeda F, Pascual-Leone A. The role of the dorsolateral prefrontal cortex during sequence learning is specific for spatial information. *Cerebral Cortex* 2001;11:628–635. [PubMed: 11415965]
- Rushworth MFS, Krams M, Passingham RE. The attentional role of the left parietal cortex: the distinct lateralization and localization of motor attention in the human brain. *Journal of Cognitive Neuroscience* 2001;13:698–710. [PubMed: 11506665]
- Seidler RD, Purushotham A, Kim SG, Ugurbil K, Willingham D, Ashe J. Cerebellum activation associated with performance change but not motor learning. *Science* 2002;296:2043–2046. [PubMed: 12065841]
- van der Graaf FHCE, Maguire RP, Leenders KL, de Jong BM. Cerebral activation related to implicit sequence learning in a double serial reaction time task. *Brain Research* 2006;1081:179–190. [PubMed: 16533501]
- Willingham DB, Koroshetz WJ. Evidence for dissociable motor skills in Huntington's disease patients. *Psychobiology* 1993;21:173–182.

Figure 1. Patient Lesions. The lesions were traced on a standard brain by a consulting neurologist using MRIcro software. Brains are depicted with the right hemisphere on the left.

Figure 2.

Reaction time by trial epoch. In epochs 1–4, participants pressed keys in response to a repeated 8-trial sequence of visual stimuli. During epoch 5, the stimuli appeared in pseudorandom order to test learning.

Patient characteristics. Edu = education in years; MCA=middle cerebral artery; L=left; Moya-Moya = a rare cerebrovascular disorder; AVM= arteriovenous malformation, I = inferior, S = superior.

Table 1

Subject	Age	Sex	Edu	Injury Date	Etiology	Lesion Location	Frontal Involvement
592	40	F	13	2001	MCA infarct, Moya Moya	S frontoparietal	+
474	47	F	12	1995	MCA infarct	I parietal, L caudate	-
312	57	F	17	1981	AVM Resection	S parietal, cerebellar atrophy	-
560	60	M	12	2003	MCA infarct	I parietal	-
564	73	F	10	2001	Infarct	S parietal, frontal	+
444	75	F	12	2002	MCA infarct	I temperoparietal	-
Mean	58.6	5F/1M	12.7				