Published in final edited form as:

Nat Rev Cancer. 2008 November ; 8(11): 875–879. doi:10.1038/nrc2522.

Ageing, oxidative stress and cancer: paradigms in parallax

Christopher C. Benz and Christina Yau

Buck Institute for Age Research, Novato, California 94945, USA.

Abstract

Two paradigms central to geroscience research are that aging is associated with increased oxidative stress and increased cancer risk. Therefore, it could be deduced that cancers arising with ageing will show evidence of increased oxidative stress. Recent studies of gene expression in age-controlled breast cancer cases indicate that this deduction is false, posing parallax views of these two paradigms, and highlighting the unanswered question: does ageing cause or simply permit cancer development?

The origins of the free radical theory of ageing date back to the 1950s, when Harman proposed that oxygen free radicals formed endogenously from normal metabolic processes that use oxygen, have an essential role in the ageing process¹. Over the following decades, equivocal evidence emerged regarding the accumulation of oxidatively damaged proteins and DNA with normal ageing². Such macromolecular damage, if not sufficiently repaired, can cause the progressive failure of cellular machinery, organ ageing and the onset of age-related disease. Many pathologies have been linked to oxidative stress, including atherosclerosis, hypertension, ischaemia–reperfusion injury, inflammation, cystic fibrosis, diabetes, Parkinson and Alzheimer diseases, and cancer. Given that mitochondria are the major intracellular generators of reactive oxygen species (ROS), which are the inevitable consequence of oxidative ATP production from electron transport along the mitochondrial inner membrane, this theory has largely become known as the mitochondrial free radical theory of ageing. Although supporting evidence is substantial, it is also largely correlative, and counter-evidence and variants of this theory also exist^{2–6} (BOX 1). Currently, most experts believe that mitochondrially generated oxidative stress is particularly important in age-related diseases but it is not the sole cause of ageing⁷. This Perspective will discuss some new and unexpected findings regarding the association of oxidative stress with ageing, as evaluated in the common age-related disease breast cancer.

Correspondence to C.C.B. e-mail: cbenz@buckinstitute.org.

DATABASES

Entrez Gene:

<http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=gene>
BCL2

National cancer institute: <http://www.cancer.gov>
breast cancer

UniProtKB:

<http://www.uniprot.org/>
endothelin 2 | ER | PR | RANK | SP1 | SRC | thymidine phosphorylase | TGFβ |

FURTHER INFORMATION

C. C. Benz's homepage:

http://www.buckinstitute.org/site/index.php?Itemid=96&id=103&option=com_content&task=view

TFSEARCH:

<http://www.cbrc.jp/research/db/TFSEARCH.html>

ALL LINKS ARE ACTIVE IN THE ONLINE PDF

Ageing, breast cancer and epigenetics

Even in animals with vastly different lifespans, the incidence of malignancies generally increases with age. Overall, the cancer incidence in humans increases exponentially with age, with 75% of newly diagnosed cases occurring in susceptible populations aged 55 years or older, which supports the epidemiological conclusion that age is the strongest demographic risk factor for most human malignancies^{8,9}. Given the worrying social, economic and medical consequences of an ageing worldwide population, there is a pressing need to understand the biological link or links between cancer and ageing. Despite the awareness that breast cancer and other epithelial cancers are primarily age-related diseases, molecular and cellular hypotheses explaining the relationship between cancer and ageing have only recently emerged^{10–13}. At the subcellular level, normal human ageing has been linked to increased genomic instability^{14,15}, global and promoter-specific epigenetic changes^{16,17}, and the altered expression of genes involved in cell division and extracellular matrix remodeling^{14, 15}. These associations have led to the hypothesis that the cancer-prone phenotype of an older individual results from the combined effects of cumulative mutational load, increased epigenetic gene silencing, telomere dysfunction and an altered stromal milieu¹⁸. However, clinical assessment of this hypothesis is challenging, as human malignancies such as breast cancer are phenotypically and genotypically heterogeneous. This heterogeneity must be accounted for in any study design that tries to evaluate differences between early- and late-onset forms of the same cancer. When one prospectively designed study evaluated early- and late-onset forms of the same histological type of sporadic human breast cancer, oestrogen receptor (ER)-positive breast cancer, the conclusion that the biology of this disease was indeed age-dependent was tempered by a surprising lack of support for the more general cancer-ageing hypothesis¹⁹. Rather than displaying a ‘mutator’ phenotype predisposed to genetic instability, accelerated proliferation and more aggressive growth, late-onset ER-positive breast cancers, like many other late-onset human malignancies, appear less aggressive and are associated with a better patient prognosis than early-onset forms of histologically identical ER-positive breast cancers. Remarkably, this prospective study showed that during the normal female lifespan ER-positive breast cancers that arose over 30 years apart did so by fundamentally different epigenetic programmes and not by any detectable differences in genomic mechanisms¹⁹. In addition, an age signature composed of 128 genes that were differentially expressed between these early- and late-onset breast cancers proved to be >80% accurate at discerning younger ER-positive breast cancers from older ER-positive breast cancers in two other independent data sets¹⁹. Do these epigenetic differences simply reflect age-associated changes in the mammary gland²⁰ or can they be explained by the accumulated effects of oxidative stress thought to accompany ageing?

Breast cancer and oxidative stress

As a putative aetiological factor for both ageing and age-related diseases such as ER-positive breast cancer, oxidative stress is an attractive mechanism with which to explain age-dependent differences in gene expression and cancer biology. Although there is a variety of evidence that suggests that reactive oxygen and nitrogen species contribute to the age-related development of cancers, the cellular sources and carcinogenic mechanisms of these molecules remain unclear²¹. ROS are critical mediators of growth factor receptor signalling²² and are involved in oestrogen-inducible cancer cell proliferation^{23,24}. Not only has the carcinogenic potential of oestrogen exposure been attributed to its oxidation chemistry^{25,26}, but oxidative stress pathways activated during cell immortalization and transformation have also been correlated with clinical prognosis in patients with breast cancer²⁷. Non-transformed breast epithelial cells experience variable fluctuations in the production of ROS from mitochondrial respiration, peroxisomal β -oxidation and cytochrome P450 xenobiotic metabolism²⁸. In transformed breast epithelial cells, constitutively activated mitogenic pathways increase intracellular ROS

production. Transformed metabolic pathways, including increased thymidine phosphorylase expression²⁹ and the lacto-peroxidase metabolism of oestrogen³⁰ further increase the levels of endogenous ROS. Inadequate tumour neovascularization results in excess oxidative stress as well as glucose deprivation, hypoxia-activating kinase cascades and decreasing antioxidant defences^{31,32}. In turn, hypoxia stimulates the expression of chemoattractants such as endothelin 2 in breast cancer³³, which recruit ROS-producing macrophages that accumulate within the hypoxic regions of various tumour types³⁴. Whether it results from organ ageing, endogenous metabolites, malignant transformation pathways or rapid neoplastic growth and metabolism (BOX 2), oxidative stress accompanies breast cancer development and progression to varying degrees, and may therefore be a plausible underlying mechanism for much of the biological and clinical diversity of ER-positive breast cancers. How can oxidative stress alter aspects of tumour biology, such as the endocrine pathways that drive ER-positive breast cancer?

Box 1|Mitochondrial free radical theory of ageing

In 1956, Denham Harman first proposed that the oxygen free radicals that are endogenously formed from normal metabolic processes in a variety of organisms cause ageing. By 1972, Harman recognized the dominant role of mitochondria in the generation of intracellular reactive oxygen species (ROS) and revamped his proposal into the mitochondrial free radical theory of ageing. With accumulating evidence, and following decades of studies that have involved both invertebrate and vertebrate model systems, there is continued controversy over whether an accumulation of macromolecular damage caused by chronic ROS production limits mammalian lifespan or whether it primarily contributes to the onset of age-related disease. Accepted refinements to this theory now include the chemical participation of reactive nitrogen species (RNS) as well as ROS, recognition that mitochondrial (mt) DNA damage also accumulates with ageing, and appreciation that the balance of intracellular antioxidant and macromolecular repair mechanisms is crucial in determining the cell fate responses to both acute and chronic oxidative stress. It is also important to note that the free radical theory of ageing is not mutually exclusive of other ageing mechanisms (for example, cell senescence, telomere shortening and genomic instability).

Evidence in support of the mitochondrial free radical ageing theory in mammals includes the following:

- mitochondrial ROS production and mtDNA damage (for example, deletions, mutations and base modifications) increase with age in various mammals, including mice and humans
- injection of chemically uncoupled or aged mitochondria induces cellular degeneration of young cells
- knock-in mice with catalase overexpression localized to the mitochondria exhibit reduced levels of mtDNA damage and have an extended lifespan
- caloric restriction reduces mitochondrial ROS production and mtDNA damage and extends lifespan

Evidence that challenges the mitochondrial free radical ageing theory in mammals includes the following:

- transgenic mice that express an error-prone mtDNA polymerase show accelerated signs of ageing and have a decreased lifespan but are not subject to ROS overproduction or oxidative stress

- knockout mice heterozygous for the superoxidase dismutase 2 gene have increased oxidative damage in their nuclear DNA and mtDNA, but exhibit neither signs of accelerated ageing nor a reduced lifespan

Oxidative stress and ER activity

In addition to DNA damage, there are at least two major consequences of excess ROS production for proteins that affect ER path-ways and the endocrine responsiveness of ER-positive breast cancer: direct oxidative injury to protein structure and ROS-induced kinase signalling. Among the intracellular proteins that are most vulnerable to direct oxidant damage are redox-sensitive nuclear transcription factors, such as ER³⁵ and SP1 (REF 36), whose zinc finger cysteine residues are readily oxidized preventing their DNA-binding function. In ER-positive breast cancers, loss of SP1 DNA-binding activity has been correlated with ageing in association with an increase in the levels of the oxidative stress marker phospho-ERK5 (extracellular signal-related kinase 5, also known as mitogen-activated protein kinase 7 (MAPK7))³⁷ in the tumour. Although a complete loss of the DNA-binding function of the ER that is extracted from primary human breast cancers has not been specifically linked to ageing, this loss has been shown to occur in up to one-third of all ER-positive primary breast cancers and has been correlated with loss of progesterone receptor (PR) expression³⁸, because the DNA-binding and transactivating functions of both ER and SP1 are needed for the optimal oestrogenic stimulation of genes such as *PR* and *BCL2*, ER-positive breast cancers that have been subjected to sufficient levels of oxidative stress would be expected to show suppressed expression of *PR*, *BCL2* and other oestrogen-inducible genes. The second major consequence of oxidative stress is its association with kinase-dependent signal transduction. In addition to their role in the mediation of growth factor receptor signalling, ROS directly inhibit protein tyrosine phosphatases and thereby stimulate SRC, Janus kinase, Ras family members, protein kinase C and MAPK signalling³⁹. These activated kinase pathways are known to modulate ER activity⁴⁰ and have been implicated in endocrine resistance^{41–44}. In particular, excess MAPK signalling in ER-positive breast cancer cells impairs oestrogen-inducible gene transcription⁴⁵ and induces a profile of gene expression similar to that of ER-negative breast cancer cells⁴⁶. Thus, oxidative stress can potentially alter the phenotype of an ER-positive breast cancer, in some cases to such an extent that the endocrine responsiveness of the tumour is lost.

Box 2|Cancer and oxidative stress

Oxidative stress in the form of excess reactive oxygen species (ROS) or reactive nitrogen species (RNS) can have either deleterious or beneficial effects on a cell, and may be generated by intracellular or extracellular sources. Oxidative stress may cause carcinogenesis by mutating nuclear or mitochondrial DNA, or by causing structural damage of intracellular lipids and proteins. A growing tumour mass may also produce intracellular and extracellular oxidative stress that can transiently or permanently modify its malignant features.

Endogenous sources of tumour ROS or RNS include an impaired mitochondrial genome or proteome, supporting the Warburg hypothesis; activated extra-mitochondrial growth and metabolism pathways; and xenobiotic metabolism.

Exogenous sources of tumour ROS or RNS include ischaemia and reperfusion; inadequate neovascularization; inflammatory cell infiltrate; and activated or damaged stroma.

Oxidant gene signatures

Is there clinical evidence for the presence of oxidative stress in ER-positive human breast cancers? Although many gene expression studies have attempted to identify the suite of endocrine-responsive genes expressed in ER-positive breast cancers^{47–51}, only one study tried to delineate a subset of endocrine-responsive genes that were also susceptible to modulation by oxidative stress in order to identify and characterize oxidatively stressed ER-positive breast cancers⁵². In this recent study, an ER-positive human breast cancer cell line was subjected to either oestrogen deprivation or ER knockdown and profiled to identify a complete set of genes regulated by oestrogen and ER (designated oestrogen+ER). In addition, the cell line was subjected to stress by three different chemical oxidants (diamide, hydrogen peroxide and menadione), and profiled in order to produce an oxidant signature. The intersection of both the oestrogen+ER and oxidant gene signatures yielded an oxidant–oestrogen+ER signature that was composed of 62 endocrine-responsive genes that were commonly susceptible to the different forms of oxidative stress. Network analysis of the oxidant–oestrogen+ER gene signature from this study highlighted the activation of cancer pathways that regulate cell growth and invasion⁵². The signature derived from the model was used to interrogate a public repository of expression microarray data that characterized 394 ER-positive primary breast cancers, and look for correlations with tumour parameters, patient age at diagnosis and survival outcome. ER-positive breast cancers with higher expression levels of this oxidant–oestrogen+ER gene signature showed a loss of PR expression, high tumour grade and reduced patient survival⁵². Given the link between ageing and oxidative stress, and the expectation that excess levels of oxidative stress would be evident in the breast cancers that arose in older individuals, it was surprising that the ER-positive breast cancers that were diagnosed at an older age did not show higher levels of expression of either the endocrine-specific oxidant–oestrogen+ER signature or the more general ER-independent oxidant signature. In fact, a more refined experimental analysis indicated that oxidative stress was more evident in the ER-positive breast cancers that were diagnosed at a younger age than those that were diagnosed at an older age⁵².

Shared TNF and TGF β pathways

The fact that the most common forms of breast cancer that arise later in life lack evidence of oxidative stress suggests that the putative link between organ ageing and oxidative stress is obscured by the presence of malignancy-transforming metabolic pathways and cellular environments that generate excess ROS. This notion is consistent with two other observations that relate to ER-positive breast cancers and their proliferative potential: first, breast cancers that are enriched for the oxidant–oestrogen+ER signature have higher expression levels of cell proliferation genes than breast cancers that are not enriched for the signature⁵²; and, second, early-onset ER-positive breast cancers have higher expression levels of cell proliferation genes than late-onset breast cancers¹⁹. Tumours with greater proliferative potential have increased levels of intracellular ROS production owing to excessive growth factor signalling, and are also subjected to increased extracellular ROS production from repetitive cycles of ischaemia and reperfusion as neovascularization attempts to keep up with the rapidly expanding tumour mass. If tumour proliferation and growth rates largely explain breast cancer oxidative stress, and both oxidative stress and proliferation gene signatures are more characteristic of early-onset breast cancers, then it might be possible to identify other tumour-associated pathways that are shared by both early-onset and oxidatively stressed ER-positive breast cancers. Comparative network analysis of the age¹⁹ and oxidant–oestrogen+ER⁵² gene signatures reveals two such pathways, the tumour necrosis factor (TNF) and transforming growth factor- β (TGF β) signalling pathways, which are common to both oxidatively stressed and early-onset ER-positive breast cancers (FIG. 1). Interestingly, signals from RANK (receptor activator of nuclear factor κ B (NF κ B)), a TNF receptor superfamily member that is commonly implicated

in breast cancer, and ligand-stimulated TGF β receptors converge to activate intracellular NF κ B and AP1 (FIG. 2). These transcription factor complexes stimulate gene expression programmes that enhance breast cancer cell survival, proliferation, invasiveness and angiogenesis⁵³. Indeed, analysis of the proximal promoter regions (2.5 kb) of the age and oxidant–oestrogen+ER signature genes shown in FIG. 1 indicates that at least 75% of these TNF- or TGF β -regulated genes have either AP1 or NF κ B consensus binding elements. both early-onset¹⁹ and oxidatively stressed⁵² ER-positive breast cancers are associated with an increased risk of metastatic recurrence and reduced patient survival. In addition, ER-positive breast cancers with activation of both NF κ B and AP1 are known to be associated with poor patient prognosis and resistance to endocrine therapy⁵⁴. Therefore, this comparative network analysis provides a supporting rationale for the development of NF κ B and AP1 inhibitors to treat early-onset ER-positive breast cancers that have a more aggressive and treatment-resistant clinical phenotype due at least in part to oxidative stress.

Evolving paradigms

Does ageing cause or simply permit cancer development? Despite a long-standing awareness that breast cancer and other cancers are primarily age-related and the general belief that ageing predisposes to cancer development, the nascent field of geroscience is only beginning to inform oncology of the relationship between cancer and ageing. Therefore, the molecular and cellular hypotheses posited to explain this relationship remain largely untested^{10–13}. Additional age cohort studies of the type described above for ER-positive breast cancer^{19,20} are needed to generalize about the age-dependent biological differences that drive ER-negative breast cancer, as well as other age-related epithelial malignancies. There is growing epidemiological evidence that cancer incidence decelerates or is even suppressed after the age of 80, and experimental investigation of this theory may substantially modify the currently accepted paradigm that ageing promotes cancer development⁵⁵. Is oxidative stress a cause or consequence of ageing, and is such oxidative stress manifested in all ageing organs or in a few? The same cause–effect question may be asked of the mechanistic relationship between oxidative stress and cancer development. If increasing mitochondrial dysfunction promotes normal cell ageing, and excess ROS from increased growth factor signalling and ischaemia–reperfusion injury promotes cancer cell aggressiveness, is there any mechanistic commonality between these disparate cell fates that are linked to oxidative stress? Aged, oxidatively stressed and senescent stroma are permissive for malignant epithelial transformation in various experimental models⁵⁶. However, human organs that are at risk for age-related malignancies such as breast cancer have not been studied sufficiently to determine whether such stromal changes or senescent cell populations actually predate and promote human cancer *in vivo*. It is now apparent that many late-onset malignancies such as ER-positive breast cancer are biologically and clinically more indolent than their earlier-onset counterparts. The increase in the incidence of these late-onset malignancies up to the age of 80 may reflect the existence of a more permissive stromal environment that enables the emergence of cancers that would be unable to thrive in younger hosts. If cancer aggressiveness decreases progressively with ageing and out of pace with its increasing incidence, then perhaps an inflection point is ultimately reached in the human lifespan, whereupon a microscopic malignancy becomes unable to make a clinical appearance. This challenging area of biology and oncology must continue to be examined and questioned from many different angles, as parallax views of our evolving cancer and ageing paradigms are needed to provide a sufficient depth of understanding to successfully confront the growing health-care burden of our ageing population.

References

1. Harman D. Aging: a theory based on free radical and radiation chemistry. *J. Gerontol* 1956;11:298–300. [PubMed: 13332224]

2. Muller FL, Lustgarten MS, Jang Y, Richardson A, Van Remmen H. Trends in oxidative aging theories. *Free Radic. Biol. Med* 2007;43:477–503. [PubMed: 17640558]
3. Finkel T, Holbrook NJ. Oxidants, oxidative stress and the biology of ageing. *Nature* 2000;408:239–247. [PubMed: 11089981]
4. Raha S, Robinson BH. Mitochondria, oxygen free radicals, disease and ageing. *Trends Biochem. Sci* 2000;25:502–508. [PubMed: 11050436]
5. Krishnan KJ, Greaves LC, Reeve AK, Turnbull D. The ageing mitochondrial genome. *Nucleic Acids Res* 2007;35:7399–7405. [PubMed: 17913753]
6. Gruber J, Schaffer S, Halliwell B. The mitochondrial free radical theory of ageing — where do we stand? *Front Biosci* 2008;13:6554–6479. [PubMed: 18508680]
7. Vijg J, Campisi J. Puzzles, promises and a cure for ageing. *Nature* 2008;454:1065–1071. [PubMed: 18756247]
8. Edwards B, et al. Annual report to the nation on the status of cancer, 1973–1999, featuring implications of age and aging on U.S. cancer burden. *Cancer* 2002;94:2766–2792. [PubMed: 12173348]
9. Thun, M.; Jemal, A. *Cancer Epidemiology*. Hamilton: BC Decker; 2006.
10. Ershler WB, Longo DL. Aging and cancer: issues of basic and clinical science. *J. Natl Cancer Inst* 1997;89:1489–1497. [PubMed: 9337345]
11. Balducci L, Ershler WB. Cancer and ageing: a nexus at several levels. *Nature Rev. Cancer* 2005;5:655–662. [PubMed: 16056261]
12. Benz CC, Campisi J, Cohen HJ, Ershler WB, Irminger-Finger I. Meeting report: translational research at the aging and cancer interface. *Cancer Res* 2007;67:4560–4563. [PubMed: 17510382]
13. Pawelec G, Solana R. Are cancer and aging different sides of the same coin? Conference on cancer and ageing. *EMBO Rep* 2008;9:234–238. [PubMed: 18259218]
14. Geigl JB, et al. Analysis of gene expression patterns and chromosomal changes associated with aging. *Cancer Res* 2004;64:8550–8557. [PubMed: 15574761]
15. Ly DH, Lockhart DJ, Lerner RA, Schultz PG. Mitotic misregulation and human aging. *Science* 2000;287:2486–2492. [PubMed: 10741968]
16. Issa JP. Aging, DNA methylation and cancer. *Crit. Rev. Oncol. Hematol* 1999;32:31–43. [PubMed: 10586353]
17. Richardson B. Impact of aging on DNA methylation. *Ageing Res. Rev* 2003;2:245–261. [PubMed: 12726774]
18. DePinho RA. The age of cancer. *Nature* 2000;408:248–254. [PubMed: 11089982]
19. Yau C, et al. Aging impacts transcriptomes but not genomes of hormone-dependent breast cancers. *Breast Cancer Res* 2007;9:R59. [PubMed: 17850661]
20. Benz CC. Impact of aging on the biology of breast cancer. *Crit. Rev. Oncol. Hematol* 2008;66:65–74. [PubMed: 17949989]
21. Halliwell B. Oxidative stress and cancer: have we moved forward? *Biochem. J* 2007;401:1–11. [PubMed: 17150040]
22. Lander HM. An essential role for free radicals and derived species in signal transduction. *FASEB J* 1997;11:118–124. [PubMed: 9039953]
23. Felty Q, Singh KP, Roy D. Estrogen-induced G₁/S transition of G₀-arrested estrogen-dependent breast cancer cells is regulated by mitochondrial oxidant signaling. *Oncogene* 2005;24:4883–4893. [PubMed: 15897899]
24. Oberley TD, Allen RG, Schultz JL, Lauchner LJ. Antioxidant enzymes and steroid-induced proliferation of kidney tubular cells. *Free Radic. Biol. Med* 1991;10:79–83. [PubMed: 2050299]
25. Patel MM, Bhat HK. Differential oxidant potential of carcinogenic and weakly carcinogenic estrogens: Involvement of metabolic activation and cytochrome P450. *J. Biochem. Mol. Toxicol* 2004;18:37–42. [PubMed: 14994278]
26. Bhat HK, Calaf G, Hei TK, Loya T, Vadgama JV. Critical role of oxidative stress in estrogen-induced carcinogenesis. *Proc. Natl Acad. Sci. USA* 2003;100:3913–3918. [PubMed: 12655060]
27. Dairkee SH, et al. Oxidative stress pathways high-lighted in tumor cell immortalization: association with breast cancer outcome. *Oncogene* 2007;26:6269–6279. [PubMed: 17471242]

28. Beckman KB, Ames BN. The free radical theory of aging matures. *Physiol. Rev* 1998;78:547–581. [PubMed: 9562038]
29. Moghaddam A, et al. Thymidine phosphorylase is angiogenic and promotes tumor growth. *Proc. Natl Acad. Sci. USA* 1995;92:998–1002. [PubMed: 7532308]
30. Sipe HJ Jr, Jordan SJ, Hanna PM, Mason RP. The metabolism of 17 β -estradiol by lactoperoxidase: a possible source of oxidative stress in breast cancer. *Carcinogenesis* 1994;15:2637–2643. [PubMed: 7955118]
31. Blackburn RV, et al. Metabolic oxidative stress activates signal transduction and gene expression during glucose deprivation in human tumor cells. *Free Radic. Biol. Med* 1999;26:419–430. [PubMed: 9895234]
32. Li C, Jackson RM. Reactive species mechanisms of cellular hypoxia-reoxygenation injury. *Am. J. Physiol. Cell Physiol* 2002;282:C227–C241. [PubMed: 11788333]
33. Grimshaw MJ, Naylor S, Balkwill FR. Endothelin-2 is a hypoxia-induced autocrine survival factor for breast tumor cells. *Mol. Cancer Ther* 2002;1:1273–1281. [PubMed: 12516960]
34. Grimshaw MJ, Wilson JL, Balkwill FR. Endothelin-2 is a macrophage chemoattractant: implications for macrophage distribution in tumors. *Eur. J. Immunol* 2002;32:2393–2400. [PubMed: 12207323]
35. Liang X, et al. Oxidant stress impaired DNA-binding of estrogen receptor from human breast cancer. *Mol. Cell Endocrinol* 1998;146:151–161. [PubMed: 10022773]
36. Wu X, Bishopric NH, Discher DJ, Murphy BJ, Webster KA. Physical and functional sensitivity of zinc finger transcription factors to redox change. *Mol. Cell Biol* 1996;16:1035–1046. [PubMed: 8622648]
37. Quong J, et al. Age-dependent changes in breast cancer hormone receptors and oxidant stress markers. *Breast Cancer Res. Treat* 2002;76:221–236. [PubMed: 12462383]
38. Scott GK, Kushner P, Vigne JL, Benz CC. Truncated forms of DNA-binding estrogen receptors in human breast cancer. *J. Clin. Invest* 1991;88:700–706. [PubMed: 1864980]
39. Valko M, Rhodes CJ, Moncol J, Izakovic M, Mazur M. Free radicals, metals and antioxidants in oxidative stress-induced cancer. *Chem. Biol. Interact* 2006;160:1–40. [PubMed: 16430879]
40. Levin ER. Bidirectional signaling between the estrogen receptor and the epidermal growth factor receptor. *Mol. Endocrinol* 2003;17:309–317. [PubMed: 12554774]
41. Kirkegaard T, et al. AKT activation predicts outcome in breast cancer patients treated with tamoxifen. *J. Pathol* 2005;207:139–146. [PubMed: 16088978]
42. Knowlden JM, et al. Elevated levels of epidermal growth factor receptor/c-erbB2 heterodimers mediate an autocrine growth regulatory pathway in tamoxifenresistant MCF-7 Cells. *Endocrinology* 2003;144:1032–1044. [PubMed: 12586780]
43. Nabha SM, et al. Upregulation of PKC- δ contributes to antiestrogen resistance in mammary tumor cells. *Oncogene* 2005;24:3166–3176. [PubMed: 15735693]
44. Gee JM, et al. Epidermal growth factor receptor/HER2/insulin-like growth factor receptor signalling and oestrogen receptor activity in clinical breast cancer. *Endocr. Relat. Cancer* 2005;12:S99–S111. [PubMed: 16113104]
45. Oh AS, et al. Hyperactivation of MAPK induces loss of ER α expression in breast cancer cells. *Mol. Endocrinol* 2001;15:1344–1359. [PubMed: 11463858]
46. Creighton CJ, et al. Activation of mitogen-activated protein kinase in estrogen receptor α -positive breast cancer cells *in vitro* induces an *in vivo* molecular phenotype of estrogen receptor α -negative human breast tumors. *Cancer Res* 2006;66:3903–3911. [PubMed: 16585219]
47. Creighton CJ, et al. Genes regulated by estrogen in breast tumor cells *in vitro* are similarly regulated *in vivo*. *Genome Biol* 2006;7:R28. [PubMed: 16606439]
48. Coser KR, et al. Global analysis of ligand sensitivity of estrogen inducible and suppressible genes in MCF7/BUS breast cancer cells by DNA microarray. *Proc. Natl Acad. Sci. USA* 2003;100:13994–13999. [PubMed: 14610279]
49. Lin CY, et al. Discovery of estrogen receptor α target genes and response elements in breast tumor cells. *Genome Biol* 2004;5:R66. [PubMed: 15345050]
50. Oh DS, et al. Estrogen-regulated genes predict survival in hormone receptor-positive breast cancers. *J. Clin. Oncol* 2006;24:1656–1664. [PubMed: 16505416]

51. Vendrell JA, et al. Estrogen regulation in human breast cancer cells of new downstream gene targets involved in estrogen metabolism, cell proliferation and cell transformation. *J. Mol. Endocrinol* 2004;32:397–414. [PubMed: 15072547]
52. Yau C, Benz CC. Genes responsive to both oxidant stress and loss of estrogen receptor (ER) function identify a poor prognosis group of ER-positive primary breast cancers. *Breast Cancer Res* 2008;10:R61. [PubMed: 18631401]
53. Zhou Y, et al. Enhanced N κ B and AP-1 transcriptional activity associated with antiestrogen resistant breast cancer. *BMC Cancer* 2007;7:59. [PubMed: 17407600]
54. Zhou Y, et al. Activation of nuclear factor κ B (NF κ B) identifies a high-risk subset of hormone-dependent breast cancers. *Int. J. Biochem. Cell Biol* 2005;37:1130–1144. [PubMed: 15743683]
55. Harding C, Pompei F, Lee EE, Wilson R. Cancer suppression at old age. *Cancer Res* 2008;68:4465–4478. [PubMed: 18519710]
56. Campisi J, d'Adda di Fagagna F. Cellular senescence: when bad things happen to good cells. *Nature Rev. Mol. Cell Biol* 2007;8:729–740. [PubMed: 17667954]
57. Zhang H, Ramanathan Y, Soteropoulos P, Recce ML, Toias PP. EZ-Retrieve: a web-server for batch retrieval of coordinate-specified human DNA sequences and underscoring putative transcription factor-binding sites. *Nucleic Acid Res* 2002;30:e121. [PubMed: 12409480]
58. Heinemeyer T, et al. Databases on transcriptional regulation: TRANSFAC, TRRD and COMPEL. *Nucleic Acid Res* 1998;26:364–370.

Acknowledgments

The authors' work is supported in part by National Institutes of Health grants R01-AG020521, R01-CA71468, P01-AG025901, U54-RR024346/RL1-AG032113, P50-CA58207, and Hazel P. Munroe memorial funding to the Buck Institute.

Figure 1. Pathways that are shared by oxidatively stressed and early-onset breast cancers

Top-scoring gene networks, which are determined by ingenuity Pathways systems analysis of age¹⁹ and oxidative stress⁵² gene signatures in oestrogen receptor (ER)-positive breast cancers, commonly identify tumour necrosis factor (TNF) and transforming growth factor- β (TGF β) pathway nodes. Ariadne Pathway studio was used to identify the respective TNF- and TGF β - regulated (direct or indirect) targets from the 126-gene age signature (**a**) and the 62-gene oxidant-oestrogen+ER signature (intersection of oestrogen- and ER-regulated signature and oxidant-stressed signature; **b**). Regulatory directions were determined from primary references provided by Pathway studio, with positive (+) regulatory directions denoted by arrow-heads and negative regulatory directions denoted by bars (lines without either arrowheads or bars indicate ambiguous regulatory direction). Unigene symbols in red ovals indicate signature genes that are upregulated by oxidative stress or in early-onset breast cancer; those in green ovals indicate signature genes that are downregulated by oxidative stress or in early-onset breast cancer. At least 75% of the signature genes that have been shown to be regulated by TNF and TGF β (**a**, **b**) contain nuclear factor κ B (NF κ B) or AP1 consensus element binding sites within their proximal promoters, as determined by EZRetrieve and TFSEARCH^{57–58} (see TFSEARCH in Further information). ABCC3, ATP-binding cassette 3; ANG, angiogenin; AREG, amphiregulin; CDC2, cell division cycle 2; CLDN1, claudin 1; CTGF, connective tissue growth factor; CXCR4, chemokine (C-X-C motif) receptor 4; CXCL13, chemokine (C-X-C motif) ligand 13; CYP1A1, cytochrome P450 1A1; EGR1, early growth response 1; DAB2, disabled homologue 2; EFNA1, ephrin A1; ESR1, oestrogen receptor 1; GAL, galanin prepropeptide; IGFBP5, insulin-like growth factor binding protein 5; ITPR1, inositol 1,4,5-triphosphate receptor 1; KRT17, kera-tin 17; LAMC2, laminin- γ 2; LCN2, lipocalin 2; LEF1, lymphoid enhancer-binding factor 1; LRP2, low density lipoprotein-related protein 2; MMP9, matrix metalloproteinase 9; PLN, phospholamban; PLIN, perilipin; PTHLH, parathyroid hormone-like hormone; PTGES, prostaglandin E synthase; SERPINA1, serpin peptidase inhibitor 1; SOD3, superoxide dismutase 3; TAGLN, transgelin; TNC, tenascin C; TNFSF10, tumour necrosis factor (ligand) superfamily 10.

Figure 2. Signalling pathways in oxidatively stressed and early onset breast cancers

RANK (receptor activator of nuclear factor κ B (NF κ B)) is a tumour necrosis factor (TNF) receptor superfamily member that is commonly implicated in breast cancer, and ligand-stimulated RANK and transforming growth factor- β (TGF β) receptors are shown as convergent signal transduction pathways that are capable of activating intracellular NF κ B and AP1 transcription factor complexes. These can induce gene expression programmes that promote breast cancer cell survival, proliferation, invasiveness and angiogenesis. ERK, extracellular signal-regulated kinase; I κ B, inhibitor of NF κ B; IKK, I κ B kinase; JNK, JUN N-terminal kinase; MEK, MAPK/ERK kinase; MKK, mitogen-activated protein kinase kinase; TRAF6, TNF receptor-associated factor 6; UBC13, ubiquitin-conjugating enzyme E2 13; UEV1A, ubiquitin-conjugating enzyme E2 variant 1A.