Published in final edited form as:

Schizophr Res. 2008 August ; 103(1-3): 11–21. doi:10.1016/j.schres.2008.04.035.

Corpus Callosal Area Differences and Gender Dimorphism In Neuroleptic-Naïve, Recent-Onset Schizophrenia And Healthy Control Subjects

John P John, M.B.B.S., M.D.^{1,*}, Mohammed Shakeel K., M.A. (Psychology)¹, and Sanjeev Jain, M.B.B.S., M.D., D.P.M.¹

¹Department of Psychiatry, National Institute of Mental Health and Neurosciences (NIMHANS), Bangalore.

Abstract

The study of corpus callosal morphometry is important to unravel the underlying connectivity disturbance in schizophrenia. We studied the corpus callosal area in schizophrenia subjects compared to healthy subjects, while controlling for several confounders that could affect morphometric measures of the corpus callosum (CC). Areas of the whole CC and its subregions obtained by two geometric partitioning schemes were studied in 23 right-handed neuroleptic-naïve, recent-onset, schizophrenia patients and compared with 23 right-handed age-, sex- and education-matched healthy subjects. The patients did not differ from controls in whole CC area. On tripartite division of the CC, the area of the anterior sub-region was significantly higher in patients compared to controls. On radial division into 5 sub-regions, the anterior truncus area was significantly higher in patients compared to controls. There was a significant effect of gender (F>M) on the area measures; however there was no significant diagnosis*gender effect. Age, age of onset, duration of illness and psychopathology ratings did not show any significant correlations with whole CC area and area of CC sub-regions. The finding of increased area of the anterior truncus that possibly comprises white fibres connecting the temporal association cortices could be indicative of an “abnormal functional hyperconnection” involving these regions in positive symptom schizophrenia. Additionally, the finding of females having larger areas of the whole CC and of the anterior and middle sub-regions could reflect a “normal hyperconnection” underlying increased ambilaterality in females.

1. Introduction

Schizophrenia is conceptualized as a disorder wherein aberrant cortico-cortical connectivity might play an important role (McGuire and Frith, 1996). Inter-hemispheric connectivity disturbances in schizophrenia are reflected as loss of normal asymmetry resulting in reduced laterality (Gruzelier, 1999). The corpus callosum (CC), being the largest mass of connecting white fibres in the brain has therefore evoked much interest among schizophrenia researchers. The majority of CC fibres constitute homotopic interconnections between association cortices of the hemispheres (Aboitiz et al., 1992) Callosal connections between homologous prefrontal and temporoparietal association areas are largely comprised of poorly myelinated, small-caliber (<2 nm diameter) and slowly conducting fibres, whereas those connecting the primary

*Address for correspondence: John P John, MD, Assistant Professor of Psychiatry, National Institute of Mental Health and Neurosciences (NIMHANS), P.B. No. 2900, Dharmaram P.O., Hosur Road, Bangalore, 560 029, Karnataka, INDIA, Telephone Numbers: +91-80-26995349 / 26995306, Fax: +91-80-26564822, Email: jppj@nimhans.kar.nic.in / jppjinc@yahoo.com.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

and secondary sensorimotor areas comprise fast-conducting highly myelinated large-calibre fibres ($>3\mu\text{m}$ diameter) (Aboitiz et al., 1992).

Following the initial report of increased thickness of the CC in postmortem brains of schizophrenia patients by Rosenthal and Bigelow (1972) and the subsequent MRI replication of this finding by Nasrallah et al (1986), the morphometry of CC has been extensively studied in schizophrenia. Shenton et al. (2001), reviewing morphometric studies in schizophrenia, pointed out that among 27 MRI studies of the CC, 17 (63%) reported positive findings while 10 (37%) reported negative findings. The findings across studies have been largely inconsistent, with whole CC areas being reported variously as increased (Nasrallah et al., 1986; Jacobsen et al., 1997), decreased (Woodruff et al., 1993) or not different (Uematsu and Kaiya, 1988; Gunther et al., 1991) between schizophrenia subjects and healthy controls.

Partitioning the CC and comparing the morphometry of the sub-regions between schizophrenia patients and controls has the potential to reveal more subtle morphological abnormalities (Jacobsen et al., 1997). Different partitioning schemes have been proposed for partitioning the CC into 3 to 7 sub-regions (de Lacoste et al., 1985; Witelson, 1989; Woodruff et al., 1993; Rumsey et al., 1996; Jancke et al., 1997; Bachmann et al., 2003; Teipel et al., 2004; Keshavan et al., 2007). Amongst these, the most popular method has been the Witelson (1989)'s scheme, which divides the CC into 7 sub-divisions using criteria based on the results of experimental work with monkeys and humans. However, none of the above partitioning schemes, including that by Witelson (1989), exactly mirrors the texture of the CC at the cellular level (Hofer and Frahm, 2006). Nevertheless, structural MRI (sMRI)-based geometric partitioning schemes continue to be widely used for studying the morphometry of the CC in relation to clinical and other biological variables, owing to the ease of studying the morphometry of CC that these techniques provide. We therefore, chose to geometrically partition the CC into 3 [anterior (rostrum+genu+anterior midbody), mid-body and posterior (isthmus+splenium) (Aboitiz et al., 2003)] and 5 [rostrum and genu, anterior truncus, middle truncus, posterior truncus and isthmus and splenium) (Hampel et al., 1998)] equidistant sub-regions for group comparisons.

Previous studies that divided the CC into 3 to 5 sub-regions found increased thickness in the middle and anterior but not the posterior, parts of the callosal body (Nasrallah et al., 1986), larger anterior third region (Uematsu and Kaiya, 1988), smaller mid-anterior and mid-posterior quarters (Woodruff et al., 1993), or no regional difference (Hauser et al., 1989) in schizophrenia subjects compared to healthy controls.

The possible reasons for the inconsistent findings of whole CC area and that of the subregions in schizophrenia subjects include confounding effects of gender (Witelson, 1989; de Lacoste et al., 1990), handedness (Witelson and Goldsmith, 1991), age (Woodruff et al., 1995; Keshavan et al., 2007), chronicity of illness (Woodruff et al., 1993; Tibbo et al., 1998), heterogeneity in symptom profiles of the samples studied (Gunther et al., 1991; Tibbo et al., 1998) and medication status along with differences in the CC partitioning schemes employed by the different studies as mentioned above. With a view to controlling for these confounders, we aimed at studying the CC morphometry in neuroleptic-naïve, recent-onset, schizophrenia (NRS) subjects in comparison to age, sex and education-matched healthy subjects (HS). Only right-handed subjects were included in the study. We measured the whole CC area as well as the areas of sub-regions derived by semi-automated partitioning of the mid-sagittal CC area respectively into 3 and 5 equidistant partitions. We hypothesized that whole CC areas will not be different between NRS patients and HS, whereas significant differences in area would emerge in the CC sub-regions, that connect cortical regions implicated in recent-onset schizophrenia.

2. Methods

The study was carried out at the National Institute of Mental Health and Neurosciences (NIMHANS), Bangalore India, with due approval from the NIMHANS Ethics Committee, thus conforming to the ethical standards laid down in the 1964 Declaration of Helsinki. Written informed consent was obtained from all the subjects (and their legally qualified representatives in case of patients) prior to enrollment into the study.

2.1. Subjects

2.1.1. Patients—Twenty three right-handed NRS patients were recruited into the study by purposive sampling, from those who attended the outpatient services of NIMHANS. The inclusion criteria were: a DSM-IV diagnosis of schizophrenia or schizophreniform disorder, right-handedness [as assessed by modified Annett's inventory (1976)], duration from onset of illness ≤ 5 years and age between 17–50 years. The diagnosis of schizophrenia or schizophreniform disorder was arrived at using criteria from the Diagnostic and Statistical Manual for Mental Disorders-Fourth edition (DSM-IV) based on the consensus of a research psychiatrist who conducted a semi-structured interview and a trained research assistant who used the Mini International Neuropsychiatric Interview (MINI) Plus (Sheehan et al., 1998). The exclusion criteria included previous exposure to psychotropic drugs; significant suicidal or homicidal risk or other disruptive behaviour which warranted immediate interventions and history of ECT within the previous 6 months. The baseline severity of schizophrenia psychopathology was evaluated using the Positive and Negative Syndrome Scale (PANSS) (Kay et al., 1987) for all subjects by one trained rater (V.A.) who had established good inter-rater reliability with another trained rater (B.R.) [Intra-class correlation co-efficient (ICC) for: PANSS positive symptom subscale = 0.87; negative symptom subscale = 0.92; general psychopathology subscale = 0.85; total score = 0.84]. Overall clinical status was assessed using Clinical Global Impression-Severity (CGI-S) (Guy, 1976). The psychopathology and CGI-S ratings were carried out during the initial one week following recruitment, with the psychopathology during the previous week being considered for rating. In a majority of patients, who were co-operative for psychopathology assessment on the day of recruitment itself, the PANSS ratings were completed in the untreated state; however, in the remaining patients, psychopathology ratings could only be completed following initiation of treatment, but within one week of recruitment.

2.1.2. Healthy Subjects—Twenty three age-, sex- and education-matched right-handed healthy individuals were recruited by word of mouth. These subjects, predominantly constituted by hospital staff and attendants of hospitalized patients, were interviewed by a trained research assistant using MINI-Plus and ascertained to have no present or past Axis I or II psychiatric disorders; in addition, there was no history of psychotic disorders in their first degree relatives, as ascertained using an unstructured clinical interview of the recruited subjects. Further, these subjects did not have any medical/neurological condition requiring regular medications; current exposure to psychotropic medications was also ruled out.

The socio-demographic and clinical characteristics of the subjects are shown in Table 1. The patients and controls were matched for gender, age and educational status. All the patients had a rating of ≥ 3 on delusions and or hallucinatory behavior on PANSS.

2.2. Image acquisition and pre-processing

MR scans were acquired on a Siemens MAGNETOM Vision 1.5 Tesla MRI machine at NIMHANS using an MPRAGE sequence (TR = 9.7 ms, TE = 4 ms, flip angle = 10° , section thickness = 1.25 mm, field of view = 256×256 mm, matrix = 256×256 , number of acquisitions = 3, number of slices = 128, orientation = sagittal, scanning time = 6min, 36s) that acquires

1.25 mm × 1 mm × 1 mm voxels across the entire cranium. The MR scans were then reoriented to transverse plane and interpolated in to 0.5 mm × 0.5 mm × 0.5 mm isotropic voxels using trilinear interpolation. The scans were preprocessed and rotated into AC-PC co-ordinates using Mayo 3D brain Atlas module of Analyze™ 7.0 software (Robb et al. 1989), as previously described in John et al. (2007)

2.3. Corpus Callosum measurements

Outlining of the whole CC as well as grid and radial partitioning were performed using Analyze™ (Robb et al., 1989) on the mid-sagittal section. We referred to the guidelines provided by Woodruff et al. (1993) and Talairach and Tournoux (1993) for identifying the mid-sagittal section of the CC, on which morphometric measurements were made. One expert rater and a trained rater (J.P.J. and M.S.K. respectively) independently identified the mid-sagittal section, selecting the slice which best satisfied both or at least one of the above sets of identifying criteria. Since the CC area measurement is made only on one slice per MRI, the data are vulnerable to errors resulting from inconsistencies in the choice of the “best mid-sagittal slice” (McCarley et al., 1999). Combination of both the above sets of criteria resulted in a highly reliable identification of the midsagittal slice (Intra-class Correlation Coefficient: standardized item alpha=0.99).

Manual tracing of the CC and generation of the object map on the mid-sagittal section was performed using the ‘Edit’ function of Analyze™ (Fig 1). The CC measurements included the whole CC area, areas of the three individual segments generated by the tripartite division as well as areas of the 5 sub-regions generated by the radial division. Partitioning into 3 and 5 sub-regions respectively was performed on the whole CC object map using the “region-of-interest (ROI)” module of Analyze™. For tripartite division, the “grid” option was used to divide the object (whole CC) vertically into 3 equal partitions (CC-I to CC-III), which were saved as anterior, middle and posterior sub-regions respectively and their areas generated (Fig 1). For radial division into 5 sub-regions, the whole CC object was radially divided into 5 equal partitions (CC1–CC5). The origin of the radial lines corresponded to the co-ordinates of the mid-point of the base of the minimum enclosing rectangle (MER) encompassing the CC (Details of the radial division given in Appendix I). These 5 equal partitions were saved as rostrum and genu (CC1), anterior truncus (CC2), middle truncus (CC3), posterior truncus (CC4), and isthmus and splenium (CC5) (Fig 1) and their areas were generated. The reliability (standardized item alpha) of the CC area measurements carried out independently by the raters (J.P.J. and M.S.K.) ranged from 0.89 for the whole CC area to 0.99 for the anterior sub-region obtained using tripartite division. Subsequent to establishing satisfactory reliability of the measurements, outlining of the CC and partitioning into sub-regions on the mid-sagittal section as well as estimation of areal measures were performed by the same trained rater (M.S.K.) on all the 46 MRIs. This rater was blind to the group status, sociodemographic data, psychopathology ratings, and the research hypotheses while performing the MRI analysis and measurements. Forty one out of the 46 brains fulfilled both the Woodruff et al. (1993)’s and Talairach and Tournoux (1993)’s criteria for identifying the mid-sagittal section. The mid-sagittal section fully satisfying at least one of the criteria sets was selected in the remaining 5 brains. No significant difference in coronal tilt was noted between MRIs of patients and controls.

Total brain volume (TBV) and intracranial volume (ICV) were estimated using the VBM2 module of SPM2 (www.fil.ion.ucl.ac.uk/spm/). Reliability of TBV and ICV estimates using SPM2 and Analyze™-based manual segmentation methods was estimated on 3 test brains (standardized item alpha=0.99) (John et al., 2007). The TBV was not significantly different between NRS and HS [HS=986.58 mm² (s.d.=109.95); NRS=973.99 mm² (s.d.=147.19);

$F=1.685$; $p<0.202$], as assessed using GLM ANCOVA with diagnosis and gender as fixed factors and age and ICV as co-variables.

2.4. Statistical Analysis

Descriptive statistics were expressed as mean and standard deviation (s.d.) of the whole CC area as well as the areas of the sub-regions (Table 2). All the CC area measures including whole CC area, areas of the tripartite sub-divisions as well as the areas of the 5 sub-regions derived by radial division were normally distributed in both HS and NRS subjects, as assessed using Shapiro-Wilks test of normality. However, the TBV in NRS subjects contained two outliers, while the TBV in HS was normally distributed. For the TBV to be used as a co-variate in the parametric statistics mentioned below, the two outliers were omitted from analysis to ensure that the data satisfied basic assumptions for conducting the parametric tests.

Whole CC area was compared between NRS subjects and HS by General Linear Model (GLM) Analysis of Co-variance (ANCOVA) with diagnosis (NRS vs HS) and gender as fixed factors and age and TBV as co-variables. Areas of the CC sub-regions obtained by geometric partitioning were compared between NRS subjects and HS using GLM Multivariate Analysis of Covariance (MANCOVA) with diagnosis and gender as fixed factors and age and TBV as co-variables. Subsequently, post-hoc univariate ANCOVA was carried out individually for areas of each sub-region. Similarly, the areas of the sub-regions were compared across genders in both NRS and HS groups using GLM ANCOVA with gender as fixed factor and age and TBV as co-variables. Correlation between age, psychopathology ratings and CC areas were carried out by calculating the partial correlation co-efficient with TBV as co-variate. The level of statistical significance was set *a priori* at $p<0.05$.

3. Results

There was no significant difference in whole CC area between NRS subjects and HS (Table 2). There was a significant effect of gender (females>males) on whole CC area ($F=18.774$; $p<0.000$). GLM MANCOVA with the areas of the 3 CC sub-regions (CC-I, CC-II and CC-III) derived by tripartite division as dependent variables, diagnosis and gender as fixed factors and age and TBV as co-variables, however, revealed significant group differences [$F(3,36)=5.369$; $p<0.004$]. There was a significant effect of gender on areas of the tripartite sub-regions [$F(3,36)=6.903$; $p<0.001$], while there was no significant diagnosis*gender effect [$F(3,36)=0.603$; $p<0.617$]. Similarly, GLM MANCOVA with the areas of the 5 CC sub-regions derived by radial division (CC1–CC5) as dependent variables, revealed significant group differences [$F(5,34)=4.916$; $p<0.002$]. There was a significant effect of gender on areas of the radial sub-divisions [$F(5,34)=4.175$; $p<0.005$], while there was no significant diagnosis*gender effect [$F(5,34)=0.520$; $p<0.759$]. Post-hoc univariate GLM ANCOVA revealed significantly higher areas of the CC-I of the tripartite division and the CC2 of the radial division in NRS subjects compared to HS. The comparison of the whole CC area and those of the sub-regions derived by geometric partitioning between NRS subjects and HS are given in Table 2. Scatter plots depicting the distribution of areas of CC-I and CC-2 are shown in Figure 2. The gender effect was significant for all the areal measurements except that of isthmus and splenium ($p<0.086$).

A GLM MANCOVA done separately for NRS subjects and HS with the three CC sub-regions derived by tripartite division as dependent variables, gender as fixed factor and age and TBV as co-variables revealed significant gender differences (females>males) in both HS [$F(3,17)=4.398$; $p<0.018$] and NRS subjects [$F(3,15)=3.587$; $p<0.0391$]. With the radial sub-divisions, the GLM-MANCOVA did not reach significance for HS [$F(5,15)=2.657$; $p<0.065$] but revealed significant gender differences in NRS subjects [$F(5,13)=3.128$; $p<0.045$]. Planned univariate gender-wise comparison of whole CC area and that of the sub-regions, separately for NRS subjects and HS using GLM ANCOVA with age and TBV as co-variables are given

in Table 3. Whole CC area was significantly higher in females in both HS and NRS groups. On tripartite division, females had significantly higher areas of the CC-I and CC-II sub-regions in the HS group, whereas the areas of all three sub-regions were higher in the NRS group. With the radial partitioning, control females had higher areas of CC1, CC2 and CC3, whereas schizophrenia females had higher areas of CC1 and CC2.

Age, age of onset, duration of illness and psychopathology ratings did not show any significant correlations with whole CC area and area of CC sub-regions.

4. Discussion

Previous morphometric studies of the whole CC and its sub-regions derived using various partitioning schemes in heterogeneous schizophrenia samples have yielded intriguing results. The inconsistencies in the findings that have emerged from previous MRI-based studies have been suggested to be due to the influence of various confounders mentioned earlier, as well as due to the variability in the methods of CC partitioning. We attempted to extend the findings of previous studies by controlling for most of these confounders in a comparative study of the mid-sagittal CC area in a sample of right-handed NRS subjects compared to age, sex and education-matched right-handed HS. All the NRS subjects had ratings of ≥ 3 on delusions and or hallucinatory behavior on PANSS. In view of the fact that no partitioning scheme exactly mirrors the texture of the CC at the cellular level (Hofer and Frahm, 2006), we chose to estimate the CC areas by employing three progressively finer and overlapping partitioning methods—whole CC area as well as geometric partitioning into 3 and 5 equidistant sub-regions using grid and radial partitioning respectively. Equidistant geometric partitioning in sMRI-based CC morphometric studies would help to ensure uniformity and ease of comparison across studies.

There was no significant difference in whole CC area, controlling for age and TBV, between NRS subjects and HS. This finding is in concordance with many previous reports which found no significant differences in the whole CC area in schizophrenia subjects (Rossell et al., 2001; Goghari, 2005). However, many authors have reported smaller whole CC areas in schizophrenia patients (Woodruff, 1993; Tibbo 1998; Keshavan, 2002; Bachmann 2003). All these studies had significant differences with respect to sample characteristics as compared to the present study. The study by Woodruff et al. (1993) included subjects with right/left/mixed handedness, the study samples were not age-, or sex-matched, and included chronically ill patients (duration of illness= 11.3 ± 6.6 years; range=2–28 years), who had been on long-term treatment with neuroleptics. The study by Tibbo et al. (1998) also had similar methodological differences with respect to the sample characteristics compared to the present study. The first-episode, neuroleptic-naïve schizophrenia sample in the study by Keshavan et al. (2007) comprised of patients with a much longer duration of illness (212.09 ± 280.91 weeks) as compared to the present study (69.96 ± 63.35 weeks). Further, both the patient and control groups comprised of subjects who differed in their handedness (R/L: patients=23/8; controls=28/3). Bachmann et al. (2003) studied first-episode schizophrenia patients (n=31) compared to healthy controls (n=12); however the sample sizes were unequal and the subjects were not individually matched for age, sex or educational status, and consequently, the gender distribution in the patient and control groups were substantially different.

The tripartite (Rumsey et al., 1996; Anstey et al., 2007) and radial divisions (Georgy et al., 1992; Teipel et al., 2004) have been used in previous morphometric studies of the CC in several neuropsychiatric conditions other than schizophrenia. In the present study, multivariate analyses revealed significant differences between NRS and HS for areal measures of sub-regions derived by both partitioning schemes. Post-hoc univariate analysis revealed that NRS subjects had significantly higher areas of CC-I (anterior sub-region) and CC2 (anterior truncus). This finding of increased area of the anterior CC is in agreement with previous reports

of increased area, thickness or width of this region in schizophrenia patients (Nasrallah et al., 1986; Uematsu and Kaiya, 1988; Narr et al., 2000). Larger anterior and posterior CC areas have been reported in childhood-onset schizophrenia (Jacobsen et al., 1997). Moreover, an enlarged whole CC size has been suggested to be associated with positive-symptom, good-prognosis schizophrenia (Gunther et al., 1991). Of the above studies, Nasrallah et al. (1986) divided the maximum antero-posterior length of the CC into 4 equidistant partitions using lines perpendicular to the ventral callosal wall, while Uematsu and Kaiya (1988) sub-divided the CC using the tripartite division. Narr et al. (2000) and Jacobsen et al. (1997) employed the Witelson et al.'s (1989) scheme for CC partitioning.

Functional topographic organization of the CC in the human brain is not fully understood. In primates, fibres from frontal cortical areas are shown to traverse the CC through the rostrum and genu, while those from the primary sensorimotor, posterotemporal, parietal and occipital cortical areas pass through the truncus and splenium (Pandya et al., 1971). In humans, fibres from the inferior frontal cortex and anterior inferior parietal regions have been shown to course through the rostrum and genu of the CC, while those from the parieto-occipital junction pass through the posterior truncus and splenium (de Lacoste et al., 1985). The rostrum, genu and the anterior truncus project mainly in the frontal and posterotemporal lobe while the rest of the callosal body and splenium send their fibres to the parietal and occipital cortices, as shown by intra-operative electrophysiologic stimulation of the CC in patients with epilepsy (Tan et al., 1991). Thus, there is no clear consensus regarding the topographical organization of fibres coursing through the human CC. In the present study, the significantly larger area of the anterior sub-region of the CC (CC-I), derived by the tripartite grid partitioning was further localized to the anterior truncus (CC2) using the radial partitioning into 5 sub-regions. As mentioned in the introduction, the callosal connections between homologous prefrontal and temporoparietal association areas, comprised of small-caliber (<2 nm diameter) fibres, are most dense in the anterior sub-region of the CC (Aboitiz et al., 1992). Therefore our finding of increased callosal area in the anterior sub-region of the CC may indicate a cortical abnormality in the prefrontal and/or temporoparietal association areas. The absence of significant findings in the CC1 area derived by radial partitioning, which connects the prefrontal association areas, indicates that the cortical abnormality may be localizable to the temporal association cortex. This finding is in keeping with the consistently reported left superior temporal gyrus (STG) volume reduction in positive symptom schizophrenia (review, Shenton et al., 2001). The finding of increased callosal area in the anterior truncus may indicate "abnormal functional hyperconnection" (David, 1993) involving these regions, in accordance with the notion that schizophrenia is caused by difficulties in filtering and selecting relevant information (Braff and Geyer, 1989). Therefore, the increased area limited to the anterior truncus of the CC may reflect the underlying cortical abnormality localizable to the left STG. This abnormal hyperconnection involving the temporal association cortices might result in reduced laterality underlying the positive symptoms of schizophrenia, as suggested by previous authors (Gruzelier, 1999). However, in view of the fact that all the schizophrenia subjects in our sample had severe psychosis at the time of the study, the possibility that this focal increase in CC area may be state-related also needs to be considered.

Alterations of white matter connectivity have been demonstrated in first episode schizophrenia using the relatively newer technique of Diffusion tensor imaging (DTI) (Federspiel et al., 2006). Moreover, Brambilla et al. (2005) demonstrated a link between positive symptoms and the anterior callosal apparent diffusion co-efficient (ADC) measures, in consonance with the findings of the present study.

Corpus callosal abnormalities are not restricted to schizophrenia, but have been reported in other psychiatric conditions such as autism (Hardan et al., 2006; Vidal et al., 2006), attention-deficit hyperactivity disorder (Hynd et al., 1991), dyslexia (Rumsey, 1996) and bipolar

affective disorder (Coffman et al., 1990). A variety of neurological conditions (review, Georgy et al., 1993) including multiple sclerosis, neoplastic disorders, vascular conditions, chronic alcoholism (Marchiafava-Bignami disease), fetal alcohol syndrome and HIV encephalopathy also involve structural abnormalities of the CC. The topography of corpus callosal atrophy has been suggested to provide an index of region-specific neuronal loss in the neocortex (Pantel et al., 1999; Hampel et al., 1998) in Alzheimer's disease. An integration of CC abnormalities in the various neuropsychiatric conditions and the corresponding behavioural features could pave the way for a better understanding of the connectivity disturbances underlying these disorders.

Females had significantly larger area of the whole CC as well as the anterior and middle callosal sub-regions in both the NRS and HS groups (Table 2). Reports of gender effects on CC morphology in schizophrenia have been inconsistent (Bishop and Wahlsten, 1997). Many previous studies have found a larger CC area relative to brain size in women, especially in the splenium or isthmus (Witelson, 1989; de Lacoste et al., 1990;). We did not find a significant gender*diagnosis interaction (Table 1) in agreement with two previous studies in first-episode schizophrenia patients (Bachmann et al., 2003; Keshavan et al., 2007). Our finding of increased area in females of whole CC as well as anterior and middle sub-regions in both NRS and HS groups provides strong support to the previous evidence for sexual dimorphism in CC areas, considering that we have attempted to control for most of the confounders, including handedness and TBV. The sexual dimorphism of CC area is commonly interpreted as reflecting increased inter-hemispheric connectivity underlying increased ambilaterality in females (McGlone, 1980; Jancke et al., 1997). It is well known that women with schizophrenia have better premorbid functioning, a more benign illness course, lower levels of disability and better integration into the community than men (Morgan et al., 2008). The callosal hyperconnection, along with the faster development of the CC noted in females may protect them against the 'misconnectivity phenomena' in the frontal lobes that males may encounter at a younger age, contributing to negative symptoms (Crow et al., 2007).

The major methodological advantage of the present study is that the schizophrenia subjects were neuroleptic-naïve. The schizophrenia sample was carefully selected to avoid possible confounding effects of gender, handedness, age, chronicity of illness, as well as clinical heterogeneity. Stringent rules were adopted to identify the mid-sagittal slice, resulting in very high inter-rater reliability in identifying the slice. Areas of CC sub-regions were computed using highly reliable geometric partitioning methods. We adjusted for total brain size by including TBV as a co-variate in group comparisons, thereby ruling out its effect on CC area. Nevertheless, the findings of the study should be interpreted in the light of certain methodological limitations which include modest sample sizes and non-equitable distribution of males and females in both NRS and HS samples (10:13). In this study on neuroleptic-naïve, recent-onset schizophrenia patients, limited power afforded by the modest sample sizes is indeed a limitation, as has been the case with similar studies in the past. It also needs to be pointed out that there were no inclusion/exclusion criteria stated *a priori* to limit the patients to only positive symptom schizophrenia subjects. However, post-hoc analysis of the NRS sample revealed that all the subjects had a rating of ≥ 3 on delusions and/or hallucinatory behavior on PANSS. Even though the patient sample comprised of recent-onset (≤ 5 years duration) schizophrenia subjects, there was substantial variance in the duration of illness (69.96 ± 63.35 weeks; range=6–260 weeks), with four patients having a duration of illness of less than 6 months (schizophreniform disorder) (Table 1). However, all the major findings persisted even on reanalysis of the data excluding these 4 subjects. Another limitation of CC partition studies is the lack of a one-to-one correlation between the various partitioning methods and the texture of the CC at the cellular level.

In summary, right-handed, NRS subjects all of whom had scores of ≥ 3 on delusions and/or hallucinatory behavior were found to have increased area of the anterior trunci sub-region of CC, in comparison to age-, sex-, and education-matched right-handed HS. This finding of increased area in a specific region of the CC that could contain fibres connecting the temporal association cortices is in consonance with information processing models of schizophrenia which suggest “abnormal functional hyperconnection” as underlying the positive symptoms. Additionally, female subjects were found to have higher areas of the whole CC as well as anterior and middle sub-regions when compared to males, in both HS and NRS samples. This finding could be reflective of a “normal hyperconnection” underlying increased ambilaterality in females. The findings of the study also highlight the importance of controlling for the various confounders affecting CC morphometry to obtain internally consistent results.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

References

- Aboitiz F, Montiel J. One hundred million years of interhemispheric communication: the history of the corpus callosum. *Braz J Med Biol Res* 2003;36:409–420. [PubMed: 12700818]
- Aboitiz F, Scheibel AB, Fisher RS, Zaidel E. Fiber composition of the human corpus callosum. *Brain Res* 1992;598:143–153. [PubMed: 1486477]
- Allen LS, Richey MF, Chai YM, Gorski RA. Sex differences in the corpus callosum of the living human being. *J Neurosci* 1991;11:933–942. [PubMed: 2010816]
- Anstey KJ, Mack HA, Christensen H, Li SC, Reglade-Meslin C, Maller J, Kumar R, Dear K, Eastel S, Sachdev P. Corpus callosum size, reaction time speed and variability in mild cognitive disorders and in a normative sample. *Neuropsychologia* 2007;45:1911–1920. [PubMed: 17240409]
- Bachmann S, Pantel J, Flender A, Bottmer C, Essig M, Schroder J. Corpus callosum in first-episode patients with schizophrenia--a magnetic resonance imaging study. *Psychol Med* 2003;33:1019–1027. [PubMed: 12946086]
- Bilder RM, Wu H, Bogerts B, Degreef G, Ashtari M, Alvir JM, Snyder PJ, Lieberman JA. Absence of regional hemispheric volume asymmetries in first-episode schizophrenia. *Am J Psychiatry* 1994;151:1437–1447. [PubMed: 8092337]
- Bishop KM, Wahlsten D. Sex differences in the human corpus callosum: myth or reality? *Neurosci Biobehav Rev* 1997;21:581–601. [PubMed: 9353793]
- Braff, D.; Geyer, M. Sensorimotor gating and the neurobiology of schizophrenia: Human and animal model studies. In: Schultz, S.; Tamminga, C., editors. *Schizophrenia: Scientific progress*. New York: Oxford University Press; 1989. p. 124-136.
- Brambilla P, Cerini R, Gasparini A, Versace A, Andreone N, Vittorini E, Barbui C, Pelizza L, Nose M, Barlocco L, Marrella G, Gregis M, Tournikioti K, David AS, Keshavan MS, Tansella M. Investigation of corpus callosum in schizophrenia with diffusion imaging. *Schizophr Res* 2005;79:201–210. [PubMed: 15953707]
- Casanova MF, Sanders RD, Goldberg TE, Bigelow LB, Christison G, Torrey EF, Weinberger DR. Morphometry of the corpus callosum in monozygotic twins discordant for schizophrenia: a magnetic resonance imaging study. *J Neurol Neurosurg Psychiatry* 1990;53:416–421. [PubMed: 2351971]
- Chua SE, Sharma T, Takei N, Murray RM, Woodruff PW. A magnetic resonance imaging study of corpus callosum size in familial schizophrenic subjects, their relatives, and normal controls. *Schizophr Res* 2000;41:397–403. [PubMed: 10728717]
- Coffman JA, Bornstein RA, Olson SC, Schwarzkopf SB, Nasrallah HA. Cognitive impairment and cerebral structure by MRI in bipolar disorders. *Biol Psychiatry* 1990;27:1188–1196. [PubMed: 2354225]
- Clarke JM, Zaidel E. Anatomical-behavioral relationships: corpus callosum morphometry and hemispheric specialization. *Behav Brain Res* 1994;64:185–202. [PubMed: 7840886]

- Clarke S, Kraftsik R, Van der Loos H, Innocenti GM. Forms and measures of adult and developing human corpus callosum: is there sexual dimorphism? *J Comp Neurol* 1989;280:213–230. [PubMed: 2925893]
- Cowell PE, Kertesz A, Denenberg VH. Multiple dimensions of handedness and the human corpus callosum. *Neurology* 1993;43:2353–2357. [PubMed: 8232955]
- Crow TJ, Paez P, Chance SA. Callosal misconnectivity and the sex difference in psychosis. *Int Rev Psychiatry* 2007;19:449–457. [PubMed: 17671877]
- Crow TJ, Poulter M, Lofthouse R, Chen G, Shah T, Bass N, Morganti C, Vita A, Smith C, Boccio-Smith A, et al. Male siblings with schizophrenia share alleles at the androgen receptor above chance expectation. *Am J Med Genet* 1993;48:159–160. [PubMed: 8291571]
- David AS. Callosal transfer in schizophrenia: too much or too little? *J Abnorm Psychol* 1993;102:573–579. [PubMed: 8282926]
- de Lacoste MC, Adesanya T, Woodward DJ. Measures of gender differences in the human brain and their relationship to brain weight. *Biol Psychiatry* 1990;28:931–942. [PubMed: 2275951]
- de Lacoste MC, Kirkpatrick JB, Ross ED. Topography of the human corpus callosum. *J Neuropathol Exp Neurol* 1985;44:578–591. [PubMed: 4056827]
- DeLisi LE, Sakuma M, Kushner M, Finer DL, Hoff AL, Crow TJ. Anomalous cerebral asymmetry and language processing in schizophrenia. *Schizophr Bull* 1997;23:255–271. [PubMed: 9165636]
- DeLisi LE, Tew W, Xie S, Hoff AL, Sakuma M, Kushner M, Lee G, Shedlack K, Smith AM, Grimson R. A prospective follow-up study of brain morphology and cognition in first-episode schizophrenic patients: preliminary findings. *Biol Psychiatry* 1995;38:349–360. [PubMed: 8547454]
- Falkai P, Schneider T, Greve B, Klieser E, Bogerts B. Reduced frontal and occipital lobe asymmetry on the CT-scans of schizophrenic patients. Its specificity and clinical significance. *J Neural Transm Gen Sect* 1995;99:63–77. [PubMed: 8579808]
- Federspiel A, Begre S, Kiefer C, Schroth G, Strik WK, Dierks T. Alterations of white matter connectivity in first episode schizophrenia. *Neurobiology of Disease* 2006;22:702–709. [PubMed: 16624566]
- Frumin M, Golland P, Kikinis R, Hirayasu Y, Salisbury DF, Hennen J, Dickey CC, Anderson M, Jolesz FA, Grimson WE, McCarley RW, Shenton ME. Shape differences in the corpus callosum in first-episode schizophrenia and first-episode psychotic affective disorder. *Am J Psychiatry* 2002;159:866–868. [PubMed: 11986146]
- Georgy BA, Hesselink JR, Jernigan TL. MR imaging of the corpus callosum. *AJR Am J Roentgenol* 1993;160:949–955. [PubMed: 8470609]
- Goghari VM, Lang DJ, Flynn SW, Mackay AL, Honer WG. Smaller corpus callosum subregions containing motor fibers in schizophrenia. *Schizophr Res* 2005;73:59–68. [PubMed: 15567078]
- Gruzeliier J. Functional neuro-psychophysiological asymmetry in schizophrenia: a review and reorientation. *Schizophr Bull* 1999;25:91–120. [PubMed: 10098916]
- Gunther W, Petsch R, Steinberg R, Moser E, Streck P, Heller H, Kurtz G, Hippus H. Brain dysfunction during motor activation and corpus callosum alterations in schizophrenia measured by cerebral blood flow and magnetic resonance imaging. *Biol Psychiatry* 1991;29:535–555. [PubMed: 1905162]
- Guy, W. Superintendent of Documents. Washington, DC: US Government Printing Office; 1976. ECDEU Assessment Manual for Psychopharmacology.
- Hampel H, Teipel SJ, Alexander GE, Horwitz B, Teichberg D, Schapiro MB, Rapoport SI. Corpus callosum atrophy is a possible indicator of region- and cell type-specific neuronal degeneration in Alzheimer disease: a magnetic resonance imaging analysis. *Arch Neurol* 1998;55:193–198. [PubMed: 9482361]
- Hardan AY, Muddasani S, Vemulapalli M, Keshavan MS, Minshew NJ. An MRI study of increased cortical thickness in autism. *Am J Psychiatry* 2006;163:1290–1292. [PubMed: 16816240]
- Hauser P, Dauphinais ID, Berrettini W, DeLisi LE, Gelernter J, Post RM. Corpus callosum dimensions measured by magnetic resonance imaging in bipolar affective disorder and schizophrenia. *Biol Psychiatry* 1989;26:659–668. [PubMed: 2804188]
- Highley JR, Esiri MM, McDonald B, Cortina-Borja M, Herron BM, Crow TJ. The size and fibre composition of the corpus callosum with respect to gender and schizophrenia: a post-mortem study. *Brain* 1999;122(Pt 1):99–110. [PubMed: 10050898]

- Hofer S, Frahm J. Topography of the human corpus callosum revisited--comprehensive fiber tractography using diffusion tensor magnetic resonance imaging. *Neuroimage* 2006;32:989–994. [PubMed: 16854598]
- Hoff AL, Neal C, Kushner M, DeLisi LE. Gender differences in corpus callosum size in first-episode schizophrenics. *Biol Psychiatry* 1994;35:913–919. [PubMed: 8080890]
- Holloway RL, Anderson PJ, Defendini R, Harper C. Sexual dimorphism of the human corpus callosum from three independent samples: relative size of the corpus callosum. *Am J Phys Anthropol* 1993;92:481–498. [PubMed: 8296877]
- Hynd GW, Clikeman MS, Lorys AR, Novay CS, Eliopoulos D, Lyytinen H. Corpus callosum morphology in attention deficit hyperactivity disorders: morphometric analysis of MRI. *J Learning Disabil* 1991;24:141–146.
- Jacobsen LK, Giedd JN, Rajapakse JC, Hamburger SD, Vaituzis AC, Frazier JA, Lenane MC, Rapoport JL. Quantitative magnetic resonance imaging of the corpus callosum in childhood onset schizophrenia. *Psychiatry Res* 1997;68:77–86. [PubMed: 9104755]
- Jancke L, Staiger JF, Schlaug G, Huang Y, Steinmetz H. The relationship between corpus callosum size and forebrain volume. *Cereb Cortex* 1997;7:48–56. [PubMed: 9023431]
- John JP, Yashavantha BS, Gado M, Veena R, Jain S, Ravishankar S, Csernansky JG. A proposal for MRI-based parcellation of the frontal pole. *Brain Struct Funct* 2007;212:245–253. [PubMed: 17929054]
- Johnson SC, Farnworth T, Pinkston JB, Bigler ED, Blatter DD. Corpus callosum surface area across the human adult life span: effect of age and gender. *Brain Res Bull* 1994;35:373–377. [PubMed: 7850489]
- Keshavan MS, Diwadkar VA, Harenski K, Rosenberg DR, Sweeney JA, Pettegrew JW. Abnormalities of the corpus callosum in first episode, treatment naive schizophrenia. *J Neurol Neurosurg Psychiatry* 2002;72:757–760. [PubMed: 12023420]
- Keshavan MS, Prasad KM, Pearlson G. Are brain structural abnormalities useful as endophenotypes in schizophrenia? *Int Rev Psychiatry* 2007;19:397–406. [PubMed: 17671872]
- LaMantia AS, Rakic P. Axon overproduction and elimination in the corpus callosum of the developing rhesus monkey. *J Neurosci* 1990;10:2156–2175. [PubMed: 2376772]
- Machiyama, Y.; Watanabe, Y.; Machiyama, R. Neuroanatomical studies of the corpus callosum in schizophrenia: evidence of aberrant interhemispheric fibre connection. In: Takahashi, R.; Flor-Henry, P.; Gluzelier, J., editors. *Cerebral Dynamics, Laterality and Psychopathology*. New York: Elsevier; 1987. p. 411–412.
- Matthew RJ, Partain CL, Prakash R, Kulkarni MV, Logan TP, Wilson WH. A study of the septum pellucidum and corpus callosum in schizophrenia with MR imaging. *Acta Psychiatr Scand* 1985;72:414–421. [PubMed: 4091025]
- McCarley RW, Wible CG, Frumin M, Hirayasu Y, Levitt JJ, Fischer IA, Shenton ME. MRI anatomy of schizophrenia. *Biol Psychiatry* 1999;45:1099–1119. [PubMed: 10331102]
- McGlone J. Sex differences in functional brain asymmetry. *Cortex* 1978;14:122–128. [PubMed: 16295117]
- McGuire PK, Frith CD. Disordered functional connectivity in schizophrenia. *Psychol Med* 1996;26:663–667. [PubMed: 8817700]
- Morgan VA, Castle DJ, Jablensky AV. Do women express and experience psychosis differently from men? Epidemiological evidence from the Australian National Study of Low Prevalence (Psychotic) Disorders. *Aust N Z J Psychiatry* 2008;42:74–82. [PubMed: 18058447]
- Narr KL, Thompson PM, Sharma T, Moussai J, Cannestra AF, Toga AW. Mapping morphology of the corpus callosum in schizophrenia. *Cereb Cortex* 2000;10:40–49. [PubMed: 10639394]
- Nasrallah HA, Andreasen NC, Coffman JA, Olson SC, Dunn VD, Ehrhardt JC, Chapman SM. A controlled magnetic resonance imaging study of corpus callosum thickness in schizophrenia. *Biol Psychiatry* 1986;21:274–282. [PubMed: 3947709]
- Pandya DN, Karol EA, Heilbronn D. The topographical distribution of interhemispheric projections in the corpus callosum of the rhesus monkey. *Brain Res* 1971;32:31–43. [PubMed: 5000193]
- Pantel J, Schroder J, Jauss M, Essig M, Minakaran R, Schonknecht P, Schneider G, Schad LR, Knopp MV. Topography of callosal atrophy reflects distribution of regional cerebral volume reduction in Alzheimer's disease. *Psychiatry Res Neuroimaging* 1999;90:181–192.

- Preuss UW, Meisenzahl EM, Frodl T, Zetzsche T, Holder J, Leinsinger G, Hegerl U, Hahn K, Moller HJ. Handedness and corpus callosum morphology. *Psychiatry Res* 2002;116:33–42. [PubMed: 12426032]
- Price G, Bagary MS, Cercignani M, Altmann DR, Ron MA. The corpus callosum in first episode schizophrenia: a diffusion tensor imaging study. *J Neurol Neurosurg Psychiatry* 2005;76:585–587. [PubMed: 15774453]
- Raine A, Harrison GN, Reynolds GP, Sheard C, Cooper JE, Medley I. Structural and functional characteristics of the corpus callosum in schizophrenics, psychiatric controls, and normal controls. A magnetic resonance imaging and neuropsychological evaluation. *Arch Gen Psychiatry* 1990;47:1060–1064. [PubMed: 2241507]
- Rajarethinam RP, DeQuardo JR, Nalepa R, Tandon R. Superior temporal gyrus in schizophrenia: a volumetric magnetic resonance imaging study. *Schizophr Res* 2000;41:303–312. [PubMed: 10708339]
- Reinartz SJ, Coffman CE, Smoker WR, Godersky JC. MR imaging of the corpus callosum: normal and pathologic findings and correlation with CT. *AJR Am J Roentgenol* 1988;151:791–798. [PubMed: 3262282]
- Robb RA, Hanson DP, Karwoski RA, Larson AG, Workman EL, Stacy MC. Analyze: a comprehensive, operator-interactive software package for multidimensional medical image display and analysis. *Comput Med Imaging Graph* 1989;13:433–454. [PubMed: 2688869]
- Rosenthal R, Bigelow LB. Quantitative brain measurements in chronic schizophrenia. *Br J Psychiatry* 1972;121:259–264. [PubMed: 5073779]
- Rossell SL, Shapleske J, Fukuda R, Woodruff PW, Simmons A, David AS. Corpus callosum area and functioning in schizophrenic patients with auditory--verbal hallucinations. *Schizophr Res* 2001;50:9–17. [PubMed: 11378310]
- Rossi A, Stratta P, Gallucci M, Passariello R, Casacchia M. Quantification of corpus callosum and ventricles in schizophrenia with nuclear magnetic resonance imaging: a pilot study. *Am J Psychiatry* 1989;146:99–101. [PubMed: 2912255]
- Rumsey JM, Casanova M, Mannheim GB, Patronas N, De Vaughn N, Hamburger SD, Aquino T. Corpus callosum morphology, as measured with MRI, in dyslexic men. *Biol Psychiatry* 1996;39:769–775. [PubMed: 8731517]
- Shenton ME, Dickey CC, Frumin M, McCarley RW. A review of MRI findings in schizophrenia. *Schizophr Res* 2001;49:1–52. [PubMed: 11343862]
- Steinmetz H, Jancke L, Kleinschmidt A, Schlaug G, Volkmann J, Huang Y. Sex but no hand difference in the isthmus of the corpus callosum. *Neurology* 1992;42:749–752. [PubMed: 1565226]
- Steinmetz H, Staiger JF, Schlaug G, Huang Y, Jancke L. Corpus callosum and brain volume in women and men. *Neuroreport* 1995;6:1002–1004. [PubMed: 7632882]
- Sunderland S. The projection of the cerebral cortex on the pons and cerebellum in the macaque monkey. *J Anat* 1940;74:201–226. [PubMed: 17104808]
- Talairach, J.; Tournoux, P. Referentially oriented cerebral MRI anatomy: Atlas of stereotaxic anatomical correlations for gray and white matter. New York: Thieme Medical publishers, Inc; 1993.
- Tan YL, Chen BH, Yang JD, Zhang J, Wang YC, Chai SH, Wang ZY, Li QH. Localization of functional projections from corpus callosum to cerebral cortex. *Chin Med J (Engl)* 1991;104:851–857. [PubMed: 1752144]
- Teipel SJ, Alexander GE, Schapiro MB, Moller HJ, Rapoport SI, Hampel H. Age-related cortical grey matter reductions in non-demented Down's syndrome adults determined by MRI with voxel-based morphometry. *Brain* 2004;127:811–824. [PubMed: 14985261]
- Tibbo P, Nopoulos P, Arndt S, Andreasen NC. Corpus callosum shape and size in male patients with schizophrenia. *Biol Psychiatry* 1998;44:405–412. [PubMed: 9777169]
- Uematsu M, Kaiya H. The morphology of the corpus callosum in schizophrenia. An MRI study. *Schizophr Res* 1988;1:391–398. [PubMed: 3154526]
- Vidal CN, Nicolson R, DeVito TJ, Hayashi KM, Geaga JA, Drost DJ, Williamson PC, Rajakumar N, Sui Y, Dutton RA, Toga AW, Thompson PM. Mapping corpus callosum deficits in autism: an index of aberrant cortical connectivity. *Biol Psychiatry* 2006;60:218–225. [PubMed: 16460701]

- Weis S, Jellinger K, Wenger E. Morphometry of the corpus callosum in normal aging and Alzheimer's disease. *J Neural Transm Suppl* 1991;33:35–38. [PubMed: 1753249]
- Witelson SF. Hand and sex differences in the isthmus and genu of the human corpus callosum. A postmortem morphological study. *Brain* 1989;112(Pt 3):799–835. [PubMed: 2731030]
- Witelson SF, Goldsmith CH. The relationship of hand preference to anatomy of the corpus callosum in men. *Brain Res* 1991;545:175–182. [PubMed: 1860044]
- Woodruff PW, McManus IC, David AS. Meta-analysis of corpus callosum size in schizophrenia. *J Neurol Neurosurg Psychiatry* 1995;58:457–461. [PubMed: 7738554]
- Woodruff PW, Pearlson GD, Geer MJ, Barta PE, Chilcoat HD. A computerized magnetic resonance imaging study of corpus callosum morphology in schizophrenia. *Psychol Med* 1993;23:45–56. [PubMed: 8475214]
- Woodruff PW, Phillips ML, Rushe T, Wright IC, Murray RM, David AS. Corpus callosum size and inter-hemispheric function in schizophrenia. *Schizophr Res* 1997;23:189–196. [PubMed: 9075296]

Figure 1.
Corpus callosum partitioning schemes (intended for reproduction in black and white)

Figure 2.

Scatterplots depicting the distribution of areas of (a) the anterior third (CC-I) and (b) anterior truncus (CC-1) of the corpus callosum in schizophrenia and healthy subjects

Table 1

Sociodemographic and clinical characteristics of the study samples

Socio-demographic and illness characteristics	Subjects (N=46)	
	Schizophrenia [n=23]	Controls [n=23]
Gender		
Male	10(43.5%)	10 (43.5%)
Female	13 (56.5%)	13 (56.5%)
Age (years)	30.13 (s.d.=5.79); range: 20–40	30.13 (s.d.=5.05); range: 20–39
Education		
Illiterate/ Primary education	11	11
Secondary/Pre-university	09	09
Graduate/Post-graduate	03	03
Diagnosis		
Paranoid schizophrenia	08	
Undifferentiated schizophrenia	11	
Schizophreniform disorder	04	
Age of onset of illness (years)	28.96 (s.d.=5.44); range: 20–40	NA
Duration of illness (weeks)	69.96 (s.d.=63.354); range: 6–260 weeks	
PANSS positive total score	24.61 (4.49)	
PANSS negative total score	25.04 (6.33)	
PANSS global psychopathology total score	46.48 (8.98)	
PANSS total score	96.13 (15.46)	
CGI score	5.00 (0.74)	

Table 2
Comparison of whole CC area and areas of sub-regions between schizophrenia subjects and healthy subjects

	Mean area (mm ²) (s.d)		Group		Gender		Group × gender	
	Schiz (n=23)	Control (n=23)	F	Sig.	F	Sig.	F	Sig.
Whole CC	663.33(73.82)	662.88(77.59)	0.612	0.439	18.774*	.000	.081	.777
CC-I (Anterior)	274.07(30.50)	259.88(36.92)	7.027*	.012	16.231*	.000	.045	.834
CC-II (Middle)	153.50(25.62)	150.57(24.62)	1.133	.294	15.822*	.000	.261	.613
CC-III (Posterior)	235.76(30.51)	252.43(29.27)	2.952	.094	6.157*	.018	.650	.425
CC1 (Rostrum and genu)	194.51(20.58)	193.73(24.50)	.608	.440	13.313*	.001	.015	.904
CC2 (Anterior truncus)	111.00(16.00)	100.09(18.42)	14.807*	.000	18.828*	.000	.039	.844
CC3 (Middle truncus)	91.80(15.41)	87.89(18.16)	1.800	.188	11.831*	.001	.196	.660
CC4 (Posterior truncus)	92.49(21.04)	99.04(18.98)	.973	.330	6.178*	.017	.587	.448
CC5 (Isthmus & splenium)	173.01(20.94)	181.79(22.40)	.926	.342	3.105	.086	.171	.681

Table 3
Gender-wise comparison of whole CC areas and areas of sub-regions in healthy subjects and schizophrenia patients

	Healthy Controls				Schizophrenia			
	Mean area (mm ²)		F	Sig.	Mean area (mm ²)		F	Sig.
	Males (n=10)	Females (n=13)			Males (n=10)	Females (n=13)		
Cc whole	637.22(90.78)	682.62(62.32)	9.492*	.006	613.44(42.21)	694.04(73.23)	10.338*	.005
CC-I(Anterior)	253.93 (42.82)	264.46 (32.74)	9.689*	.006	257.25 (23.19)	284.42 (30.53)	10.220*	.005
CC-II(Middle)	137.38(28.25)	160.71(16.01)	12.711*	.002	139.31(18.73)	162.23(25.94)	4.771*	.043
CC-III(Posterior)	245.93(29.07)	257.44(29.57)	1.394	.252	216.88(15.90)	247.38(31.95)	5.003*	.039
C1(Rostrum and genu)	185.50(23.90)	200.44(23.66)	11.414*	.003	181.97(16.52)	202.23(19.42)	5.497*	.031
C2(Anterior truncus)	97.73(21.50)	101.90(16.34)	6.234*	.022	102.84(11.32)	116.02(16.75)	19.781*	.000
C3 (Middle truncus)	79.60 (21.45)	94.27 (12.57)	9.316*	.007	82.94(14.41)	97.25(13.78)	3.571	.076
C4(Posteriortruncus)	94.98 (20.15)	102.17(18.21)	1.912	.183	80.91(17.03)	99.62(20.59)	4.259	.055
C5 (Isthmus andsplenium)	179.40(25.28)	183.63(20.79)	0.456	.508	164.22(15.27)	178.42(22.62)	3.276	.088