Calcium dynamics integrated into signalling pathways that influence vertebrate axial patterning

Christina M. Freisinger, Igor Schneider, Trudi A. Westfall
and Diane C. Slusarski*

Department of Biology, University of Iowa, Iowa City, IA 52242, USA

Many aspects of animal development including fertilization as well as organ formation and function are dependent upon the dynamic release of calcium (Ca^{2+}) ions. Although the controlled release and/or accumulation of Ca^{2+} ions has been extensively studied, how the release dynamics produce a specific biological output in embryonic development is less clear. We will briefly summarize Ca^{2+} sources, highlight data on endogenous Ca^{2+} release in vertebrate embryos relevant to body plan formation and cell movement, and integrate pharmacological and molecular-genetic studies to lend insight into the signalling pathways involved. Finally, based on *in vivo* imaging in zebrafish genetic mutants, we will put forward the model that distinct Ca^{2+} release dynamics lead to antagonism of the developmentally important Wnt/ β -catenin signalling pathway, while sustained Ca^{2+} release modulates cell polarization or directed migration.

Keywords: calcium; Wnt; β -catenin; zebrafish; dorsal–ventral axis; left–right patterning

1. CALCIUM SOURCES

Although critical for many processes, calcium ions (Ca^{2+}) are not metabolized by the cell. Instead, Ca^{2+} enters the cell across either the plasma membrane or the membrane of intracellular organelles. Depending on the location of the ion channels and the extent and duration of the channel opening, local or global changes in Ca^{2+} levels in the cytosol of the cell can result. Intricate crosstalk and feedback between release circuits can stimulate Ca^{2+} -induced Ca^{2+} release, influencing neighbouring receptors and potentially triggering a regenerative wave (Berridge 1997; Berridge *et al.* 2003; Roderick *et al.* 2003). In addition, continued stimulation and/or depletion of endoplasmic reticulum (ER) stores activate a store-operated Ca^{2+} entry influx pathway located at the plasma membrane (Parekh & Putney 2005).

In non-excitable (non-neuronal) cells, a majority of intracellular Ca^{2+} release occurs through inositol 1,4,5-trisphosphate (IP_3)-sensitive Ca^{2+} channels present in the ER membrane (reviewed in Berridge *et al.* 2003). The phosphatidylinositol (PI) cycle is activated in response to many hormones and growth factors that bind to cell surface receptors. Two predominant receptor classes are the G-protein-coupled receptor class and the receptor tyrosine kinase class. Extracellular ligand stimulation of these receptors activates a PI-specific phospholipase C (PLC). Activated PLC converts membrane-bound phosphatidylinositol (4,5)-bisphosphate (PIP_2) into IP_3 and lipophilic diacylglycerol (DAG). IP_3 subsequently binds to receptors (IP_3R) located principally

on the ER triggering the rapid release of Ca^{2+} into the cytosol of the cell. At the same time, DAG produced by PIP_2 hydrolysis can act as an additional second messenger to further activate downstream targets such as protein kinase C (PKC).

Relevant to this discussion is the fact that Ca^{2+} release is heterogeneous. Specific cellular responses can be triggered by differences in the amplitude, frequency and duration of intracellular Ca^{2+} oscillations. Such oscillations can be derived from changes in upstream steps within the PI cycle, such as G-protein activity, PLC activity and IP_3 levels (Hirose *et al.* 1999; Luo *et al.* 2001; McCarron *et al.* 2004; Thore *et al.* 2004; Nomikos *et al.* 2005; Rey *et al.* 2005). Oscillatory small molecules such as IP_3 may be transmitted to other cells via gap junctions (Lin *et al.* 2004), a phenomenon that may be of significance in the regulation of axis induction in the zebrafish blastula (see below). Feedback from activated Ca^{2+} -binding proteins adds another layer of complexity to the dynamics of Ca^{2+} release and removal. For example, IP_3R activity integrates signals from small molecules and proteins, including PKC and Ca^{2+} /calmodulin-dependent protein kinase II (CaMKII; Nadif Kasri *et al.* 2002; Assefa *et al.* 2004; Patterson *et al.* 2004).

2. CALCIUM AND THE VERTEBRATE BODY PLAN

After fertilization, the next major developmental programme involves the establishment of the primary axes, in which regions of the embryo receive signals to determine the cells that will contribute to the dorsal (back) or ventral (belly) tissue as well as anterior (head/top) and posterior (tail/bottom) regions. A number of studies have linked PI-cycle activity with

* Author for correspondence (diane-slusarski@uiowa.edu).

One contribution of 11 to a Discussion Meeting Issue 'Calcium signals and developmental patterning'.

body plan formation. Classical work using lithium, an inhibitor of inositol turnover (Berridge *et al.* 1989), induced expansion of dorsal structures in *Xenopus* (Kao *et al.* 1986; Kao & Elinson 1989, 1998), and similar effects were obtained in the zebrafish (*Danio rerio*) embryo (Stachel *et al.* 1993; Aanstad & Whitaker 1999). Lithium-induced embryonic defects are rescued by supplying an intermediate of the PI cycle, *myo*-inositol (Busa & Gimlich 1989). In further support, zebrafish and *Xenopus* embryos injected with antibodies that disrupt IP₃R function displayed expanded dorsal structures with the loss of ventral structures (Kume *et al.* 1997; Westfall *et al.* 2003b). Treatment of zebrafish embryos with inhibitors that target different steps in the PI cycle generated dorsalized or axis-duplication phenotypes (figure 1a, inset). The dorsalization defects generated by the use of pharmacological inhibitors that target inositol turnover in the zebrafish were likewise rescued by *myo*-inositol (Westfall *et al.* 2003b). The findings of an involvement for PI-cycle activity in axis induction are consistent with the observed spontaneous increase in IP₃ levels in the *Xenopus* embryo at the blastula stage (Busa & Gimlich 1989; Maslanski *et al.* 1992). Moreover, *in vivo* imaging of calcium release dynamics in the zebrafish embryo identified rapid aperiodic Ca²⁺ release that persists until the midblastula transition stage (Reinhard *et al.* 1995; Slusarski *et al.* 1997b; Slusarski & Corces 2000). The idea that increased IP₃ levels may trigger Ca²⁺ release during these stages has been corroborated by drug inhibition studies (Slusarski *et al.* 1997a). The effects of lithium were most pronounced when exposure occurred on the ventral side of the embryo, suggesting that in the embryo PI-cycle activity is normally high on the ventral side and low on the dorsal side. Consistent with this is the described role of activated CaMKII for ventral fates in *Xenopus* embryos (Kühl *et al.* 2000a).

3. THE WNT SIGNALLING NETWORK

The Wnt family of growth factors and components of their signalling pathways have diverse roles in development and disease. Wnt signalling influences many aspects of embryonic patterning, cell proliferation as well as the maintenance and differentiation of stem cells, and is critical in axis formation (figure 1b; reviewed in Moon *et al.* 2004; Kohn & Moon 2005; Clevers 2006). In the absence of the so-called canonical Wnt signalling (Wnt/β-catenin), β-catenin is rapidly sequestered in a cytoplasmic degradation complex containing axin, the adenomatous polyposis tumour suppressor protein (APC) and the serine threonine kinase GSK-3β. GSK-3 phosphorylation of β-catenin targets the latter for proteasomal degradation (figure 1b). Wnt binding to its co-receptors Frizzled and LRP5/6 (low-density lipoprotein receptor-related protein) activates a cytoplasmic phosphoprotein (Dishevelled, Dsh) which downregulates GSK-3 and inhibits the degradation of β-catenin. Stabilized β-catenin protein interacts with the members of the LEF/TCF transcription factor family in the nucleus to promote the activation of downstream target genes involved in axis specification.

Another endogenous target of lithium is the β-catenin degradation complex component GSK-3, which when inhibited promotes dorsal axis induction (Klein & Melton 1996; Stambolic *et al.* 1996). The fact that exogenous *myo*-inositol can suppress the effects of GSK-3 inhibition (Hedgepeth *et al.* 1997) further supports the notion of communication between PI-cycle activity and Wnt/β-catenin signalling to regulate axis induction.

The Wnt network has layers of complexity including the fact that different Wnt ligands can activate distinct cellular outputs. In vertebrate embryos, overexpression of a subset of Wnts induces hyperdorsalization and ectopic axes by virtue of increased Wnt/β-catenin signalling activity (Moon *et al.* 1993b; Moon & Kimelman 1998). Additional Wnts (including *Wnt*-5, -4 and -11) appear to act independently of β-catenin function (Dale 1998; Kühl *et al.* 2000b). Emerging evidence suggests that the ability of Wnt ligands to activate different signalling pathways, β-catenin dependent (canonical) and β-catenin independent (non-canonical), appears to be controlled by the timing of expression and receptor context, not to mention the correct combination of intracellular effectors. In the zebrafish embryo, *Wnt*-5 overexpression results in an increase in the frequency of intracellular Ca²⁺ release in a manner that is dependent on G-protein activity and the PI cycle (Slusarski *et al.* 1997a,b), thus linking Wnt activity to IP₃-dependent Ca²⁺ release and defining the Wnt/Ca²⁺ signalling pathway.

The concept that Wnt/Ca²⁺, either in parallel or as part of a complex signalling network, appears to interact with the Wnt/β-catenin pathway in early axis specification was initially suggested by the apparent antagonism of certain pairs of Wnt ligands when expressed in *Xenopus* and zebrafish embryos (Moon *et al.* 1993b; Slusarski *et al.* 1997b). Expression of ligands that activate Wnt/β-catenin signalling in these embryos, such as *Wnt*-8, results in ectopic axis induction. However, when *Wnt*-5 is co-expressed with *Wnt*-8, the *Wnt*-8 axis induction phenotype is suppressed. Stimulating Ca²⁺ release, via activated serotonin receptor, also antagonizes *Wnt*-8-induced expansion of dorsal domains (Slusarski *et al.* 1997b), supporting that *Wnt*-5 antagonism of Wnt/β-catenin is mediated by Ca²⁺ release. On the other hand, pharmacological or genetic reduction of the Wnt/Ca²⁺ pathway in zebrafish embryos or mouse limb buds generates ectopic accumulation of nuclear β-catenin and activation of β-catenin transcriptional targets (Topol *et al.* 2003; Westfall *et al.* 2003a,b). Additionally, inhibition of G-protein function dorsalizes *Xenopus* embryos (Kume *et al.* 2000). These observations are consistent with a model in which IP₃-dependent Ca²⁺ release, promoted by Wnt/Ca²⁺ signalling activity, negatively regulate the Wnt/β-catenin signalling pathway and therefore axis induction (figure 1b).

Various studies have shown that there are common components between the Wnt/Ca²⁺ and the planar cell polarity pathways (Wnt/PCP), another non-canonical Wnt pathway involved in the polarization of cells in *Drosophila* and vertebrate species (Kohn & Moon 2005; Solnica-Krezel 2005). During gastrulation, vertebrate

Figure 1. Signal transduction pathways in vertebrate axis formation. During embryogenesis, the basic body plan generates a dorsal-ventral and anterior-posterior orientation of tissues and organs. (a) Lateral view of a wild-type zebrafish 48 hour larvae with normal axial patterning with anterior to the right and dorsal to the top. Key tissues are noted. Inset: a two-headed embryo as a result of PI-cycle inhibition. (b) A simplified schematic of the Wnt signalling network. β -catenin-independent pathways are encased in green region and β -catenin-dependent components are in the yellow region. β -cat, β -catenin. (c) A single ratiometric image taken from a time course during the cellular blastoderm stage of a fura-2-injected zebrafish embryo. The image is pseudo-coloured to represent high calcium levels as warm colouring (yellow) and low calcium levels as blue. Arrows denote transient fluxes observed in the enveloping layer and the arrowhead designates the yolk syncytial layer region. (d) *hecate* mutant embryo lacking dorsal-anterior structures such as the eyes and brain reflective of a ventralized phenotype. (e) *ppt* mutant embryo with a shortened anterior-posterior axis and kinked tail.

embryos undergo a variety of morphogenetic movements to thicken (dorsal convergence) and elongate (axis extension) the embryo, also called convergence extension (CE; Keller 2002). Ca^{2+} mobilization associated with waves of tissue contraction can be observed in *Xenopus* explants (Wallingford *et al.* 2001a). In intact zebrafish embryos, intercellular Ca^{2+} waves have been observed at the margin during gastrulation (Gilland *et al.* 1999). The relationship between Ca^{2+} waves and cell movement is supported by the finding that, in *Xenopus* embryos, pharmacological inhibition of such waves results in CE defects without altering cell fate (Wallingford *et al.* 2001a,b). Wnt genes that result in the activation of Ca^{2+} release in the blastula embryo, such as *Wnt-5* (Slusarski *et al.* 1997b; Westfall *et al.* 2003a), can also alter morphogenetic movements later during gastrulation when misexpressed (Moon *et al.* 1993a; Ungar & Moon 1995). Core components involved in the polarization of epithelial cells in the *Drosophila* cuticle are also necessary for the polarization of migrating cells during vertebrate gastrulation (Solnica-Krezel 2005). Indeed, several core components of Wnt/PCP have been shown to activate Ca^{2+} release in zebrafish including Frizzled-2, Dsh and the intracellular protein Prickle (Slusarski *et al.* 1997a; Sheldahl *et al.* 2003; Veeman *et al.* 2003). The fact that interference with either Ca^{2+} release or Wnt/PCP signalling results in CE defects suggests non-canonical Wnt signalling activity can be characterized as a complex network with cellular outputs defined by Ca^{2+} modulation and polarized cell movement (figure 1b).

4. CALCIUM DYNAMICS AND BIOLOGICAL OUTPUTS

The Ca^{2+} activity in the zebrafish blastula is observed in the enveloping layer (EVL) and yolk syncytial layer (YSL; figure 1c; Reinhard *et al.* 1995; Slusarski *et al.* 1997b; Slusarski & Corces 2000). Although present in overlapping stages, the EVL-specific Ca^{2+} fluxes are present in a cell or small cluster of cells lasting for short intervals. The YSL-specific Ca^{2+} , on the other hand, displays sustained elevation in a population of cells. The distinct dynamics of Ca^{2+} increases and the bimodal role of Wnt/ Ca^{2+} , in β -catenin antagonism and polarized cell movement, led us to hypothesize that the rapid aperiodic Ca^{2+} release is coupled to Wnt/ β -catenin antagonism and the sustained Ca^{2+} levels integrates into polarizing cells or their directed migration.

Support for this theory comes from the analysis of Ca^{2+} release dynamics in zebrafish mutants. A mutation in the zebrafish maternal gene *hecate* produces ventralized embryos (figure 1d) that lack nuclear β -catenin (Lyman Gingerich *et al.* 2005). Consistent with our theory, these mutant embryos display increased Ca^{2+} release frequency in the EVL (Lyman Gingerich *et al.* 2005). Suppression of the Ca^{2+} dynamics with pharmacological reagents was sufficient to rescue the defects in dorsal cell fate specification observed in these mutants (Lyman Gingerich *et al.* 2005), highlighting the relationship between Ca^{2+} release and β -catenin antagonism in a genetic context.

In the zebrafish, *Wnt-5* has been shown to correspond to the genetic mutation *pipetail* (*ppt*; Rauch *et al.* 1997). The *ppt* zygotic mutants have axis

extension defects, reflected in a shorter anterior–posterior length and kinks in the tail, resembling a pipe (figure 1e; Hammerschmidt *et al.* 1996; Kilian *et al.* 2003). Analysis of zygotic *ppt* mutant embryos revealed reduced Ca^{2+} levels in the YSL region (Westfall *et al.* 2003a). Suggestive that reduced Ca^{2+} levels are central to the *ppt* defects is the ability to rescue the mutant phenotype with the expression of activated CaMKII (Westfall *et al.* 2003a). These data raise the possibility that the YSL-specific Ca^{2+} dynamics contributes to polarized cell movements during vertebrate gastrulation.

5. CALCIUM AND LEFT–RIGHT ASYMMETRY

There are many cases in development where a signalling cassette is used in multiple processes. Thus, it would be predicted that transient Ca^{2+} release will correlate with β -catenin antagonism or sustained Ca^{2+} signalling with cell polarization at additional stages or in different tissues. Section 6 describes the orientation of organs relative to the body axis and the implications of both transient and sustained Ca^{2+} modulations in this process. Although vertebrates appear bilaterally symmetrical from the outside, the heart, lungs, liver and gut are carefully positioned across the left–right (LR) axis. The development of this asymmetry is highly conserved across species and most likely will use conserved signalling molecules.

Embryonic organ laterality is preceded by molecular and physiological asymmetries. Shown in figure 2a is a schematic of developmental structures and gene products implicated in LR patterning with a focus on zebrafish. At the morphological level, non-involuting dorsal mesoderm cells, the dorsal forerunner cells (DFCs), migrate ahead of the dorsal blastoderm during gastrulation (figure 2b, arrow; Cooper & D'Amico 1996; Melby *et al.* 1996). At the start of gastrulation, these cells express key signalling molecules such as nodal-related *squint*, a Brachyury homologue *no tail* and *left–right dynein* (*lrd1*; Schulte-Merker *et al.* 1994; Feldman *et al.* 1998; Essner *et al.* 2002). The DFCs are highly endocytotic and readily take up vital dyes (Cooper & D'Amico 1996) allowing for the visualization of their migration to the tailbud region where they undergo morphogenesis and form a ciliated structure, the Kupffer's vesicle (KV), during early somite stages (figure 2c; Cooper & D'Amico 1996; Melby *et al.* 1996). Ablation of the DFCs or mechanical disruption of the KV disrupts LR patterning (Amack & Yost 2004; Essner *et al.* 2005). Endogenous Ca^{2+} release activity in and around the DFC region during epiboly suggests striking similarities to the transient activity both in a cell or a few cells (figure 2d, arrow) and in a region of sustained high Ca^{2+} (figure 2d, arrowhead) previously described during the cellular blastoderm stages (figure 1c; Schneider *et al.* 2008). Whether this activity correlates with β -catenin antagonism and polarized cell movement bears further study.

The LR signals have been proposed to be modulated by gap junctional communication and/or asymmetrical H^+/K^+ -ATPase expression during *Xenopus* early cleavage (reviewed in Levin 2005).

Figure 2. Left-right pattern formation in the zebrafish. (a) Schematic noting key structures and gene products implicated in LR patterning. (b) Epiboly-staged embryo oriented with the animal pole towards the top, yolk region at the bottom and the dorsal shield region in the centre. The arrow notes the DFCs migrating ahead of the shield region. (c) Vital-dye-labelled DFCs (with Syto-11) at the early somite stages. The combined bright field and fluorescent image focused on the tailbud showing the DFCs forming the KV in the black box. The midline of the embryo can be distinguished by the compact notochord cells above the KV. (d) A ratiometric image taken from a time course of an epiboly-staged fura-2-injected embryo. Arrow notes a transient flux while the arrowhead notes a region of high sustained Ca^{2+} . (e) Lateral view of a somite-stage embryo at a time molecular asymmetric markers begin to be expressed. The KV is morphologically visible in the tail (black box). (f) Whole-mount *in situ* hybridization at approximately 22 somite stage with the *southpaw* and *no tail* probes. A dorsal view, anterior to the top, shows left-sided expression of *southpaw* and midline expression of *no tail*. (g) Whole-mount *in situ* hybridization of a 48 hours post-fertilization larva with cardiac myosin light chain to illustrate heart morphogenesis. The arrow points to the heart tube.

However, conserved molecular asymmetries in vertebrates become apparent only during later somite stages after the KV/node has formed (figure 2e). Of note is the conserved left-sided expression of the secreted transforming growth factor- β (TGF- β) related factor *nodal* (reviewed in Ahmad et al. 2004). In zebrafish, the *nodal*-related gene *southpaw* (*spaw*; Long et al. 2003) is the earliest asymmetric marker expressed in the left lateral plate mesoderm (figure 2f). Downstream targets of *southpaw*, including *lefty1* and *pitx2*, also have conserved asymmetric expression in vertebrates (reviewed in Hamada et al. 2002; Wright & Halpern 2002). *lefty1* contributes to a barrier at the midline of the embryo, preventing the left-sided signal from propagating to the right, and *pitx2* contributes to the subsequent morphogenetic

changes in the organs, such as looping to form the chambers of the heart (figure 2g).

Asymmetric Ca^{2+} levels across the mouse and chick node have been implicated in LR axis determination (McGrath et al. 2003; Raya et al. 2004). Elevated Ca^{2+} is thought to act via an unknown mechanism to induce left-sided gene expression in concert with or independent of nodal activity (Brennan et al. 2002; Hashimoto et al. 2004; Marques et al. 2004). Rotation of monocilia generates a leftward fluid flow in the node (Nonaka et al. 1998; Okada et al. 1999) and has been proposed to stimulate mechanosensory cilia and trigger elevated intracellular Ca^{2+} levels at the left edge of the mouse node (McGrath et al. 2003). The zebrafish KV contains monociliated cells similar to those found in the mouse node (Supp et al. 1997, 1999;

Brueckner 2001; Hashimoto *et al.* 2004; Essner *et al.* 2005) and these cilia have been shown to beat in the same direction, possibly establishing Ca^{2+} asymmetries (Kramer-Zucker *et al.* 2005). Indeed, an intracellular Ca^{2+} flux with a left-sided bias near the zebrafish KV has been detected and is proposed to be required for normal LR patterning (Sarmah *et al.* 2005). In an alternate model, leftward flow of vesicular particles containing *sonic hedgehog* (*shh*) acts in a distinct atypical signalling pathway to activate Ca^{2+} on the left side of the node (Tanaka *et al.* 2005).

In contrast to intracellular Ca^{2+} release, a role for left-sided elevation of extracellular Ca^{2+} has also been proposed. In the chick node, extracellular Ca^{2+} levels appear to be higher transiently on the left side, although it is unclear whether intracellular Ca^{2+} is also increased. This asymmetry was abolished after a treatment with omeprazole, an inhibitor of H^+/K^+ ATPase, which also caused LR defects. This led to the proposal that differential H^+/K^+ ATPase activity during gastrulation sets up a spatial gradient of extracellular Ca^{2+} , which is subsequently transduced through Notch to activate asymmetric gene expression (Raya *et al.* 2004). In zebrafish, pharmacological disruption of H^+/K^+ ATPase activity leads to LR asymmetry defects (Kawakami *et al.* 2005) and early H^+/K^+ ATPase inhibition disrupts cilia number and length in the KV (Adams *et al.* 2006). However, immunodetection of H^+/K^+ ATPase did not display obvious asymmetries in zebrafish (Kawakami *et al.* 2005). Moreover, the H^+/K^+ ATPase inhibitor used, omeprazole, also induces ectopic *shh* expression in chick (Raya *et al.* 2004). Although it is unclear whether extracellular Ca^{2+} regulation is conserved in other organisms, mouse embryos that are homozygous for mutations in the *Polycystin-2* (*Pkd2*) gene, a Ca^{2+} -permeable cation-selective channel, exhibit loss of asymmetric Ca^{2+} levels across the node and LR defects (McGrath *et al.* 2003). Thus, several vertebrate models support a role for Ca^{2+} signalling in the establishment of LR asymmetry, but many questions and issues remain to be addressed, such as the Ca^{2+} sources, the Ca^{2+} -dependent responders and the mechanism linking the Ca^{2+} flux in and around the node to asymmetry in the lateral plate mesoderm. Also, it has not been investigated whether there are multiple stages of Ca^{2+} -dependent events involving the polarized migration and coalescence of the DFCs into a KV, initiation of asymmetric expression and the subsequent maintenance of laterality signals.

6. CONCLUSIONS AND FUTURE DIRECTIONS

In conclusion, progression from egg to embryo requires a dynamic interaction of signal transduction networks. The continuous exchange of information between cells, through both direct contact and diffusible molecules, influences gene expression and cell behaviour. *In vivo* imaging studies are a critical step in the comprehensive analysis of Ca^{2+} signalling in development. Coupling *in vivo* imaging with molecular, genetic or pharmacological tools will determine the mechanism by which Ca^{2+} signalling is modulated and interpreted in the

embryo. Future studies will reconstruct the spatial and temporal dynamics of Ca^{2+} release and incorporate this activity into known signalling pathways, thus providing the capacity to discern the true nature of the cellular basis of pattern formation. Knowledge about how intricate Ca^{2+} signals are integrated into developmental pathways, in particular the Wnt growth factors, raises the possibility to address the pathophysiology of diseases that result from misregulation of this network.

Animal care and experiments were performed in accordance with institutional guidelines for animal experiment ethics.

We are grateful to our collaborators who have contributed to the research reviewed here. We thank Dr Pelegri and Dr Rebagliati as well as members of the Slusarski lab for their insightful discussions. This work was supported by the National Institutes of Health to D.C.S. and American Heart Association predoctoral fellowships to C.M.F. and I.S.

REFERENCES

- Aanstad, P. & Whitaker, M. 1999 Predictability of dorso-ventral asymmetry in the cleavage stage zebrafish embryo: an analysis using lithium sensitivity as a dorso-ventral marker. *Mech. Dev.* **88**, 33–41. (doi:10.1016/S0925-4773(99)00171-9)
- Adams, D. S., Robinson, K. R., Fukumoto, T., Yuan, S., Albertson, R. C., Yelick, P., Kuo, L., McSweeney, M. & Levin, M. 2006 Early, H^+ -V-ATPase-dependent proton flux is necessary for consistent left–right patterning of non-mammalian vertebrates. *Development* **133**, 1657–1671. (doi:10.1242/dev.02341)
- Ahmad, N., Long, S. & Rebagliati, M. 2004 A southpaw joins the roster: the role of the zebrafish *nodal-related* gene *southpaw* in cardiac LR asymmetry. *Trends Cardiovasc. Med.* **14**, 43–49. (doi:10.1016/j.tcm.2003.11.001)
- Amack, J. D. & Yost, H. J. 2004 The T box transcription factor no tail in ciliated cells controls zebrafish left–right asymmetry. *Curr. Biol.* **14**, 685–690. (doi:10.1016/j.cub.2004.04.002)
- Assefa, Z. *et al.* 2004 Caspase-3-induced truncation of type 1 inositol trisphosphate receptor accelerates apoptotic cell death and induces inositol trisphosphate-independent calcium release during apoptosis. *J. Biol. Chem.* **279**, 43 227–43 236. (doi:10.1074/jbc.M403872200)
- Berridge, M. J. 1997 The AM and FM of calcium signalling. *Nature* **386**, 759–760. (doi:10.1038/386759a0)
- Berridge, M., Downes, C. & Hanley, M. 1989 Neural and developmental actions of lithium: a unifying hypothesis. *Cell* **59**, 411–419. (doi:10.1016/0092-8674(89)90026-3)
- Berridge, M. J., Bootman, M. D. & Roderick, H. L. 2003 Calcium signalling: dynamics, homeostasis and remodelling. *Nat. Rev. Mol. Cell Biol.* **4**, 517–529. (doi:10.1038/nrm1155)
- Brennan, J., Norris, D. P. & Robertson, E. J. 2002 Nodal activity in the node governs left–right asymmetry. *Genes Dev.* **16**, 2339–2344. (doi:10.1101/gad.1016202)
- Brueckner, M. 2001 Cilia propel the embryo in the right direction. *Am. J. Med. Genet.* **101**, 339–344. (doi:10.1002/1096-8628(20010715)101:4<339::AID-AJMG1442>3.0.CO;2-P)
- Busa, W. & Gimlich, R. 1989 Lithium-induced teratogenesis in frog embryos prevented by a polyphosphoinositide cycle intermediate or a diacylglycerol analog. *Dev. Biol.* **132**, 315–324. (doi:10.1016/0012-1606(89)90228-5)
- Clevers, H. 2006 Wnt/beta-catenin signaling in development and disease. *Cell* **127**, 469–480. (doi:10.1016/j.cell.2006.10.018)

- Cooper, M. S. & D'Amico, L. A. 1996 A cluster of noninvoluting endocytic cells at the margin of the zebrafish blastoderm marks the site of embryonic shield formation. *Dev. Biol.* **180**, 184–198. (doi:10.1006/dbio.1996.0294)
- Dale, T. 1998 Signal transduction by the Wnt family of ligands. *Biochem. J.* **329**, 209–223.
- Essner, J. J., Vogan, K. J., Wagner, M. K., Tabin, C. J., Yost, H. J. & Brueckner, M. 2002 Conserved function for embryonic nodal cilia. *Nature* **418**, 37–38. (doi:10.1038/418037a)
- Essner, J. J., Amack, J. D., Nyholm, M. K., Harris, E. B. & Yost, H. J. 2005 Kupffer's vesicle is a ciliated organ of asymmetry in the zebrafish embryo that initiates left–right development of the brain, heart and gut. *Development* **132**, 1247–1260. (doi:10.1242/dev.01663)
- Feldman, B., Gates, M. A., Egan, E. S., Dougan, S. T., Rennebeck, G., Sirotkin, H. I., Schier, A. F. & Talbot, W. S. 1998 Zebrafish organizer development and germ-layer formation require nodal-related signals [see comments]. *Nature* **395**, 181–185. (doi:10.1038/26013)
- Gilland, E., Miller, A. L., Karplus, E., Baker, R. & Webb, S. E. 1999 Imaging of multicellular large-scale rhythmic calcium waves during zebrafish gastrulation. *Proc. Natl Acad. Sci. USA* **96**, 157–161. (doi:10.1073/pnas.96.1.157)
- Hamada, H., Meno, C., Watanabe, D. & Saijoh, Y. 2002 Establishment of vertebrate left–right asymmetry. *Nat. Rev. Genet.* **3**, 103–113. (doi:10.1038/nrg732)
- Hammerschmidt, M. et al. 1996 Mutations affecting morphogenesis during gastrulation and tail formation in the zebrafish, *Danio rerio*. *Development* **123**, 143–151.
- Hashimoto, H., Rebagliati, M., Ahmad, N., Muraoka, O., Kurokawa, T., Hibi, M. & Suzuki, T. 2004 The Cerberus/Dan-family protein Charon is a negative regulator of Nodal signaling during left–right patterning in zebrafish. *Development* **131**, 1741–1753. (doi:10.1242/dev.01070)
- Hedgepeth, C. M., Conrad, L. J., Zhang, J., Huang, H.-C., Lee, V. M. & Klein, P. S. 1997 Activation of the Wnt signaling pathway: a molecular mechanism for lithium action. *Dev. Biol.* **185**, 82–91. (doi:10.1006/dbio.1997.8552)
- Hirose, K., Kadowaki, S., Tanabe, M., Takeshima, H. & Iino, M. 1999 Spatiotemporal dynamics of inositol 1,4,5-trisphosphate that underlies complex Ca^{2+} mobilization patterns. *Science* **284**, 1527–1530. (doi:10.1126/science.284.5419.1527)
- Kao, K. R. & Elinson, R. P. 1989 Dorsalization of mesoderm induction by lithium. *Dev. Biol. (Orl.)* **132**, 81–90. (doi:10.1016/0012-1606(89)90207-8)
- Kao, K. R. & Elinson, R. P. 1998 The legacy of lithium effects on development. *Biol. Cell* **90**, 585–590. (doi:10.1016/S0248-4900(99)80016-1)
- Kao, K., Masui, Y. & Elinson, R. 1986 Lithium-induced respecification of pattern in *Xenopus laevis* embryos. *Nature* **322**, 371–373. (doi:10.1038/322371a0)
- Kawakami, Y., Raya, A., Raya, R. M., Rodriguez-Esteban, C. & Belmonte, J. C. 2005 Retinoic acid signalling links left–right asymmetric patterning and bilaterally symmetric somitogenesis in the zebrafish embryo. *Nature* **435**, 165–171. (doi:10.1038/nature03512)
- Keller, R. 2002 Shaping the vertebrate body plan by polarized embryonic cell movements. *Science* **298**, 1950–1954. (doi:10.1126/science.1079478)
- Kilian, B., Mansukoski, H., Barbosa, F. C., Ulrich, F., Tada, M. & Heisenberg, C. P. 2003 The role of Ppt/Wnt5 in regulating cell shape and movement during zebrafish gastrulation. *Mech. Dev.* **120**, 467–476. (doi:10.1016/S0925-4773(03)00004-2)
- Klein, P. S. & Melton, D. A. 1996 A molecular mechanism for the effect of lithium on development. *Proc. Natl Acad. Sci. USA* **93**, 8455–8459. (doi:10.1073/pnas.93.16.8455)
- Kohn, A. D. & Moon, R. T. 2005 Wnt and calcium signaling: β -catenin-independent pathways. *Cell Calcium* **38**, 439–446. (doi:10.1016/j.ceca.2005.06.022)
- Kramer-Zucker, A. G., Olale, F., Haycraft, C. J., Yoder, B. K., Schier, A. F. & Drummond, I. A. 2005 Cilia-driven fluid flow in the zebrafish pronephros, brain and Kupffer's vesicle is required for normal organogenesis. *Development* **132**, 1907–1921. (doi:10.1242/dev.01772)
- Kühl, M., Sheldahl, L. C., Malbon, C. C. & Moon, R. T. 2000a Ca^{2+} /calmodulin-dependent protein kinase II is stimulated by Wnt and frizzled homologs and promotes ventral cell fates in *Xenopus*. *J. Biol. Chem.* **275**, 12 701–12 711. (doi:10.1074/jbc.275.17.12701)
- Kühl, M., Sheldahl, L. C., Park, M., Miller, J. R. & Moon, R. T. 2000b The Wnt/ Ca^{2+} pathway—a new vertebrate Wnt signaling pathway takes shape. *Trends Genet.* **16**, 279–283. (doi:10.1016/S0168-9525(00)02028-X)
- Kume, S., Muto, A., Inoue, T., Suga, K., Okano, H. & Mikoshiba, K. 1997 Role of inositol 1,4,5-trisphosphate receptor in ventral signaling in *Xenopus* embryos. *Science* **278**, 1940–1943. (doi:10.1126/science.278.5345.1940)
- Kume, S., Saneyoshi, T. & Mikoshiba, K. 2000 Desensitization of IP₃-induced Ca^{2+} release by overexpression of a constitutively active Gq α protein converts ventral to dorsal fate in *Xenopus* early embryos. *Dev. Growth Differ.* **42**, 327–335. (doi:10.1046/j.1440-169x.2000.00519.x)
- Levin, M. 2005 Left–right asymmetry in embryonic development: a comprehensive review. *Mech. Dev.* **122**, 3–25. (doi:10.1016/j.mod.2004.08.006)
- Lin, G. C., Rurangirwa, J. K., Koval, M. & Steinberg, T. H. 2004 Gap junctional communication modulates agonist-induced calcium oscillations in transfected HeLa cells. *J. Cell Sci.* **117**, 881–887. (doi:10.1242/jcs.00942)
- Long, S., Ahmad, N. & Rebagliati, M. 2003 The zebrafish nodal-related gene *southpaw* is required for visceral and diencephalic left–right asymmetry. *Development* **130**, 2303–2316. (doi:10.1242/dev.00436)
- Luo, X., Popov, S., Bera, A. K., Wilkie, T. M. & Muallem, S. 2001 RGS proteins provide biochemical control of agonist-evoked $[\text{Ca}^{2+}]_i$ oscillations. *Mol. Cell* **7**, 651–660. (doi:10.1016/S1097-2765(01)00211-8)
- Lyman Gingerich, J., Westfall, T. A., Slusarski, D. C. & Pelegri, F. 2005 *hecate*, a zebrafish maternal effect gene, affects dorsal organizer induction and intracellular calcium transient frequency. *Dev. Biol.* **286**, 427–439. (doi:10.1016/j.ydbio.2005.07.031)
- Marques, S., Borges, A. C., Silva, A. C., Freitas, S., Cordenonsi, M. & Belo, J. A. 2004 The activity of the Nodal antagonist Cerl-2 in the mouse node is required for correct L/R body axis. *Genes Dev.* **18**, 2342–2347. (doi:10.1101/gad.306504)
- Maslanski, J., Leshko, L. & Busa, W. 1992 Lithium-sensitive production of inositol phosphates during amphibian embryonic mesoderm induction. *Science* **256**, 243–245. (doi:10.1126/science.1314424)
- McCarron, J. G., MacMillan, D., Bradley, K. N., Chalmers, S. & Muir, T. C. 2004 Origin and mechanisms of Ca^{2+} waves in smooth muscle as revealed by localized photolysis of caged inositol 1,4,5-trisphosphate. *J. Biol. Chem.* **279**, 8417–8427. (doi:10.1074/jbc.M311797200)
- McGrath, J., Somlo, S., Makova, S., Tian, X. & Brueckner, M. 2003 Two populations of node monocilia initiate left–right asymmetry in the mouse. *Cell* **114**, 61–73. (doi:10.1016/S0092-8674(03)00511-7)
- Melby, A. E., Warga, R. M. & Kimmel, C. B. 1996 Specification of cell fates at the dorsal margin of the zebrafish gastrula. *Development* **122**, 2225–2237.
- Moon, R. T. & Kimelman, D. 1998 From cortical rotation to organizer gene expression: toward a molecular explanation

- of axis specification in *Xenopus*. *Bioessays* **20**, 536–545. (doi:10.1002/(SICI)1521-1878(199807)20:7<536::AID-BIES4>3.0.CO;2-I)
- Moon, R. T., Campbell, R. M., Christian, J. L., McGrew, L. L., Shih, J. & Fraser, S. 1993a Xwnt-5A: a maternal Wnt that affects morphogenetic movements after over-expression in embryos of *Xenopus laevis*. *Development* **119**, 97–111.
- Moon, R. T., Christian, J. L., Campbell, R. M., McGrew, L. L., DeMarais, A. A., Torres, M., Lai, C.-J., Olson, D. J. & Kelly, G. M. 1993b Dissecting Wnt signalling pathways and Wnt-sensitive developmental processes through transient misexpression analyses in embryos of *Xenopus laevis*. *Dev. Suppl.* 85–94.
- Moon, R. T., Kohn, A. D., De Ferrari, G. V. & Kaykas, A. 2004 WNT and beta-catenin signalling: diseases and therapies. *Nat. Rev. Genet.* **5**, 691–701. (doi:10.1038/nrg1427)
- Nadif Kasri, N., Bultynck, G., Sienaert, I., Callewaert, G., Erneux, C., Missiaen, L., Parys, J. B. & De Smedt, H. 2002 The role of calmodulin for inositol 1,4,5-trisphosphate receptor function. *Biochim. Biophys. Acta* **1600**, 19–31.
- Nomikos, M., Blayney, L. M., Larman, M. G., Campbell, K., Rossbach, A., Saunders, C. M., Swann, K. & Lai, F. A. 2005 Role of phospholipase C-zeta domains in Ca^{2+} -dependent phosphatidylinositol 4,5-bisphosphate hydrolysis and cytoplasmic Ca^{2+} oscillations. *J. Biol. Chem.* **280**, 31 011–31 018. (doi:10.1074/jbc.M500629200)
- Nonaka, S., Tanaka, Y., Okada, Y., Takeda, S., Harada, A., Kanai, Y., Kido, M. & Hirokawa, N. 1998 Randomization of left–right asymmetry due to loss of nodal cilia generating leftward flow of extraembryonic fluid in mice lacking KIF3B motor protein. *Cell* **95**, 829–837. (doi:10.1016/S0092-8674(00)81705-5)
- Okada, Y., Nonaka, S., Tanaka, Y., Saijoh, Y., Hamada, H. & Hirokawa, N. 1999 Abnormal nodal flow precedes situs inversus in *iv* and *inv* mice. *Mol. Cell* **4**, 459–468. (doi:10.1016/S1097-2765(00)80197-5)
- Parekh, A. B. & Putney Jr, J. W. 2005 Store-operated calcium channels. *Physiol. Rev.* **85**, 757–810. (doi:10.1152/physrev.00057.2003)
- Patterson, R. L., Boehning, D. & Snyder, S. H. 2004 Inositol 1,4,5-trisphosphate receptors as signal integrators. *Annu. Rev. Biochem.* **73**, 437–465. (doi:10.1146/annurev.biochem.73.071403.161303)
- Rauch, G. J., Hammerschmidt, M., Blader, P., Schauerte, H. E., Strahle, U., Ingham, P. W., McMahon, A. P. & Hafter, P. 1997 Wnt5 is required for tail formation in the zebrafish embryo. *Cold Spring Harb. Symp. Quant. Biol.* **62**, 227–234.
- Raya, A., Kawakami, Y., Rodriguez-Esteban, C., Ibanes, M., Rasskin-Gutman, D., Rodriguez-Leon, J., Buscher, D., Feijo, J. A. & Izpisua Belmonte, J. C. 2004 Notch activity acts as a sensor for extracellular calcium during vertebrate left–right determination. *Nature* **427**, 121–128. (doi:10.1038/nature02190)
- Reinhard, E., Yokoe, H., Niebling, K. R., Allbritton, N. L., Kuhn, M. A. & Meyer, T. 1995 Localized calcium signals in early zebrafish development. *Dev. Biol. (Orl.)* **170**, 50–61. (doi:10.1006/dbio.1995.1194)
- Rey, O., Young, S. H., Yuan, J., Slice, L. & Rozengurt, E. 2005 Amino acid-stimulated Ca^{2+} oscillations produced by the Ca^{2+} -sensing receptor are mediated by a phospholipase C/inositol 1,4,5-trisphosphate-independent pathway that requires G12, Rho, filamin-A, and the actin cytoskeleton. *J. Biol. Chem.* **280**, 22 875–22 882. (doi:10.1074/jbc.M503455200)
- Roderick, H. L., Berridge, M. J. & Bootman, M. D. 2003 Calcium-induced calcium release. *Curr. Biol.* **13**, R425. (doi:10.1016/S0960-9822(03)00358-0)
- Sarmah, B., Latimer, A. J., Appel, B. & Wente, S. R. 2005 Inositol polyphosphates regulate zebrafish left–right asymmetry. *Dev. Cell* **9**, 133–145. (doi:10.1016/j.devcel.2005.05.002)
- Schneider, I., Houston, D. W., Rebagliati, M. R. & Slusarski, D. C. 2008 Calcium fluxes in dorsal forerunner cells antagonize β -catenin and alter left–right patterning. *Development* **135**, 75–84. (doi:10.1242/dev.004713)
- Schulte-Merker, S., van Eeden, F. J., Halpern, M. E., Kimmel, C. B. & Nusslein-Volhard, C. 1994 No tail (*ntl*) is the zebrafish homologue of the mouse T (Brachyury) gene. *Development* **120**, 1009–1015.
- Sheldahl, L. C., Slusarski, D. C., Pandur, P., Miller, J. R., Kühl, M. & Moon, R. T. 2003 Dishevelled activates Ca^{2+} flux, PKC, and CamKII in vertebrate embryos. *J. Cell Biol.* **161**, 769–777. (doi:10.1083/jcb.200211094)
- Slusarski, D. C. & Corces, V. G. 2000 Calcium imaging in cell–cell signaling. In *Developmental biology protocols*, vol. 1 (eds R. S. Tuan & C. W. Lo), pp. 253–261. Totowa, NJ: Humana Press.
- Slusarski, D. C., Corces, V. G. & Moon, R. T. 1997a Interaction of Wnt and a Frizzled homologue triggers G-protein-linked phosphatidylinositol signalling. *Nature* **390**, 410–413. (doi:10.1038/37138)
- Slusarski, D. C., Yang-Snyder, J., Busa, W. B. & Moon, R. T. 1997b Modulation of embryonic intracellular Ca^{2+} signaling by Wnt-5A. *Dev. Biol.* **185**, 114–120. (doi:10.1006/dbio.1996.8463)
- Solnica-Krezel, L. 2005 Conserved patterns of cell movements during vertebrate gastrulation. *Curr. Biol.* **15**, R213–R228. (doi:10.1016/j.cub.2005.03.016)
- Stachel, S., Grunwald, D. & Meyers, P. 1993 Lithium perturbation and *goosecoid* expression identify a dorsal specification pathway in the pregastrula zebrafish. *Development* **117**, 1261–1274.
- Stambolic, V., Ruel, L. & Woodgett, J. R. 1996 Lithium inhibits glycogen synthase kinase-3 activity and mimics wingless signalling in intact cells [published erratum appears in *Curr. Biol.* 1997 Mar 1;7(3):196]. *Curr. Biol.* **6**, 1664–1668. (doi:10.1016/S0960-9822(02)70790-2)
- Supp, D. M., Witte, D. P., Potter, S. S. & Brueckner, M. 1997 Mutation of an axonemal dynein affects left–right asymmetry in *inversus viscerum* mice. *Nature* **389**, 963–966. (doi:10.1038/40140)
- Supp, D. M., Brueckner, M., Kuehn, M. R., Witte, D. P., Lowe, L. A., McGrath, J., Corrales, J. & Potter, S. S. 1999 Targeted deletion of the ATP binding domain of left–right dynein confirms its role in specifying development of left–right asymmetries. *Development* **126**, 5495–5504.
- Tanaka, Y., Okada, Y. & Hirokawa, N. 2005 FGF-induced vesicular release of Sonic hedgehog and retinoic acid in leftward nodal flow is critical for left–right determination. *Nature* **435**, 172. (doi:10.1038/nature03494)
- Thore, S., Dyachok, O. & Tengholm, A. 2004 Oscillations of phospholipase C activity triggered by depolarization and Ca^{2+} influx in insulin-secreting cells. *J. Biol. Chem.* **279**, 19 396–19 400. (doi:10.1074/jbc.C400088200)
- Topol, L., Jiang, X., Choi, H., Garrett-Beal, L., Carolan, P. J. & Yang, Y. 2003 Wnt-5a inhibits the canonical Wnt pathway by promoting GSK-3-independent β -catenin degradation. *J. Cell Biol.* **162**, 899–908. (doi:10.1083/jcb.200303158)
- Ungar, A. R. & Moon, R. T. 1995 *Wnt4* affects morphogenesis when misexpressed in the zebrafish embryo. *Mech. Dev.* **52**, 153–164. (doi:10.1016/0925-4773(95)00386-F)
- Veeman, M. T., Slusarski, D. C., Kaykas, A., Louie, S. H. & Moon, R. T. 2003 Zebrafish *prickle*, a modulator of

- noncanonical Wnt/fz signaling, regulates gastrulation movements. *Curr. Biol.* **13**, 680–685. (doi:10.1016/S0960-9822(03)00240-9)
- Wallingford, J. B., Ewald, A. J., Harland, R. M. & Fraser, S. E. 2001a Calcium signaling during convergent extension in *Xenopus*. *Curr. Biol.* **11**, 652–661. (doi:10.1016/S0960-9822(01)00201-9)
- Wallingford, J. B., Vogeli, K. M. & Harland, R. M. 2001b Regulation of convergent extension in *Xenopus* by Wnt5a and Frizzled-8 is independent of the canonical Wnt pathway. *Int. J. Dev. Biol.* **45**, 225–227.
- Westfall, T. A., Brimeyer, R., Twedt, J., Gladon, J., Olberding, A., Furutani-Seiki, M. & Slusarski, D. C. 2003a Wnt-5/*pipetail* functions in vertebrate axis formation as a negative regulator of Wnt/ β -catenin activity. *J. Cell Biol.* **162**, 889–898. (doi:10.1083/jcb.200303107)
- Westfall, T. A., Hjertos, B. & Slusarski, D. C. 2003b Requirement for intracellular calcium modulation in zebrafish dorsal–ventral patterning. *Dev. Biol.* **259**, 380–391. (doi:10.1016/S0012-1606(03)00209-4)
- Wright, C. V. & Halpern, M. E. 2002 Specification of left–right asymmetry. *Results Probl. Cell Differ.* **40**, 96–116.