Published in final edited form as:

Org Lett. 2008 June 5; 10(11): 2307–2310. doi:10.1021/ol8008705.

Low Temperature Organocopper-Mediated Two-Component Cross Coupling/Cycloisomerization Approach Towards N-Fused Heterocycles

Dmitri Chernyak, Surendra Babu Gadamsetty, and Vladimir Gevorgyan*

Department of Chemistry, University of Illinois at Chicago, 845 West Taylor Street Chicago, Illinois 60607-7061

Abstract

Organocopper reagents smoothly react with heterocyclic propargyl mesylates at low temperature to produce N-fused heterocycles. The copper reagent plays a “double duty” in this novel cascade transformation, which proceeds via an S_N2' substitution followed by a subsequent cycloisomerization step.

The transition metal-catalyzed cycloisomerization approach is a powerful tool for the assembly of diverse heterocyclic frameworks.¹ We have developed the base-assisted Cu-catalyzed cycloisomerization of conjugated alkynylpyridines **1** into monosubstituted N-fused heterocycles **2**² (eq 1). Later, our group and others reported alternative methods for cycloisomerization of skipped propargyl imines **3** into 1,2 or 1,3-disubstituted N-fused heterocycles **4**³ (eq 2). Although both approaches are quite general with respect to the heterocyclic cores, they still have certain boundaries. Thus, the former method is restricted to the synthesis of heterocycles which do not possess substituents at the C-1 and C-2 positions, whereas the latter approach is limited to the oxygen-based substituents at C-1 (OR^2), as these propargylic precursors **3** are normally prepared from the corresponding heterocyclic aldehydes. Accordingly, development of more general methodologies toward multisubstituted N-fused heterocyclic scaffolds is desirable. Herein, we wish to report the copper-mediated coupling/cyclization cascade reaction of propargyl mesylates **5** toward N-fused heterocyclic frameworks **6** (eq 3). Not only does this method give access to C-1 alkyl- or aryl-substituted heterocycles, unavailable by existing cycloisomerization techniques, but it also holds a promise for a quick assembly of libraries of heterocyclic compounds via the two-component coupling/cycloisomerization strategy.

(eq 1)

(eq 2)

(eq 3)

It was suggested that the copper-catalyzed cycloisomerization of **1** into the monosubstituted heterocycles **2** proceeds via a 1,3-disubstituted allenyl intermediate *i* (eq 1).² Accordingly, we aimed at the analogous 1,1,3-trisubstituted allenyl intermediates *en route* to 1,3-disubstituted N-fused heterocyclic cores. Likely, these reactive allenyl intermediates **8** can be generated *in situ* via the S_N2' -substitution⁴ of the corresponding propargyl esters **5** (eq 4).

LG = living group

(eq 4)

We chose to employ the organocopper nucleophiles,⁵ based on the reasoning that, potentially, the copper reagent (or the copper by-product) can also mediate the subsequent cycloisomerization step of **v** into the heterocycle **6**, thus playing a “double duty” in this transformation.⁶

To this end, a possible substitution/cycloisomerization cascade of pyridyl-containing propargyl alcohol derivatives **5** with various copper reagents⁷ has been examined (Table 1). It was found that the higher-order alkyl cuprate reagents were not efficient in this transformation (entries 1-3). In contrast, the lower-order cyanocuprate (MeCu(CN)Li) reacted with mesylate **5** very smoothly, affording the C-1 alkylated indolizine derivative **6a** in 90% yield (entry 5)! Attempts to substitute mesylate with another leaving group⁸ were not successful (entries 6-12).

Short optimization indicated that employment of a phenyl copper reagent was reasonably efficient for the construction of C-1 phenyl-substituted indolizine **6** (entry 16). Importantly, in all cases, the corresponding allenes **8** were detected at early stages of the reaction, thus confirming for the first time that the cycloisomerizations of propargyl imines, indeed, proceed via allenic intermediates *i*^{2,3} (eq 1) and **8** (eq 4).

Next, the generality of the substitution/cycloisomerization cascade of differently substituted propargyl mesylates **5** was examined (Table 2). Thus, secondary alkyl-substituted mesylates **5a** and **5b** underwent smooth cyclization upon treatment with primary- (**7a-c**), secondary- (**7d**), and tertiary (**7e**) lower-order cyanocuprates to produce the corresponding indolizines **6** in good to excellent yields. (Table 2, entries 1-7). Noteworthy, primary mesylate **5d** can also be employed in this reaction giving access to monosubstituted indolizine **6ga**, a heterocycle, which can be further elaborated at C-3 via a direct C-H functionalization protocol.⁹ The secondary aryl-substituted mesylates **5e-i** were also efficient in this cyclization producing C-3 arylated indolizines in good yields. Notably, ester (entry 10), chloro- (entry 11), cyano- (entry 12), and nitro- (entry 13) functionalities were perfectly tolerated under these reaction conditions. It was also shown that pyrroloquinoline (entries 15, 16) and pyrroloisoquinoline (entry 17) cores can efficiently be assembled via this protocol. As discussed above, phenyl copper-mediated cyclization of **5a** produced C-1 aryl-substituted indolizine **6af** in 71% yield. Likewise, application of the same reagent to cyclization of **5j** led to formation of arylated pyrroloquinoline **6jf**, albeit in 51% yield (entry 16, Table 2).

We also investigated efficacy of a possible stepwise approach toward N-fused heterocycles **6** (Table 3). It was found that addition of cuprate reagents to the mesylates **5** at -78°C in one hour led to formation of the allenes **8** in varying yields (27-89%).¹⁰ The next step, cycloisomerization of allenes **8** under the standard conditions,^{2a} was even less efficient, producing trace to low yields of indolizines **6** (Table 3). Evidently, due to low stability of the pyridyl allenes **8** and low yields of their cyclization under the catalytic cyclization conditions,^{2a} the two-step process is much less efficient compared to the cascade substitution/cycloisomerization process (Table 2).

We propose the following mechanism for the substitution/cycloisomerization cascade of propargyl mesylates **5** with copper reagents into N-fused heterocycles **6** (Scheme 2). First, S_N2'-substitution of mesylate in **5** leads to allene **8**, which, upon intramolecular nucleophilic attack of pyridyl nitrogen at the Cu-activated double bond of allene produces cyclic intermediate **9**. The latter can transform into product **6** either via the deprotonation-protonation sequence (path A), or through the 1,2-hydride shift in the carbenoid intermediate **9** (path B). To verify which mechanism operates, we performed a deuterium-labelling experiment. Indeed, if the reaction proceeds via the Path A, then cyclization of **5b-d** should produce **6ba**, a product with significant or total deuterium loss.¹¹

In contrast, the end-game via Path B would give **6ba-d** with complete preservation of the deuterium label.¹² It was found that, under the standard cyclization conditions, the reaction of isotopically pure **5b-d** produced indolizine **6ba** with total deuterium loss, thus strongly supporting Path A.

In summary, we have developed a novel organocopper-mediated two-component coupling/cycloisomerization cascade transformation. This mild and efficient method allows for easy synthesis of C-1 alkyl- and aryl-substituted N-fused heterocycles, such as indolizines, pyrroloquinolines, and pyrroloisoquinolines, from easily available starting materials. It deserves mentioning that these important heterocyclic scaffolds¹³ with alkyl or aryl substituent at C-1 are not available via the existing cycloisomerization methods.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

Financial support from the National Institutes of Health (GM-64444) is gratefully acknowledged.

References

1. For recent reviews, see: (a) Fürstner A, Davies PW. *Angew Chem Int Ed* 2007;46:3410. (b) Alonso F, Beletskaya IP, Yus M. *Chem Rev* 2004;104:3079. [PubMed: 15186189] (c) Nakamura I, Yamamoto Y. *Chem Rev* 2004;104:2127. [PubMed: 15137788]
2. (a) Kel'in AV, Sromek AW, Gevorgyan V. *J Am Chem Soc* 2001;123:2074. [PubMed: 11456838] (b) Kim JT, Butt J, Gevorgyan V. *J Org Chem* 2004;69:5638. [PubMed: 15307734] (c) Kim JT, Gevorgyan V. *J Org Chem* 2005;70:2054. [PubMed: 15760187] (d) Schwier T, Sromek AW, Yap DML, Chernyak D, Gevorgyan V. *J Am Chem Soc* 2007;129:9868. [PubMed: 17658805]
3. (a) Seregin IV, Gevorgyan V. *J Am Chem Soc* 2006;128:12050. [PubMed: 16967938] (b) Seregin IV, Schammel AW, Gevorgyan V. *Org Lett* 2007;9:3433. [PubMed: 17637023] (c) Smith CR, Bunnelle EM, Rhodes AJ, Sarpong R. *Org Lett* 2007;9:1169. [PubMed: 17309277] (d) Hardin AR, Sarpong R. *Org Lett* 2007;9:4547. [PubMed: 17845052] (e) Yan B, Zhou Y, Zhang H, Chen J, Liu Y. *J Org Chem* 2007;72:7783. [PubMed: 17718578] (f) Kim J, Choi HK, Won G, Lee H. *Tetrahedron Lett* 2007;48:6863.
4. For a general review, see: (a) Krause N Hashmi SK Modern Allene Chemistry Wiley-VCH Weinheim 2004 1143. See also: (b) Krause N, Hoffmann-Roeder A. *Tetrahedron* 2004;60:11671. (c) Krause N, Hoffmann-Roeder A. *Angew Chem Int Ed* 2002;41:2933.
5. For recent examples on synthesis of allenes using copper reagents via S_N2' substitution approach, see: (a) Saito A, Kanno A, Hanzawa Y. *Angew Chem Int Ed* 2007;46:3931. (b) Ghosh P, Lotesta SD, Williams LJ. *J Am Chem Soc* 2007;129:2438. [PubMed: 17288427] (c) Dieter RK, Chen N, Gore VK. *J Org Chem* 2006;71:8755. [PubMed: 17081003] (d) Pacheco MC, Gouverneur V. *Org Lett* 2005;7:1267. [PubMed: 15787483] (e) Dieter RK, Chen N, Yu H, Nice LE, Gore VK. *J Org Chem* 2005;70:2109. [PubMed: 15760194] (f) Regás D, Afonso MM, Rodríguez ML, Palenzuela JA. *J Org Chem* 2003;68:7845. [PubMed: 14510565] (g) Dieter RK, Yu H. *Org Lett* 2001;3:3855. [PubMed: 11720553]
6. For copper-mediated cascade transformations, see for example: (a) Sherman ES, Chemler SR, Tan TB, Gerlits O. *Org Lett* 2004;6:1573. [PubMed: 15128239] (b) Sherman ES, Fuller PH, Kasi D, Chemler SR. *J Org Chem* 2007;72:3896. [PubMed: 17428100]
7. For general reviews on copper reagents, see: (a) Lipshutz BH *Organometallics in Synthesis: A Manual* Schlosser MWiley-VCH Weinheim 1994 283376 (b) Krause N *Modern Organocopper Chemistry* Wiley-VCH Weinheim 2002 (c) Caprio V. *Lett Org Chem* 2006;3:339. (d) Nakamura E, Seiji M. *Angew Chem Int Ed* 2000;39:3750. (e) Woodward S. *Chem Soc Rev* 2000;29:393. (f) Krause N, Gerold A. *Angew Chem Int Ed* 1997;36:186. (g) Taylor RJ *CasyG Organocopper Reagents-A Practical Approach* Taylor RJ KOxford University Press New York 1994
8. It was also recently reported by Krause that the lower-order cyanocuprates provide high selectivity in S_N2' -substitution reaction employing propargyl acetates. See: (a) Jansen A, Krause N. *Inorg Chem Acta* 2006;359:1761. (b) Jansen A, Krause N. *Synthesis* 2002:1987.
9. For a review, see: (a) Seregin IV, Gevorgyan V. *Chem Soc Rev* 2007;36:1173. [PubMed: 17576484]. For a direct arylation of indolizines, see: (b) Park C-H, Ryabova V, Seregin IV, Sromek AW, Gevorgyan V. *Org Lett* 2004;6:1159. [PubMed: 15040747]. For a direct alkynylation of indolizines, see: (c) Seregin IV, Gevorgyan V. *J Am Chem Soc* 2007;129:7742. [PubMed: 17539646]
10. See Supporting Information for complete experimental data.
11. For substantial deuterium loss in the Cu-mediated cycloisomerization, see ref. 2a.
12. For complete preservation of the deuterium label in the cycloisomerization proceeding via a 1,2-deuterium shift in carbenoid intermediate, see (a) Sromek AW, Rubina M, Gevorgyan V. *J Am Chem*

- Soc 2005;127:10500. [PubMed: 16045332] (b) Dudnik AS, Sromek AW, Rubina M, Kim JT, Kel'in AV, Gevorgyan V. J Am Chem Soc 2008;130:1440. [PubMed: 18173272]
13. For biological activity of N-fused pyrroloheterocycles, see for example: (a) Fujita T, Matsumoto Y, Kimura T, Yokota S, Sawada M, Majima M, Ohtani Y, Kumagai Y. Br J Clin Pharmacol 2002;54:283. [PubMed: 12236849] (b) Facompré M, Tardy C, Bal-Mahieu C, Colson P, Perez C, Manzanares I, Cuevas C, Bailly C. Cancer Res 2003;63:7392. [PubMed: 14612538] (c) Reddy MV, Rao MR, Rhodes D, Hansen MS, Rubins K, Bushman FD, Venkateswarlu Y, Faulkner DJ. J Med Chem 1999;42:1901. [PubMed: 10354398] (d) Østby OB, Dalhus B, Gundersen L-L, Rise F, Bast A, Haenen GRMM. Eur J Org Chem 2000:3763. (e) Hagishita S, Yamada M, Shirahase K, Okada T, Murakami Y, Ito Y, Matsuura T, Wada M, Kato T, Ueno M, Chikazawa Y, Yamada K, Ono T, Teshirogi I, Ohtani M. J Med Chem 1996;39:3636. [PubMed: 8809154]

Scheme 2.
Proposed mechanism for cascade cyclization

Table 1

Optimization of reaction conditions

run	$[CuR]^a$	LG	solvent	yield % ^b
1	s-Bu(Me)Cu(CN)Li ₂	OMs	Et ₂ O	11
2	n-Bu ₂ Cu(CN)Li ₂	OMs	Et ₂ O	traces
3	Me ₂ Cu(CN)Li ₂	OMs	THF	0
4	MeCu(CN)Li	OMs	Et ₂ O	8
5	MeCu(CN)Li	OMs	THF	90
6	MeCu(CN)Li	OAc	THF	21
7	MeCu(CN)Li	OTFA	THF	16
8	MeCu(CN)Li	OC(O)OEt	THF	0
9	MeCu(CN)Li	OP(O)OEt ₂	THF	27
10	MeCu(CN)Li	OTf	THF	5
11	MeCu(CN)Li	OBs	THF	traces
12	MeCu(CN)Li	ONs	THF	traces
13	Ph ₂ Cu(CN)Li ₂	OMs	Et ₂ O	traces
14	Ph ₂ Cu(CN)Li ₂ /BF ₃ ·Et ₂ O	OMs	Et ₂ O-THF	9
15	PhCu(CN)Li	OMs	Et ₂ O	0
16	PhCu^c	OMs	THF	71

^a Reactions were carried with 1.0 mmol of **5** and 1.2 mmol of copper reagent at -78 °C for 1 h, then at rt for 1 h.

^b Isolated yield.

^c Prepared by mixing PhMgBr with CuBr·SMe₂ complex.

Table 2Organocopper-mediated synthesis of N-fused heterocycles^a

entry	starting material ^b 5	[CuR ²] 7	product 6	yield ^c %
1	 5a	MeCu(CN)Li (7a)	 6aa	90
2	5a	<i>n</i> -BuCu(CN)Li (7b)	 6ab	89
3	5a	<i>n</i> -HexCu(CN)Li (7c)	 6ac	81
4	5a	<i>s</i> -BuCu(CN)Li (7d)	 6ad	82
5	5a	<i>t</i> -BuCu(CN)Li (7e)	 6ae	79
6	 5b	7a	 6ba	70 ^d

entry	starting material ^b 5	[CuR ²] 7	product 6	yield ^c %
7	<p>5c</p>	7a	<p>6ca</p>	80 ^e
8	<p>5d</p>	7a	<p>6da</p>	61
9	<p>5e</p>	7a	<p>6ea</p>	75
10	<p>5f</p>	7a	<p>6fa</p>	62
11	<p>5g</p>	7a	<p>6ga</p>	74
12	<p>5h</p>	7a	<p>6ha</p>	63
13	<p>5i</p>	7a	<p>6ia</p>	77

entry	starting material ^b 5	[CuR ²] 7	product 6	yield ^c %
14	5a	PhCu (7f)	6af	71
15	5j	7a	6ja	65
16	5j	7f	6jf	51
17	5k	7a	6ka	66

^a Reaction carried out in 1.0 - 1.5 mmol scale.

^b Mesylate prepared in situ.

^c Isolated yields.

^d Additionally, 14% of allene was isolated.

^e Additionally, 4% of allene was isolated.

Table 3

Attempts on stepwise synthesis of N-fused heterocycles

run	R ¹	R ²	yield ^a % 8	yield ^a % 6	overall yield ^b % 6
1	H	Me	27 ^c	0	trace (61)
2	Me	<i>s</i> -Bu	89	42	37 (82)
3	C ₆ H ₄ - <i>p</i> -CO ₂ Me	Me	34	27	9 (62)

^a Isolated yield.

^b Overall isolated yield for a stepwise process. Yield for cascade transformation from Table 2 is given in parenthesis.

^c Trace amounts of **6** were also isolated.