Published in final edited form as:

Cancer Res. 2009 January 15; 69(2): 475–482. doi:10.1158/0008-5472.CAN-08-2649.

Transforming growth factor β induces clustering of HER2 and integrins by activating Src-FAK and receptor association to the cytoskeleton

Shizhen Emily Wang^{1,5}, Bin Xiang², Roy Zent², Vito Quaranta³, Ambra Pozzi², and Carlos L. Arteaga^{2,3,4,5}

¹ Division of Tumor Cell Biology, Beckman Research Institute of City of Hope, Duarte, California, 91010

² Department of Medicine; Vanderbilt University School of Medicine, Nashville, Tennessee 37232

³ Department of Cancer Biology; Vanderbilt University School of Medicine, Nashville, Tennessee 37232

⁴ Breast Cancer Research Program, Vanderbilt-Ingram Comprehensive Cancer Center; Vanderbilt University School of Medicine, Nashville, Tennessee 37232

Abstract

It has been proposed that cross-talk between integrin and growth factor receptor signaling such as ErbB2 (HER2) is required for activation of downstream effectors and ErbB2-mediated mammary tumorigenesis. Here we show that transforming growth factor β (TGF- β) induced focal adhesion kinase (FAK)-dependent clustering of HER2 and integrin $\alpha 6$, $\beta 1$ and $\beta 4$ in HER2-overexpressing mammary epithelial cells without altering the total and surface levels of HER2 receptors. This effect was mediated by ligand-induced EGFR activation and the subsequent phosphorylation of Src and FAK. We have previously reported that TGF- β upregulates EGFR ligand shedding through a mechanism involving the phosphorylation of TNF- α converting enzyme (TACE/ADAM17). Knock down of TACE, FAK or integrin $\alpha 6$ by siRNA or inhibition of EGFR or Src by specific inhibitors abrogated TGF- β -induced receptor clustering and signaling to PI3K-Akt. Finally, inhibition of Src-FAK reversed TGF- β -induced resistance to the therapeutic HER2 inhibitor trastuzumab in HER2-overexpressing breast cancer cells. Taken together, these data suggest that, by activating Src-FAK, TGF- β integrates ErbB receptor and integrin signaling to induce cell migration and survival during breast cancer progression.

Keywords

TGF- β ; HER2 (ErbB2); integrin; FAK; Src

Introduction

The integrin family of transmembrane receptors of the extracellular matrix (ECM) mediates a wide variety of cellular behaviors including cell adhesion, migration, proliferation and survival (1,2). In addition to transducing signals in the ECM to the cell membrane and initiating intracellular signaling through integrin-associated proteins such as focal adhesion kinase

5Corresponding authors: SEW: E-MAIL E-mail: ewang@coh.org; FAX 626-301-8972, Division of Tumor Cell Biology, Beckman Research Institute of City of Hope, 1500 East Duarte Road, KCRB Room 2007, Duarte, CA 91010, USA TEL 626-256-4673 x63118, CLA: E-MAIL E-mail: carlos.arteaga@vanderbilt.edu; FAX 615-936-1790, Division of Oncology, Vanderbilt University School of Medicine, 2200 Pierce Avenue, 777 PRB, Nashville, TN 37232-6307, USA, TEL 615-936-3524.

(FAK), integrins are also involved in growth factor signaling through co-clustering with the growth factor receptor tyrosine kinases (RTKs) such as the epidermal growth factor receptor (EGFR), HER2 (ErbB2) and platelet-derived growth factor receptor (PDGFR) (3,4). Phosphorylation of integrins by RTKs and non-receptor tyrosine kinases such as FAK and Src may, in turn, regulate “inside-out” integrin signaling by altering their affinity and avidity to ECM ligands. The membrane-proximal clustering of integrin and RTK requires FAK as the scaffold that interacts with both RTK and integrin receptors through different domains (3).

HER2 gene amplification or overexpression of HER2 is observed in about 30% of invasive breast cancers, where it correlates with poor patient prognosis. Several integrin-associated signaling molecules, including both integrin $\beta 1$ and $\beta 4$, integrin-linked kinase (ILK) and FAK, have been implicated in the initiation and progression of mammary tumors driven by oncogenes such as Neu (the rat homologue of HER2) and the polyomavirus middle T (PyVMT) (4–7). In the transgenic mouse model of MMTV (mouse mammary tumor virus)/Neu, targeted deletion of $\beta 4$ integrin was shown to suppress mammary tumor onset and invasive growth (4).

Transforming growth factor β (TGF- β), a multitasking cytokine involved in development, differentiation, tissue regeneration, and immune response, has also been shown to synergize with Neu/ErbB2 (8–11). Overexpression of active TGF- $\beta 1$ or active mutants of the type I TGF- β receptor (T β RI, Alk5) in the mammary gland of bi-transgenic mice also expressing MMTV/Neu accelerates metastases from Neu-induced mammary cancers (8–10). In the MCF10A human mammary epithelial cells that were engineered to overexpress HER2 (MCF10A/HER2), TGF- β induces cell migration and invasion (12,13). Inhibition of HER2 with the HER2-neutralizing antibody trastuzumab blocked the promigratory effect of TGF- β on HER2-overexpressing mammary epithelial cells (12), suggesting that the proto-oncogene is required for the transforming effect of TGF- β in HER2-overexpressing transformed cells.

We have previously shown that TGF- β induces HER2 translocation to the lamellipodia through a PI3K-dependent mechanism that involves activation of Rac1 and Rho1 and reorganization of actin cytoskeleton (13). Moreover, blockade of integrin signaling by an inhibitory antibody against $\beta 1$ integrin abrogates the function of TGF- β to induce motility in MCF10A/HER2 cells (12). Because TGF- β is known to influence various integrin functions such as adhesion and migration, we focused in this study on the role of integrin signaling in the crosstalk between TGF- β and HER2 in breast cancer progression.

Materials and methods

Cells, reagents and viruses

MCF10A/HER2 cells were generated and maintained as described previously (12,14). Human breast cancer cell lines MDA-MB-453 and BT-474 were purchased from the American Type Tissue Culture Collection and maintained in IMEM (Cellgro) containing 10% fetal bovine serum (FBS; Hyclone) in a humidified 5% CO₂ incubator at 37°C. Recombinant human TGF- $\beta 1$ and TGF- α was purchased from R&D Systems. Brefeldin A, PP2 and cytochalasin D were purchased from Sigma. Trastuzumab and cetuximab were purchased at the Vanderbilt University Medical Center Pharmacy. Lapatinib ditosylate (GW-572016) was from LC Laboratories. Adenoviruses encoding dominant negative (DN) p85 (A Δ p85) was described previously (15). The plasmid encoding the HA-tagged full-length mouse TACE [TACE(HA)] was described previously (16).

Matrix preparation and mass spectrometry (MS)

Preparation of deposited extracellular matrix and matrix-associated proteins was performed as described (17). Briefly, confluent MCF10A/HER2 cells were lysed on plate by incubating in

20 mM sterile NH_4OH for 5 min at room temperature (RT) followed by gently scraping and extensive washing. The remaining matrix preparation was directly scraped into reducing Laemmli sample buffer, boiled, and applied to SDS-PAGE. The gel was stained with Coomassie blue. Excised protein bands were subjected to *in-gel* digestion as described elsewhere (13).

Cell surface biotinylation, immunoprecipitation (IP) and immunoblot

Cells grown in 100-mm dishes were washed in cold PBS (pH 8.0) for 3 times before incubated with freshly prepared Sulfo-NHS-Biotin reagent (2 mM; Pierce) 30 min at 4°C. The reaction was quenched with 100 mM glycine in PBS and the cells were lysed in NP-40 lysis buffer (20 mM Tris, 150 mM NaCl, 1% Nonidet P-40, 0.1 mM EDTA, plus protease and phosphatase inhibitors). After sonication for 10 s and centrifugation (14,000 rpm), protein concentration in the supernatants was measured using the BCA protein assay reagent (Pierce). Equal amounts of protein extracts (500 μg) were subjected to precipitation using Streptavidin Magnetic Spheres (Promega) followed by SDS-PAGE and HER2 immunoblot. For immunoprecipitation, cells were washed twice with cold PBS and lysed in NP-40 lysis buffer. Immunoprecipitation and immunoblotting were performed as described (14) utilizing horseradish peroxidase-conjugated secondary antibodies (Promega). Primary antibodies include: integrin α_6 , α_3 , β_1 , β_4 , TACE (Chemicon), EGFR, HER2/ErbB2 (NeoMarkers), actin (Sigma), FAK, Src (Santa Cruz Biotechnology), P-EGFR(Y1068), P-HER2(Y1248), P-Src(Y416), P-FAK(Y397), P-Akt(S473), Akt (Cell Signaling, Beverly, Mass.), and p85 (Upstate Biotechnology). Bands were quantified using ImageJ.

Fluorescence-activated cell sorter analysis

Cells were harvested by trypsinization, washed with PBS, and exposed to monoclonal antibodies of the appropriate integrin (1:50 in DMEM) followed by a FITC-coupled rabbit anti-mouse secondary IgG (1:50). Flow cytometry was performed with a FACScan instrument (Becton Dickinson).

Cell viability, motility and invasion assays

For viability assays, cells growing on 6-well plates were serum-starved for 48 h and collected for Trypan blue staining as described elsewhere (18). Trypan blue-positive cells were counted manually and their percentage calculated over the total cell input. Transwell motility assays were performed as described previously (19). Invasion assay was performed using laminin- or collagen-coated transwells (BD Biosciences) according to the manufacturer's protocol. Cells were seeded at 2.5×10^4 cells/well on coated transwells and allowed to invade toward serum-free medium $\pm$ TGF- β . At 16 h, cells invaded through the transwell filters were stained and counted as described (19).

Three-dimensional cell culture

Cells were seeded on Growth Factor Reduced Matrigel (BD Biosciences) in 8-well chamber slides following the protocol described by Debnath *et al.* (20). Inhibitors were added into the medium 12 h after cell seeding and were replenished every 3 days. Phase contrast images were taken every 3 days. For cell number counting, cultures growing on Matrigel were trypsinized and cell numbers were measured in a Coulter counter.

RNA interference studies

Silencer siRNAs against human TACE were obtained from Qiagen in a FlexiTube format. Four species of siRNAs against different TACE target sequences (TCCCATGAAGAACACGTGTAA; CTGCAGTAAACAATCAATCTA; CAGGATGTAATTGAACGATTT; AAGAAACAGAGTGCTAATTTA) were mixed in

equal amount to generate a TACE siRNA pool (siTACE). SiRNAs against FAK target sequences (CACCTGGGTACTGGTATGGAA; AATCACACACCAAATTCGAGT; CCGGTCGAATGATAAGGTGTA; AACAAATTTATGTTCACATTAA) and integrin $\alpha 6$ target sequences (CAGGGTAATAAACTTAGGTAA; CCGGCCTGTGATTAATATTCA; AAGGATGGGTGGCAAGATATA; AACCTGTGGCTACAGGATAA) (Qiagen) were used. SiRNAs were transfected into cell lines using HiPerFect transfection reagent (Qiagen) according to the manufacturer's procedures. In a 6-well plate format, a total of 150 ng of siRNAs and 12 μ l of HiPerFect reagent were used for each transfection. Mismatched siRNAs were used as negative control (siCTRL).

Results

TGF- β induces clustering of HER2 and integrin $\alpha 6$, $\beta 1$ and $\beta 4$ at the cell membrane

We first determined the extracellular matrix (ECM) deposited by MCF10A/HER2 cells. Cells growing on tissue culture dishes were lysed as described in Materials and Methods; proteins attached to the plate were harvested and subjected to mass spectrometry (MS). Laminin-5, which consists of the laminin $\alpha 3$, $\beta 3$ and $\gamma 2$ chains, as well as integrin $\alpha 6$, $\alpha 3$, $\beta 1$ and $\beta 4$ were identified in the matrix deposited by 10A/HER2 cells by MS (Figure 1A). We also analyzed the ECM composition in cells treated with TGF- β for 24 hrs using the same approach and did not find different protein species between treated and untreated cells. As determined by flow cytometry using specific antibodies, treatment with TGF- β for 24 hrs did not affect the abundance of integrin $\alpha 6$, $\alpha 3$ and $\beta 1$ on the cell surface while slightly decreasing the surface levels of integrin $\beta 4$ (Figure 1B). By immunoprecipitation and immunoblot analysis with HER2 and integrin antibodies, TGF- β induced the association between HER2 and integrin $\alpha 6$, $\beta 1$ and $\beta 4$ at 1 hr after treatment without affecting their total levels. Although expression of integrin $\alpha 3$ was detected in these cells by immunoblot, HER2 did not associate with this integrin in the absence or presence of TGF- β (Figure 1C). In 10A/HER2 and in HER2-overexpressing MDA-MB-453 and BT474 breast cancer cells, HER2 was detected in ECM-associated complexes and this association was markedly increased by TGF- β (Figure 1D). The level of cell surface-localized HER2 as determined by surface biotinylation followed by pull-down with streptavidin beads and HER2 immunoblot, and was not affected by TGF- β in all three cell lines. Brefeldin A, an inhibitor of intracellular protein transport, abrogated the cell surface localization of HER2 and its association with the ECM (Figure 1D).

TGF- β activates Src-FAK, leading to receptor clustering and activation of PI3K-Akt

It has been shown that the membrane-proximal clustering of integrin and RTK requires the scaffolding function of FAK (3). Thus, to determine the role of the FAK in TGF- β -modulated clustering of HER2 and integrin, we performed immunoprecipitation with HER2 and integrin $\beta 1$ antibodies followed by FAK immunoblot. FAK associated with both HER2 and integrin $\beta 1$ and this association was increased by the addition of TGF- β in the same time course as it induced HER2:integrin clustering (Figure 2A). Concurrently, tyrosine phosphorylation of FAK (site Y397) and Src (Y416) was also induced by TGF- β . Phosphorylation of the Y397 Src-binding site of FAK is required for p85 binding and PI3K activation (21). In line with this report, activation of Akt was also detected along with FAK phosphorylation during the time course following TGF- β treatment (Figure 2A). Inhibition of Src kinase activity with PP2 abolished both basal and TGF- β -induced association of HER2 with FAK and integrin $\beta 1$ and the TGF- β -induced activation of FAK and Akt (Figure 2B). Finally, knock-down of FAK expression using specific siRNA abolished the ability of TGF- β to induce receptor clustering and FAK-mediated activation of Akt and severely impaired both basal and TGF- β -induced phosphorylation of HER2 and EGFR as detected with site-specific phospho-antibodies (Figure 2C). Control IP with anti-integrin $\alpha 3$ was also performed in all experiments and showed negative results (data not shown).

TGF- β induces Src-FAK through EGFR activation

In a recent study (22), we have reported that TGF- β induces the shedding of ErbB ligands TGF- α , amphiregulin and heregulin through a TACE-dependent mechanism. To explore whether this is responsible for the activation of Src-FAK pathway by TGF- β , we used the EGFR antibody cetuximab to block ligand binding to EGFR or siRNA to knock-down the expression of TACE. Both cetuximab and RNAi of TACE abrogated TGF- β -induced FAK activation and HER2 clustering with integrin (Figure 3A&B). Transfection of a full-length mouse TACE construct [TACE(HA)] reconstituted TGF- β -induced effects in cells transfected with human TACE siRNA (Figure 3B). Further, EGFR activation by its ligand TGF- α was sufficient to induce activation of the Src-FAK pathway as well as receptor clustering. This was impaired by addition of the Src kinase inhibitor PP2 (Figure 3C). In HER2-overexpressing BT-474 breast cancer cells, TGF- β also induced Src activation and HER2 clustering with integrin; both effects were abolished by TACE siRNA oligonucleotides or the addition of PP2 (Figure S1A&B).

Integrin $\alpha 6$ is essential for the clustering of HER2, integrin $\beta 1$ and FAK

Clustering between HER2 and integrin has only been observed with integrin $\alpha 6\beta 1$ and $\alpha 6\beta 4$, suggesting a unique role of $\alpha 6$ integrin on HER2 signaling. We further evaluated the function of this integrin in HER2 signaling and its crosstalk with the TGF- β network using RNA interference. Knock-down of $\alpha 6$ expression abolished the formation of the HER2:FAK:integrin complex and impaired FAK-mediated PI3K signaling to Akt (Figure 4A). In addition, the ability of TGF- β to enhance migration and survival in MCF10A/HER2 cells in serum-free medium were lost upon depletion of $\alpha 6$ integrin or FAK (Figure 4B&C). TGF- β induced invasion of 10A/HER2 cells on laminin but not collagen; this effect was also impaired when $\alpha 6$ integrin or FAK was knocked down with siRNA oligonucleotides (Figure 4D).

TGF- β induces association of HER2 and integrins with the cytoskeleton

We have previously shown that TGF- β spatially regulates HER2 signaling by inducing its association with the actin cytoskeleton and relocalization to lamellipodia in a PI3K-dependent manner (13). Integrin receptors are also anchored to the cytoskeleton through adaptor proteins such as FAK (23). In this study, TGF- β induced association of actin with both HER2 and integrin $\alpha 6$; this association was impaired by the HER2 tyrosine kinase inhibitor lapatinib ditosylate and the Src kinase inhibitor PP2 (Figure 5A). Expression of dominant-negative (DN) p85 also abolished the association of integrin $\beta 1$ with actin and HER2:FAK:integrin complex formation (Figure 5B). Treatment with cytochalasin D, a potent inhibitor of actin polymerization, impaired FAK phosphorylation and receptor clustering upon TGF- β treatment and diminished HER2 signaling in both basal and TGF- β -treated conditions (Figure 5C). Basal Akt phosphorylation was increased as a result of cytochalasin D treatment; the mechanism of this is unknown but we speculate it may reflect a compensatory cellular response to the stress from drug treatment.

Inhibition of Src-FAK reverses TGF- β -induced resistance to trastuzumab in HER2-overexpressing breast cancer cells

In a recent work we have shown that TGF- β upregulates PI3K/Akt and, in turn, desensitizes HER2-overexpressing cells to trastuzumab (22). Here we further found that FAK was phosphorylated at Y397 and associated with PI3K p85 upon TGF- β stimulation; knock-down of FAK using siRNA oligonucleotides abolished TGF- β -mediated Akt activation (Figure 2C). Therefore, we speculated that inhibition of Src-FAK might reverse TGF- β -induced resistance to trastuzumab. To explore this, we cultured HER2-overexpressing BT-474 breast cancer cells on Matrigel in the absence or presence of TGF- β and PP2 and examined their responses to trastuzumab. Addition of PP2 blocked the protective effect of TGF- β and restored trastuzumab

action in BT-474 cells (Figure 6), consistent with the ability of the Src inhibitor to inhibit P-Akt (shown in Figures 2&3).

Discussion

TGF- β is known to play dual roles in tumor progression. Whereas it functions as a tumor suppressor in normal cells or during early stages of tumorigenesis, it is often found as a pro-metastatic factor in established or invasive tumors through a combination of autocrine and paracrine mechanisms (24,25). Coexpression of activated oncogenes is a cue to the pro-tumorigenic functions of TGF- β . In cancer models that depend on Neu (ErbB2), PyVMT, or mutant Ras, TGF- β synergizes with these oncogenes to accelerate cancer progression and metastasis (8,10,26–28). Recent work (22) and data shown herein indicate that in HER2 (ErbB2)-overexpressing cells, TGF- β enhances ErbB-initiated signal transduction by increasing ErbB ligand shedding, HER2-containing heterodimers, and their crosstalk with integrins (Figure 5D). The latter is mediated by the activation of Src-FAK; inhibition of this kinase complex not only abolishes the ability of TGF- β to facilitate cancer progression but also reverses TGF- β -mediated resistance to trastuzumab. Interestingly, RNA interference of integrin $\alpha 6$ or FAK completely abrogated the pro-survival and pro-migratory functions of TGF- β in HER2-overexpressing cells (Figure 4C), suggesting that integrin-regulated interactions of HER2 with the actin cytoskeleton and not the total levels of HER2 protein can specify the qualitative response to TGF- β .

In a previous study we have examined effects of TGF- β on the cellular distribution and signaling of HER2 using MCF10A/HER2 and breast cancer lines that naturally overexpress the protooncogene. In these cells, treatment with TGF- β activates PI3K and activates the Vav2 guanine nucleotide exchange factor, which then activates Rac1 and its effector Pak1, resulting in colocalization of HER2 with these molecules as well as actin cytoskeleton components at cell protrusions (13). Here we show that FAK is required for PI3K/Akt activation by TGF- β . In addition to activating PI3K/Akt, FAK is also known to activate CAS-Crk-Rac signaling and regulate the assembly and turnover of focal adhesions at the leading lamellipodia of migrating cells. This occurs through the regulation of integrin-mediated cell adhesion and cytoskeleton rearrangement (Figure 5D).

The dual kinase Src/FAK complex has an established role in human cancers (29). In cancers, both growth factor receptor and FAK signaling are upregulated. Autophosphorylation of FAK on Y397 creates a SH2 domain binding site that can recruit and activate Src. FAK phosphorylated at this site has been found in various tumors including ovarian, cervical carcinomas and acute myeloid leukemia [reviewed in (29)]. Phosphorylation of this site is also responsible for the direct association of FAK with the regulatory subunit of PI3K, p85, as well as the activation of PI3K by ligand-bound PDGFR (21). Thus, FAK integrates both growth factor receptor tyrosine kinase and integrin networks and serves as an effector for these two signaling systems. Indeed, cells lacking FAK are refractory to EGF- or PDGF-induced tyrosine phosphorylation of EGFR and PDGFR, respectively, and cell motility (3). Consistently, FAK RNAi suppressed both basal and TGF- β -induced tyrosine phosphorylation of HER2 and EGFR (Figure 2C), suggesting that FAK-mediated cross-talk between receptor tyrosine kinases and integrins is required for receptor activation.

Integrin signaling has been shown to play a crucial role in ErbB2-mediated mammary tumorigenesis. Integrin $\beta 4$ amplifies ErbB2 signaling to promote mammary tumor onset and invasion in transgenic mice (4). Moreover, deletion of the $\beta 4$ signaling domain enhances the efficacy of gefitinib, a small molecule inhibitor of the EGFR tyrosine kinase (4). The role of the Src-FAK complex on regulating the transforming role of integrins suggests it can be targeted therapeutically in human cancers. Inhibitors of the Src kinase are currently in early

clinical development for the treatment of solid tumors including breast cancer. In addition, amplified TGF- β signaling has been shown to upregulate PI3K and desensitize the antitumor effect of HER2 antagonist trastuzumab (22). Our data herein further suggest that 1) Src-FAK is essential for the activation of PI3K that results from the interaction between HER2 and TGF- β , and 2) inhibition of Src-FAK reverses TGF- β -induced resistance to HER2 antagonist trastuzumab.

Acknowledgements

This work was supported by American Cancer Society-Institutional Research Grant #IRG-58-009-49 (SEW), NCI K99 CA125892 (SEW), NCI R01 CA62212 (CLA), R01 CA80195 (CLA), ACS Clinical Research Professorship Grant CRP-07-234 (CLA), R01 DK69921 (RZ), R01 DK075594 (RZ), R01 DK074359 (AP), Merit Award from the Department of Veterans Affairs (RZ and AP), Breast Cancer Specialized Program of Research Excellence (SPORE) Grant P50 CA98131, and Vanderbilt-Ingram Comprehensive Cancer Center Support Grant P30 CA68485.

References

- Hood JD, Chersesh DA. Role of integrins in cell invasion and migration. *Nat Rev Cancer* 2002;2:91–100. [PubMed: 12635172]
- Hynes RO. Integrins: bidirectional, allosteric signaling machines. *Cell* 2002;110:673–87. [PubMed: 12297042]
- Sieg DJ, Hauck CR, Ilic D, et al. FAK integrates growth-factor and integrin signals to promote cell migration. *Nat Cell Biol* 2000;2:249–56. [PubMed: 10806474]
- Guo W, Pylyayeva Y, Pepe A, et al. Beta 4 integrin amplifies ErbB2 signaling to promote mammary tumorigenesis. *Cell* 2006;126:489–502. [PubMed: 16901783]
- White DE, Cardiff RD, Dedhar S, Muller WJ. Mammary epithelial-specific expression of the integrin-linked kinase (ILK) results in the induction of mammary gland hyperplasias and tumors in transgenic mice. *Oncogene* 2001;20:7064–72. [PubMed: 11704830]
- White DE, Kurpios NA, Zuo D, et al. Targeted disruption of beta1-integrin in a transgenic mouse model of human breast cancer reveals an essential role in mammary tumor induction. *Cancer Cell* 2004;6:159–70. [PubMed: 15324699]
- Lahlou H, Sanguin-Gendreau V, Zuo D, et al. Mammary epithelial-specific disruption of the focal adhesion kinase blocks mammary tumor progression. *Proc Natl Acad Sci U S A* 2007;104:20302–7. [PubMed: 18056629]
- Muraoka RS, Koh Y, Roebuck LR, et al. Increased malignancy of Neu-induced mammary tumors overexpressing active transforming growth factor beta1. *Molecular and cellular biology* 2003;23:8691–703. [PubMed: 14612410]
- Muraoka-Cook RS, Shin I, Yi JY, et al. Activated type I TGFbeta receptor kinase enhances the survival of mammary epithelial cells and accelerates tumor progression. *Oncogene* 2006;25:3408–23. [PubMed: 16186809]
- Siegel PM, Shu W, Cardiff RD, Muller WJ, Massague J. Transforming growth factor beta signaling impairs Neu-induced mammary tumorigenesis while promoting pulmonary metastasis. *Proc Natl Acad Sci U S A* 2003;100:8430–5. [PubMed: 12808151]
- Seton-Rogers SE, Lu Y, Hines LM, et al. Cooperation of the ErbB2 receptor and transforming growth factor beta in induction of migration and invasion in mammary epithelial cells. *Proc Natl Acad Sci U S A* 2004;101:1257–62. [PubMed: 14739340]
- Ueda Y, Wang S, Dumont N, Yi JY, Koh Y, Arteaga CL. Overexpression of HER2 (erbB2) in human breast epithelial cells unmasks transforming growth factor beta-induced cell motility. *J Biol Chem* 2004;279:24505–13. [PubMed: 15044465]
- Wang SE, Shin I, Wu FY, Friedman DB, Arteaga CL. HER2/Neu (ErbB2) Signaling to Rac1-Pak1 Is Temporally and Spatially Modulated by Transforming Growth Factor {beta}. *Cancer Res* 2006;66:9591–600. [PubMed: 17018616]
- Wang SE, Wu FY, Shin I, Qu S, Arteaga CL. Transforming growth factor {beta} (TGF-{beta})-Smad target gene protein tyrosine phosphatase receptor type kappa is required for TGF-{beta} function. *Mol Cell Biol* 2005;25:4703–15. [PubMed: 15899872]

15. Kotani K, Ogawa W, Hino Y, et al. Dominant negative forms of Akt (protein kinase B) and atypical protein kinase Clambda do not prevent insulin inhibition of phosphoenolpyruvate carboxykinase gene transcription. *J Biol Chem* 1999;274:21305–12. [PubMed: 10409689]
16. Diaz-Rodriguez E, Montero JC, Esparis-Ogando A, Yuste L, Pandiella A. Extracellular signal-regulated kinase phosphorylates tumor necrosis factor alpha-converting enzyme at threonine 735: a potential role in regulated shedding. *Mol Biol Cell* 2002;13:2031–44. [PubMed: 12058067]
17. Falk-Marzillier J, Domanico SZ, Pelletier A, Mullen L, Quaranta V. Characterization of a tight molecular complex between integrin alpha 6 beta 4 and laminin-5 extracellular matrix. *Biochem Biophys Res Commun* 1998;251:49–55. [PubMed: 9790905]
18. Wu FY, Wang SE, Sanders ME, et al. Reduction of cytosolic p27(Kip1) inhibits cancer cell motility, survival, and tumorigenicity. *Cancer Res* 2006;66:2162–72. [PubMed: 16489017]
19. Wang SE, Narasanna A, Perez-Torres M, et al. HER2 kinase domain mutation results in constitutive phosphorylation and activation of HER2 and EGFR and resistance to EGFR tyrosine kinase inhibitors. *Cancer Cell* 2006;10:25–38. [PubMed: 16843263]
20. Debnath J, Muthuswamy SK, Brugge JS. Morphogenesis and oncogenesis of MCF-10A mammary epithelial acini grown in three-dimensional basement membrane cultures. *Methods* 2003;30:256–68. [PubMed: 12798140]
21. Chen HC, Appeddu PA, Isoda H, Guan JL. Phosphorylation of tyrosine 397 in focal adhesion kinase is required for binding phosphatidylinositol 3-kinase. *J Biol Chem* 1996;271:26329–34. [PubMed: 8824286]
22. Wang SE, Xiang B, Guix M, et al. Transforming growth factor beta engages TACE and ErbB3 to activate phosphatidylinositol-3 kinase/Akt in ErbB2-overexpressing breast cancer and desensitizes cells to trastuzumab. *Molecular and cellular biology* 2008;28:5605–20. [PubMed: 18625725]
23. Giancotti FG, Ruoslahti E. Integrin signaling. *Science* 1999;285:1028–32. [PubMed: 10446041]
24. Dumont N, Arteaga CL. Targeting the TGF beta signaling network in human neoplasia. *Cancer Cell* 2003;3:531–6. [PubMed: 12842082]
25. Massague J, Gomis RR. The logic of TGFbeta signaling. *FEBS Lett* 2006;580:2811–20. [PubMed: 16678165]
26. Muraoka RS, Dumont N, Ritter CA, et al. Blockade of TGF-beta inhibits mammary tumor cell viability, migration, and metastases. *The Journal of clinical investigation* 2002;109:1551–9. [PubMed: 12070302]
27. Muraoka-Cook RS, Kurokawa H, Koh Y, et al. Conditional overexpression of active transforming growth factor beta1 in vivo accelerates metastases of transgenic mammary tumors. *Cancer Res* 2004;64:9002–11. [PubMed: 15604265]
28. Janda E, Lehmann K, Killisch I, et al. Ras and TGF[beta] cooperatively regulate epithelial cell plasticity and metastasis: dissection of Ras signaling pathways. *J Cell Biol* 2002;156:299–313. [PubMed: 11790801]
29. McLean GW, Carragher NO, Avizienyte E, Evans J, Brunton VG, Frame MC. The role of focal-adhesion kinase in cancer - a new therapeutic opportunity. *Nat Rev Cancer* 2005;5:505–15. [PubMed: 16069815]

Figure 1.

TGF- β induces clustering of HER2 and integrin $\alpha 6$, $\alpha 3$, $\beta 1$ and $\beta 4$ at the cell membrane. **A.** Matrix and matrix-associated proteins were prepared from MCF10A/HER2 cells grown on 100-mm culture dish as described in (17). The preparations were separated by SDS-PAGE and the identities of the proteins were determined by peptide mass mapping and single-stage MS as indicated in Materials and Methods. **B.** MCF10A/HER2 cells grown on culture dishes were treated with TGF- β (2 ng/ml) or equal amount of PBS (vehicle control) for 24 h in growth medium. At the end of treatment, cells were harvested by trypsinization and stained with antibodies against integrin $\alpha 6$, $\alpha 3$, $\beta 1$ and $\beta 4$, following by a secondary FITC-conjugated antibody. Stained cells were then analyzed by flow cytometry for the level of FITC. **C.** MCF10A/HER2 cells were serum-starved for 16 h before treated with TGF- β (2 ng/ml) for 1 or 24 h. At the end of the treatment, cells were lysed for IP using antibodies against HER2, integrin $\alpha 6$, $\beta 1$ or $\beta 4$. The pull-downs were subjected to immunoblot with the indicated antibodies. **D.** MCF10A/HER2, MDA-MB-453 and BT-474 cells were serum-starved for 16 h and treated with TGF- β or PBS (vehicle control) for 1 h before subjected to matrix preparation or cell surface biotinylation as described in Materials and Methods. Prepared matrix proteins, the streptavidin pull-down for surface-located proteins, and the whole cell lysate were subjected to immunoblot with HER2 antibody. Brefeldin A (10 μ g/ml) was added to cells 1 h prior to TGF- β in the indicated lanes.

Figure 2.

TGF- β activates Src-FAK, leading to receptor clustering and activation of PI3K-Akt. **A.** MCF10A/HER2 cells were serum-starved for 16 h and treated with TGF- β for indicated time before subjected to IP using antibodies against integrin β 1 and HER2. The pull-downs and whole cell lysate were subjected to immunoblot with the indicated antibodies. **B.** MCF10A/HER2 cells were serum-starved for 16 h and pretreated with PP2 (5 μ M) or DMSO (vehicle control) for 1 h before treated with TGF- β or PBS (vehicle control) for 1 h and subjected to IP and immunoblot. **C.** MCF10A/HER2 cells grown on 6-well plates were transiently transfected with siRNA oligonucleotides targeting FAK or a control sequence. Seventy-two h after transfection, cells were serum-starved for 16 h and treated with TGF- β or PBS (vehicle control) for 1 h before subjected to IP and immunoblot. Bands were quantified using ImageJ with untreated cell samples (first lanes) set as 1.0.

Figure 3.

TGF-β induces Src-FAK through EGFR activation. **A.** MCF10A/HER2 cells were serum-starved for 16 h and pretreated with cetuximab (10 μg/ml) or PBS (vehicle control) for 1 h before treated with TGF-β or PBS (vehicle control) for 1 h and subjected to IP and immunoblot. **B.** MCF10A/HER2 cells grown on 6-well plates were transfected by siRNA oligonucleotides targeting human TACE or a control sequence. To rescue TACE expression, a plasmid encoding the full-length mouse TACE [TACE(HA)] or empty vector (vec) were co-transfected with the siRNA oligonucleotides. Seventy-two h after transfection, cells were serum-starved for 16 h and treated with TGF-β or PBS (vehicle control) for 1 h before subjected to IP and immunoblot. **C.** MCF10A/HER2 cells were serum-starved for 16 h and pretreated with PP2 or DMSO (vehicle control) for 1 h before treated with TGF-α (10 ng/ml) or PBS (vehicle control) for 1 h and subjected to IP and immunoblot. Bands were quantified using ImageJ with untreated cell samples (first lanes) set as 1.0.

Figure 4.

Integrin α 6 is essential for the clustering of HER2, integrin β 1 and FAK. **A**. MCF10A/HER2 cells grown on 6-well plates were transiently transfected with siRNA oligonucleotides targeting integrin α 6 or a control sequence. Seventy-two h after transfection, cells were serum-starved for 16 h and treated with TGF- β or PBS (vehicle control) for 1 h before subjected to IP and immunoblot. **B**. MCF10A/HER2 cells were transiently transfected with siRNA targeting integrin α 6 or the control sequence and grown on 6-well plates for 2 days to a 100% confluence. Cell monolayers were serum-starved for 16 h before wounding and the addition of 2 ng/ml TGF- β . Wound closure was captured at 24 h. Bar equals to 50 μ m. **C**. MCF10A/HER2 cells that had been transfected with siRNA targeting integrin α 6, FAK or the control sequence for 48 h were changed to serum-free medium $\pm$ TGF- β . After 48 h, cells were harvested for trypan blue staining. Each data point represents the mean $\pm$ S.D. of 3 wells. **D**. MCF10A/HER2 cells that had been transfected with siRNA targeting integrin α 6, FAK or the control sequence for 48 h were seeded at 2.5×10^4 cells/well on laminin (LN)- or collagen (COL)-coated transwells and allowed to invade toward serum-free medium $\pm$ TGF- β . At 16 h, the cells invaded through transwell filters were counted; each bar represents the mean $\pm$ S.D. of 3 wells. Bands were quantified using ImageJ with untreated cell samples (first lanes) set as 1.0.

Figure 5.

TGF- β induces association of HER2 and integrin with the cytoskeleton. **A.** MCF10A/HER2 cells were serum-starved for 16 h and pretreated with PP2 or lapatinib (1 μ M) or DMSO (vehicle control) for 1 h before treated with TGF- β or PBS (vehicle control) for 1 h and subjected to IP and immunoblot. **B.** MCF10A/HER2 cells grown on 6-well plates were infected by adenoviruses encoding the dominant negative (DN) PI3K p85 subunit or control viruses expressing β -gal (MOI 1:5). Sixteen h after infection, cells were serum-starved for 16 h and treated with TGF- β or PBS (vehicle control) for 1 h before subjected to IP and immunoblot. **C.** MCF10A/HER2 cells were serum-starved for 16 h and pretreated with cytochalasin D (5 μ M) or DMSO (vehicle control) for 1 h before treated with TGF- β or PBS (vehicle control) for 1 h and subjected to IP and immunoblot. **D.** TGF- β induces membrane-proximal clustering of HER2 and integrins by activating Src-FAK and receptor association to cytoskeleton. Signals from extracellular TGF- β are transduced into cells through TGF- β receptors. Activated T β RI induces phosphorylation of TACE, resulting in its translocation to the cell surface, where TACE cleaves EGFR pro-ligands. EGFR ligands will initiate autocrine and paracrine EGFR

signaling, which is amplified in HER2-overexpressing cells, leading to Src-FAK activation and cytoskeleton rearrangement; both are required for the membrane-proximal clustering of HER2 and integrin.

Figure 6.

Inhibition of Src-FAK reverses TGF- β -induced resistance to trastuzumab. BT-474 cells were plated in Matrigel in 8-well chambers and allowed to grow in the absence or presence of TGF- β (2 ng/ml) and PP2 (1 μ M) as indicated. Trastuzumab were added to the top medium 12 h after cell seeding. The inhibitors were replenished every 3 days. Phase contrast images shown were recorded 9 days after the initial seeding of cells. Bar equals 50 μ m. Lower panel: 9-day acini were trypsinized and total cell number was determined. Each bar graph represents the mean $\pm$ S.D. of 4 wells.