Published in final edited form as:

Org Lett. 2008 November 6; 10(21): 4771–4774. doi:10.1021/ol8018506.

B,O-Chelated Azadipyrromethenes as Near-IR Probes

Aurore Loudet[†], Rakeshwar Bandichhor[†], Kevin Burgess^{†,*}, Aniello Palma[‡], Shane O. McDonnell[‡], Michael J. Hall[‡], and Donal F. O'Shea[‡]

[†]Texas A & M University, Chemistry Department, P.O. Box 30012, College Station, Texas 77842, USA.

[‡]Centre for Synthesis and Chemical Biology, School of Chemistry and Chemical Biology, University College Dublin, Belfield, Dublin 4, Ireland.

Abstract

$\lambda_{\text{max fluor}}$ up to 782 nm
(MePh)

X = H, Br
R = H, OMe
Ar = Ph, 4-BrC₆H₄

Complementary synthetic routes to a new class of near IR fluorophores are described. These allow facile access (four synthetic steps) to the core fluorophore and substituted derivatives with emissions between 740 and 780 nm in good quantum yields.

Surprisingly few fluorescent probes emit in the near-IR region with high quantum yields.^{1, 2} Such compounds are valuable for intracellular imaging because autofluorescence in cells tends to obscure emissions at wavelengths below approximately 550 nm, but this factor becomes less of an issue at longer wavelengths. Probes that emit in the 750–900 nm region are, therefore, relatively easy to visualize *in vivo*.³ Cyanine dyes are currently the most widely used probes for this wavelength region, but their quantum yields and photostabilities are sub-optimal.⁴ Thus there is need for new fluorescent probes that emit efficiently above 750 nm.

burgess@tamu.edu and donal.f.oshea@ucd.ie.

Supporting Information **Available** Experimental procedures and characterization data for the new compounds reported. This material is available free of charge via the Internet at <http://pubs.acs.org>.

Dyes based on the 5-6-5 fused BODIPY ring system **A** are popular in biotechnology because they tend to have relatively sharp fluorescence emission characteristics, and high quantum yields.⁵ However, the emission wavelengths of most BODIPY dyes are between 530 and 630 nm, and this is sub-optimal for intracellular or tissue imaging. Previous work by Burgess *et al.*⁶⁻⁸ and others,^{9,10} successfully explored a modification strategy for shifting BODIPY-fluorescence emissions to the red (Figure 1). Adaptation of the core fluorophore scaffold by intramolecular B-O ring formation to produce the 6,6-5-6-5-6,6 ring system **B** gave a *ca* 65 nm bathochromic shift (compare **C** and **D**),¹¹ but the fluorescence emission wavelengths were less than 700 nm.¹²

O'Shea and co-workers have demonstrated that BF₂ chelated tetraaryl-substituted azadipyrrromethenes **E**, fluoresce with wavelength maxima between 670-720 nm with *para*-oriented electron donating groups being particularly effective in enhancing the red-shifted fluorescence, *eg* **F** (Figure 2).¹³⁻¹⁹ This is close to the 750 - 1,400 nm region at which light permeation through tissue is most effective. Subsequently, it was shown that derivatives with extended aromatic groups do emit in near-IR region.²⁰ The premise of this paper is that it would be advantageous to “hybridize” the structure **B** and **E** to shift their emissions beyond 750 nm. On the basis of the structures shown in Figure 1 it was therefore logical to introduce an intramolecular *B,O*-chelate on a tetraaryl aza-BODIPY scaffold *eg* **G**. The studies described here feature compound **1** of this type; it emerges that they can emit in this region.

Scheme 1 shows one approach to the synthesis of the target class **1** from the precursors **2a-d** which can be synthesized in four steps from commercially available materials.²¹ A solid state X-ray structure determination of **2a** illustrated that both 2-MeOC₆H₄ rings lie out of the plane of the central chromophore with torsion angles of 42.7° and 61.4° (Supporting Information). Compounds **1** were obtained from the corresponding **2** by treatment with boron tribromide in dichloromethane. We anticipated that demethylation of the methoxy groups with BBr₃ to the corresponding bisphenols would lead to spontaneous cyclization giving the target materials **1**. This proved correct for substrates **1a** and **1b** with products isolated in 61 and 36% yields respectively following chromatography. Furthermore, in addition to **1a** and **1b**, compounds corresponding to bromination of the pyrrole ring *i.e.* **1e-f** were also observed. This is somewhat surprising since this did not occur for synthesis of the **B** class, but it is also potentially useful if the compounds are to be modified further.⁶ Systems **1c** and **1d** were not air-stable, and were not isolated.

Prior experience with aza-BODIPY derivatives such as **F** indicated that electron releasing R-groups might red-shift the fluorescence maxima of the *B,O*-chelated compounds **1**. An alternative route, which avoids the aryl demethylation conditions used in Scheme 1, was therefore developed to allow inclusion of 4-methoxy groups (Scheme 2). This required conjugate addition of nitromethane to the chalcones **3a, g, h**. Heating of **4a, g, h** with an ammonium source gave phenolic substituted azadipyrrromethenes **5a, g, h**. Fluorophores **1a, g, h** were obtained in a one-pot BF₂ chelation and intramolecular phenolic oxygen – fluorine displacement thereby generating the structurally constraining benzo{1,3,2}oxazaborinine rings. Presumably this synthetic route could be modified to include functional groups in place of the *p*-methoxy position of **1g,h** for bioconjugation. Single crystal X-ray structures for **1h** and **1f** were obtained (Figure 3 and Supporting Information). The benzo fused 6,6-5-6-5-6,6 ring structure was shown to be tetrahedral around the boron atom. Consequently the molecule is chiral just as for the other *B,O*-chelates **B**.

Figure 4 highlights the dramatic spectroscopic contrast between the constrained derivative **1a** and its unconstrained precursor **2a**. Restrictions caused by the *B,O* bonds in **1a** gave bathochromic shifts in both the absorption (86 nm) and emission maxima (58 nm) and over a seven fold increase in the quantum yield from 0.07 to 0.51 (Figure 4, Table 1, entries 1 and 2).

These trends were observed for all derivatives of **1** with additional hypso- and bathochromatic shifts due to the further influence of the aryl and pyrrole substituents.

Introduction of bromine- (**1b**) and methoxy (**1h**) aryl-substituents resulted in a 14 nm bathochromic and 4 nm hypsochromic shift for fluorescence maxima respectively when compared to **1a** (entries 3 and 7). The longest absorbance/emission wavelength maxima (765/782 nm) were observed for derivative **1g** containing electron donating methoxy substituents on the benzo{1,3,2}oxazaborinine rings (entry 6). This pronounced red-shift for fluorescence maximum of 94 nm when compared to **2a** and 36 nm from **1a** places it into the near-IR spectral region (Figure 5). Interestingly, the tetrabromo derivative **1f** also had red-shifted absorption and emission profiles and the substituent did not result in a pronounced quenching of the fluorescence (entry 5).¹⁴

In addition the absorbance spectra showed only a small sensitivity to solvent polarity. For example, the $\lambda_{\text{max abs}}$ of **1g** in DMF is only red shifted by 5 nm when compared to 761 nm in cyclohexane (Table 2). Inspection of fluorescence maxima for **1g** and **1h** revealed a small blue shift trend in polar solvents (Table 3 and Supporting Information). Large Stokes shift values of 34-56 nm in various solvents were recorded for **2a**, however this trend did not translate to the conformationally restricted fluorophores **1a, g** (9 and 19 nm, Table 3).

In conclusion, two efficient synthetic routes to a new class of near-IR fluorophore **1** has been accomplished. The highly favorable emission wavelengths and quantum yields are strongly indicative of future applications in biotechnology. The ease at which key functional groups (halogen and ether) can be included on the core fluorophore indicate that future structural manipulation would be possible to allow adaption for specific use as *in vitro* and non-invasive *in vivo* imaging agents.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgment

Dr Burgess. thanks the members of the TAMU/LBMS-Applications Laboratory directed by Dr Shane Tichy for assistance with mass spectrometry and Dr Nattamai Bhuvanesh for assistance with crystallographic data. Support for this work was provided by The National Institutes of Health (GM72041) and by The Robert A. Welch Foundation. Dr O'Shea thanks funding support from the Program for Research in Third-Level Institutions administered by the HEA, Dr. D. Rai of the CSCB Mass Spectrometry Centre and Dr. H. Mueller-Bunz of the UCD Crystallographic Centre.

References

1. Sun C, Yang J, Li L, Wu X, Liu Y, Liu S. J. Chromatogr., B 2004:173, 90.
2. Kiyose K, Kojima H, Nagano T. Chem.--Asian J 2008:506, 515.
3. Frangioni JV. Curr. Opin. Chem. Biol 2003:626, 34. [PubMed: 14580568]
4. Benson RC, Kues HA. J. Chem. Eng. Data 1977:379, 83.
5. Boyer JH, Haag AM, Sathyamoorthi G, Soong ML, Thangaraj K, Pavlopoulos TG. Heteroat. Chem 1993:39, 49.
6. Chen J, Burghart A, Derecskei-Kovacs A, Burgess K. J. Org. Chem 2000:2900, 2906. [PubMed: 10814176]
7. Thivierge C, Bandichhor R, Burgess K. Org. Lett 2007:2135, 2138. [PubMed: 17455941]
8. Loudet A, Burgess K. Chem. Rev 2007:4891, 4932. [PubMed: 17924696]
9. Umezawa K, Nakamura Y, Makino H, Citterio D, Suzuki K. J. Am. Chem. Soc 2008:1550, 1551. [PubMed: 18193873]

10. Ulrich G, Ziesel R, Harriman A. *Angew. Chem. Int. Ed* 2008:1184, 1201.
11. Kim H, Burghart A, Welch MB, Reibenspies J, Burgess K. *Chem. Commun* 1999:1889, 90.
12. Burghart A, Kim H, Welch MB, Thoresen LH, Reibenspies J, Burgess K, Bergstroem F, Johansson LBA. *J. Org. Chem* 1999:7813, 7819.
13. Killoran J, Allen L, Gallagher JF, Gallagher WM, O'Shea DF. *Chem. Commun* 2002:1862, 1863.
14. Gorman A, Killoran J, O'Shea C, Kenna T, Gallagher WM, O'Shea DF. *J. Am. Chem. Soc* 2004:10619, 31. [PubMed: 15327320]
15. McDonnell SO, Hall MJ, Allen LT, Byrne A, Gallagher WM, O'Shea DF. *J. Am. Chem. Soc* 2005:16360, 16361. [PubMed: 16305199]
16. Killoran J, O'Shea DF. *Chem. Commun* 2006:1503, 1505.
17. Hall MJ, Allen LT, O'Shea DF. *Org. Biomol. Chem* 2006:776, 780. [PubMed: 16493459]
18. O'Shea, D.; Killoran, J.; Gallagher, W. Compounds useful as photodynamic therapeutic agents. US Pat. 7220732, 20030324. 2003.
19. Killoran J, McDonnell SO, Gallagher JF, O'Shea DF. *New J. Chem* 2008:483, 489.
20. Zhao W, Carreira EM. *Angew. Chem., Int. Ed* 2005:1677, 9.
21. Loudet A, Bandichhor R, Wu L, Burgess K. *Tetrahedron* 2008:3642, 3654.

Figure 1. Comparison of fluorescence emission wavelength maxima and quantum yields (in CHCl_3) for: **A** BODIPY class and a B-O ring-extended class **B**.

Figure 2. Representative fluorescence emission wavelength maximum and quantum yield (in CHCl_3) for **E** BF_2 -chelated tetraarylazadipyrromethenes and a new B-O ring extended class **G**.

Scheme 1.
Synthesis of Near-IR Fluorophores **1**

Scheme 2. Four steps synthesis to Near-IR Fluorophores **1**

Figure 3.

X-Ray crystal structure of **1h**. Selected bond lengths *B-O* 1.471(2) and 1.476(2); *B-N* 1.513(2) and 1.514(2); bond angles *O-B-O* 108.0(1)° and *N-B-N* 103.9(1)°.

Figure 4. Normalized absorbance (solid line) and emission (dashed line) for **2a** (blue) and **1a** (red) in CHCl_3 .

Figure 5. Normalized absorbance (a); and (b) emission spectra (excited at the respective λ_{max}) of **1a**, **b**, **e**, **f**, **g** and **h** in toluene

Table 1

Photophysical Properties of **1** in Toluene.

entry	λ_{max} (nm)	λ_{r} (nm)	fwhm (nm) ^a	Φ_{f}^b
1	2a	688	43	0.07 ¹⁰
2	1a	728	36	0.51
3	1b	745	36	0.46
4	1e	747	38	0.35
5	1f	755	36	0.36
6	1g	765	40	0.18 ^c
7	1h	728	36	0.17 ^c

^a Full width at half-maximum height.

^b Measured in 1% pyridine in toluene. ZnPtc (Φ_{f} 0.30) was used as standard

^c Measured in CHCl₃.

Table 2

Effects of the solvent polarity on the absorption.

	$\lambda_{\text{max abs}}$ (fwhm) ^a / nm				
		CHCl ₃	THF	Dioxane	Cyclohexane
					DMF
2a		642 (67)	643 (68)	636 (67)	643 (71)
1a		725 (33)	724 (34)	724 (34)	725 (36)
1g		764 (43)	764 (47)	764 (42)	766 (47)

^a Full width at half-maximum height.

Table 3

Effects of the solvent polarity on the fluorescence.

	$\lambda_{\text{max abs}}$ (Stokes shift) / nm			
	CHCl ₃	THF	Dioxane	Cyclohexane
2a	686 (44)	682 (39)	681 (45)	674 (34)
1a	737 (12)	738 (14)	738 (14)	733 (9)
1g	778 (14)	782 (18)	778 (14)	771 (10)
				696 (56)
				738 (13)
				785 (19)