Published in final edited form as:

Dev Dyn. 2008 September ; 237(9): 2415–2429. doi:10.1002/dvdy.21686.

Dynamic expression patterns of G protein-regulated inducer of neurite outgrowth 1 (GRIN1) and its colocalization with Gao implicate significant roles of Gao-GRIN1 signaling in nervous system

Ikuo Masuho^{1,2,*}, Yasumasa Mototani^{1,2,3,*}, Yoshinori Sahara^{2,3}, Junko Asami², Shun Nakamura^{2,4}, Tohru Kozasa^{1,§}, and Takayoshi Inoue²

¹ Department of Pharmacology, College of Medicine, University of Illinois at Chicago, IL 60612, USA

² Department of Biochemistry and Cellular Biology, National Institute of Neuroscience, NCNP, Tokyo 187-8502, Japan

Abstract

GRIN1 (Gprin 1) is a signaling molecule coexpression of which with constitutively active form of Gao can stimulate neurite extensions in Neuro2a cells, yet its *in vivo* roles remain elusive. Here we examine expression profiles of *GRIN1* during mouse development by *in situ* hybridization (ISH) and immunohistochemistry. ISH analysis revealed that *GRIN1* expression was limited to the nervous system at all developmental stages tested: in the central nervous system, *GRIN1* expression occurred within the entire embryonic mantle zones, while it became restricted to sets of nuclei at postnatal to adult stages. Immunohistochemistry using a GRIN1 specific antibody demonstrated that GRIN1 colocalized with Gao at neuronal dendrites and axons, but it was not detected in glial cells. These results suggest that Gao-GRIN1 pathway could mediate significant roles in neuronal migration and differentiation at embryonic stages and exert functions in wiring and/or maintenance of specific neural circuitries at postnatal to adult stages.

Keywords

GRIN1; Gprin1; Gao; gene expression; neural development; mouse

Introduction

G protein-coupled receptors (GPCRs) represent a large superfamily of cell-surface receptors (>1,000 members in human), being involved in a variety of physiological processes such as perception of sensory information, modulation of synaptic transmission, hormone release/actions, regulation of cell contraction/migration, and cell growth/differentiation. In the signaling cycles, agonist occupancy of GPCRs promotes the activation of heterotrimeric G proteins (G proteins) by catalyzing the exchange of GDP for GTP on the α subunit ($G\alpha$). GTP-bound $G\alpha$ and free $\beta\gamma$ subunits ($G\beta\gamma$) then separately activate downstream effectors, including adenylyl cyclases, phospholipases, RhoGEFs and ion channels. Hydrolysis of GTP

*These authors equally contributed to this work.

§To whom correspondence should be addressed: Tohru Kozasa, Department of Pharmacology, College of Medicine, University of Illinois, 909 South Wolcott Avenue, Chicago, IL 60612. Tel.: 312-413-0111; E-mail: tkozas@uic.edu

³Present address: Department of physiology, Tsurumi University School of Dental Medicine, Kanagawa 230-8501, Japan

⁴Tokyo University of Agriculture and Technology, Biotechnology and Life Science, Tokyo 184-8588, Japan

to GDP on $G\alpha$ promotes both dissociation of $G\alpha$ from effectors and reassociation of $G\alpha$ with $G\beta\gamma$, so that termination of the signaling cycle is dependent on the GTPase activity of $G\alpha$ (for review, see Hepler and Gilman, 1992; Neer, 1995).

G protein classes are defined according to the primary sequences of their α subunits, resulting in four main families: Gai/o , Gas , $Gaq/11$, and $G\alpha_{12/13}$. One type of $G\alpha$ subunit, Gao , is extremely abundant in the mouse brain (Strathmann et al., 1990) and constitutes about 0.5% of the membrane protein of bovine adult brain (Huff et al. 1985). In particular, it is highly enriched in growth cone membranes isolated from the fetal rat brain (Strittmatter et al., 1990). Despite its abundance in the nervous system and growth cone, functional roles of Go signaling pathway in the nervous system is still poorly characterized. At the molecular level, GPCR-mediated signaling through Go is known to be primarily mediated by its $G\beta\gamma$, yet direct involvement of Gao for effector regulation has not been clearly demonstrated, especially under the *in vivo* condition (reviewed by Wettschureck and Offermanns, 2005). At the cellular level, Go has been implicated in neuronal differentiation, but its molecular mechanism has not been well defined (Doherty et al., 1991; Strittmatter et al., 1994; Ghil et al., 2000).

To identify novel effectors of α subunits of Gi/o subfamily, we have taken advantage of the fact that Gaz can be phosphorylated by protein kinase C at a site near its amino terminus that does not interfere with interaction between Gaz and its effector, adenylyl cyclase (Kozasa and Gilman, 1996). Using the far-western method with ^{32}P -phosphorylated, $GTP\gamma S$ -bound Gaz , we have originally identified G protein-regulated inducer of neurite outgrowth 1 (GRIN1) as an effector candidate of Gai/o from a mouse embryo cDNA library (Chen et al., 1999). Homologues of GRIN1, GRIN2 and GRIN3, were further identified by database search (Chen et al., 1999; Iida and Kozasa 2004). Among different mouse tissue/organs examined, GRIN1 protein and mRNA were only copious in the brain (Chen et al., 1999). Endogenous GRIN1 protein was predominantly membrane-bound and was highly enriched in the growth cone membrane fraction of mouse embryonic brains like Gao protein (Chen et al., 1999). Although GRIN1 did not contain any domains that are homologous to other known signaling motifs, it specifically recognized activated Gai/o through amino acid residues 716 to 746 and 797 to 827 of GRIN1 (Chen et al., 1999; Nakata et al., 2005). Interestingly, coexpression of GRIN1 with the constitutively active mutant of Gao promoted the extension of neurites in mouse neuroblastoma Neuro2a cells, possibly through activation of Cdc42 small G protein (Nakata et al., 2005). These results implicated that the Gao -GRIN1 interaction might play a pivotal role in regulating neurite outgrowth.

As an initial approach to understand *in vivo* roles of GRIN1, we extensively defined distribution patterns of GRIN1 transcripts from the mouse embryonic to adult stages by *in situ* hybridization. In addition, we determined distribution patterns of GRIN1 protein in the mouse brain and its subcellular localization in primary cultured neurons by immunohistochemistry. We also compared distribution patterns of GRIN1 protein with those of Gao protein in the mouse brain and primary cultured neurons. All the results from these analyses demonstrated that *GRIN1* mRNA was primarily expressed in migrating and/or differentiating neurons and that GRIN1 and Gao proteins shared similar subcellular localization in neuronal cells, supporting the idea that Gao -GRIN1 pathway plays important roles in the neural development and/or neural circuit formation.

Results

Differentiating neuronal cells widely express *GRIN1* mRNA both in the central and peripheral nervous system during mouse embryogenesis

In order to comprehensively determine temporal and spatial expression patterns of *GRIN1* during mouse embryogenesis, we applied *in situ* hybridization (ISH) technique that allows sensitive and reliable detection of mRNAs. Since hybridization probes may not deeply penetrate into thick samples, we performed ISH in whole-mount preparations for samples up to the embryonic day (E)10.5: for samples older than E10.5, we made transverse slices of embryos/brains with ~500 µm thickness to maximize efficiency of probe hybridization and washings. Specificity of ISH was evaluated by parallel application of sense and antisense riboprobes. Our *GRIN1* sense probes did not yield any detectable staining throughout the experiments (data not shown). An exception was the case when both sense and antisense probes were trapped within the embryonic brain vesicles, producing non-specific staining in the whole-mount preparations. In such cases, we identified specific signals by comparing sense to antisense signals. Under these conditions, we were able to obtain reproducible ISH results as follows:

Expression of *GRIN1* mRNA was first detected at E9.5 within a confined region of the central nervous system (CNS) as well as subgroups of cells in the peripheral nervous system (PNS; Fig. 1C), while no intense expression was found at E8.5-9.0 (Fig. 1A and B). In the CNS, the earliest expression emerged within the ventral domain of the spinal cord at E9.5 (Fig. 1C), and this *GRIN1* mRNA positive area gradually spread into the dorso-lateral region of the spinal cord by E10.5 (Fig. 1D). Transverse slices through upper body of the E10.5 embryo demonstrated that the signal was localized within the ventral horn (Fig. 1F). At E12.5, *GRIN1* mRNA was detected in the whole area of the spinal cord except for the ventricular zone (Fig. 2C and D). This expression pattern continued during embryonic stages (Fig. 2F, 3H and 3P). Within the brain, clear *GRIN1* signals were visible in the restricted developmental units such as the ventral midbrain and rhombic lip at E10.5 (Fig. 1D and E), whereas nonspecific background staining was accumulated in the brain ventricles from E9.5 onward in whole-mount preparations (Fig. 1C). Intense signals were also detectable in the diencephalon and telencephalon at E11.5 (Fig. 2A). While the dorsal portion of the telencephalon (i.e. the future cerebral cortex) had no *GRIN1* signal until E12.5 (Fig. 2A and C), many of *GRIN1* positive cells emerged progressively to fill the area at E14.5 (Fig. 2E). The appearance of *GRIN1* mRNA in the olfactory bulb was observed at E12.5 (Fig. 2D), and the expression was maintained until adult stage (Fig. 2F, 3A, 3I, 4A and 5A). Dynamic features of *GRIN1* expression profiles in the CNS well correlated to patterns of neuronal migrations and/or maturations. For instance, *GRIN1* expression in the spinal cord started from the ventral region and this signal expanded dorsally to eventually cover entire domain, which well reflected patterns of neuronal maturations in the spinal cord (Nornes and Das, 1972). Also, the developmental changes of *GRIN1* positive domains in the telencephalon appear to mirror the migration pattern of interneurons in telencephalic vesicles, where interneurons originate in the basal domain and migrate dorsally to fill the cerebral cortex at around E.12.5 (Anderson et al., 1997).

In the PNS, the expression of *GRIN1* mRNA was first detected at E9.5 within the cranial ganglia, such as the trigeminal, facio-acoustic, glossopharyngeal and vagal ganglia, and within the dorsal root ganglia at the trunk level (Fig. 1C). The *GRIN1* signals increased in all these regions at E10.5 (Fig. 1D) and these signals were maintained until E14.5 (Fig. 2A-F). The staining signal was also observed in the sympathetic ganglia at E11.5 (data not shown) and the expression was sustained until E14.5 (Fig. 2D and F). In the neural retina, the moderate level of expression was detected at E12.5 (Fig. 2D) and E14.5 (Fig. 2F), whereas no expression was observed in other sensory organs, tongue, olfactory epithelium and

primordium of follicle of vibrissa, during embryonic stages tested (data not shown). It should be noted that positive signals were localized primarily in the differentiating neurons, while emigrating/migrating neural crest cells did not show any *GRIN1* expression both in the cranial and trunk regions.

Most importantly, no *GRIN1* expression was observed in the tissue/organs outside of the nervous system. In addition, both in the CNS and PNS, the expression was restricted in the differentiating cell populations at all developmental stages tested: cells in the ventricular zone and neural crest generating zone (i.e. roof plate) did not show any *GRIN1* expressions during mouse embryogenesis.

***GRIN1* mRNA expression is restricted to differentiating fields in the mouse developing brain**

To further examine the expression profiles of *GRIN1* mRNA during brain development, we prepared ~500 µm slices of E14.5 to adult brains and subjected them to ISH. Cells expressing *GRIN1* mRNA were found in almost entire differentiating fields in the embryonic CNS (Fig. 3), while they again became restricted to specific regions from the postnatal to adult stages (Fig. 4 and 5). The ISH results are detailed below for each brain subdivision.

Forebrain—*GRIN1* mRNA was expressed in ventral region of the telencephalon and diencephalon of E14.5 embryos (Fig. 3A-C). At this stage, the septum, striatum, piriform cortex, and thalamus strongly expressed *GRIN1* mRNA. The olfactory bulb, hippocampus, amygdala, and hypothalamus also expressed *GRIN1* mRNA but at relatively weak level. The earliest expression in the developing cerebral cortex occurred at the lateral region by E14.5, while the medial region exhibited no expression at this stage. *GRIN1* expression was not observed in the ventricular zones of the cortex and the ganglionic eminences.

To follow changes of *GRIN1* expression domains in the forebrain during development, we further examined the expression at later stages, E16.5, postnatal day 2 (P2), and adult. At E16.5, *GRIN1*-positive cells became expanded into the dorsal area of the telencephalon, so that *GRIN1* expression were found in almost all areas of the forebrain except for the ventricular zone and ganglionic eminence (Fig. 3I-K). At this stage, *GRIN1* expression persisted at a high level within the septum, striatum, piriform cortex, and thalamus, and at moderate level in the amygdala and hypothalamus. Distinct *GRIN1* expression appeared in the lateral migratory stream and nucleus of the lateral olfactory tract. Hybridization signals were greatly increased at E16.5 in the olfactory bulb, cerebral cortex, and hippocampus. Especially, significant developmental change was observed in the cerebral cortex: *GRIN1*-positive cells emerged to fill the medial cortex.

After birth, *GRIN1* expression was reduced from the ventral areas in the forebrain (Fig. 4B and C). At P2, the reduction was observed in the septum, striatum and hypothalamus. In the septum and striatum, only weak expression was detectable. In the hypothalamus, the reduction of *GRIN1* expression was significant in the lateral and anterior hypothalamus. In contrast, dorsal areas of the hypothalamus, the paraventricular nucleus and zona incerta maintained conspicuous expression of *GRIN1*. Furthermore, high level of *GRIN1* expression was maintained in lateral and dorsal areas of the forebrain; the olfactory bulb, cerebral cortex, piriform cortex, hippocampus, amygdala, and thalamus exhibited strong *GRIN1* expression (Fig. 4A-C).

At the adult stage, *GRIN1* expression was greatly decreased in the entire forebrain and was restricted to specific regions (Fig. 5A-C). The strong signals were persisted in the olfactory bulb, piriform cortex, and hippocampal formation. Weak signals were observed in the

cerebral cortex and amygdala. Thus, it is significant to note that, in the adult forebrain, robust *GRIN1* expression is confined to the areas where maturation of newborn neurons takes place.

Midbrain—In the midbrain, *GRIN1* expression was widely distributed at E14.5 (Fig. 3D-F). Moderate *GRIN1* signal was observed in the tegmentum, and weak signals were noticed in the superior and inferior colliculi. At E16.5, the *GRIN1* expressions increased slightly (Fig. 3L-O). The superior colliculus, inferior colliculus and tegmentum showed moderate expressions. At this stage, the periaqueductal gray and red nucleus could be identified due to their strong signals. These expressions were greatly reduced after birth. At P2, weak expressions were observed in the superior colliculus, inferior colliculus and tegmentum (Fig. 4E and F). Weak but clear expression was present in the periaqueductal gray. At the adult stage, *GRIN1* expressions were hardly detectable within the midbrain (Fig. 5E and F). Notably, no *GRIN1* expression was observed in the midbrain ventricular zone at all developmental stages examined.

Hindbrain—*GRIN1*-positive regions included almost whole areas of the hindbrain except for the ventricular zone at E14.5 (Fig. 3E-G). The pons, medulla and cerebellum exhibited moderate *GRIN1* signals. However, the expression in the anterior pons was barely detectable (data not shown). At this stage, some nuclei and migratory streams in the pons and medulla were identified. The cochlear nuclei were strongly positive, and conspicuous signals were detected in the trigeminal motor nucleus, posterior precerebellar extramural migration and precerebellar intramural migration. Some other nuclei of the hindbrain were positive, but these remain to be unidentified.

At E16.5, *GRIN1* expression was expanded into the anterior pons, so that moderate expressions were observed in the whole areas of the hindbrain except for the ventricular zone (Fig. 3M-O). Conspicuous signals were observed in the locus coeruleus, tegmental nucleus, cochlear nuclei, trigeminal nucleus complex, facial motor nucleus and superior olivary nucleus. Strong signals were also noticed in the pontine nuclei and anterior precerebellar extramural migration. However, the *GRIN1* expression level as a whole started to decrease from this stage.

After birth, most of these expressions were greatly decreased. At P2, clear but weak expression was recognizable in the cerebellum and little expressions were observed in the pons and medulla (Fig. 4E-G). At this stage, only pontine nuclei showed a strong signal. At the adult stage, *GRIN1* expression was hardly detectable in the hindbrain except for granule cell layer of the cerebellum (Fig. 5E-G). Again, *GRIN1* expression was not observed in the ventricular zone of the fourth ventricle at all developmental stages tested.

Spinal cord—*GRIN1* was strongly expressed in all areas except for ventricular zone at E14.5 and E16.5 (Fig. 3H and P). However, this expression was greatly reduced at P2 (Fig. 4H) and virtually no signals were detected at the adult stage (Fig. 5H).

Taken together, *GRIN1* expression was observed mostly in differentiating fields of the developing CNS. For instance, *GRIN1* expression in the striatum was high at E14.5 and E16.5 and then was decreased by P2, which well correlated with the time course of neurogenesis in the striatum (for review, see Bayer and Altman, 2004). On the other hand, proliferative ventricular zones exhibited no *GRIN1* expression throughout the neuroaxis at all developmental stages examined, supporting the notion that *GRIN1* mRNA expression has a physiological significance in the differentiating fields (see Discussion section for details).

A GRIN1 specific antibody labels *GRIN1* mRNA positive cell populations in the adult mouse brain that substantially coexpressed Gao protein

In order to determine sites of actions for GRIN1 protein, we next performed immunohistochemical staining on 30 μ m sections of adult mouse brain by utilizing an anti-GRIN1 antibody against the oligopeptides containing GRIN1 specific sequences. We have previously confirmed the specificity of this GRIN1 antibody for immunostaining through the assay of competition with the immunogenic peptides or GRIN1 protein (Chen et al., 1999). Also, pre-immune antiserum yielded no staining (data not shown). The specificity of the antibody was further demonstrated by Western blot analysis (Fig. 6): the antibody reacts specifically with an apparent molecular weight of about 110 kDa in the mouse adult brain extract, while no significant signal was detected from the liver extract. Simultaneously, we examined colocalization of GRIN1 with Gao using an anti-Gao antibody. We also performed ISH on 14 μ m sections of adult mouse brain to rigorously compare the distribution of *GRIN1* mRNA and its protein.

First of all, we found that the overall labeling pattern of immunostaining with our anti-GRIN1 antibody in mouse adult brain was similar to that obtained in our ISH analysis. For instance, immunohistochemical labeling of the olfactory bulb, piriform cortex, amygdala, hippocampus, cerebrum and cerebellum by the GRIN1 specific antibody was noted with prominent intensity (Fig. 7G-L), which was consistent with results from ISH analysis (Fig. 5A-G and 7A-F). Within these GRIN1 positive areas, conspicuous immunostaining for Gao was detected (Fig. 7M-R), and considerable overlaps between GRIN1 and Gao staining were observed at various regions of the adult brain (Fig. 7S-X). The followings are summaries of detailed expression profiles by each area.

Olfactory bulb—In the olfactory bulb, ISH positive cells were detected within confined sets of layers (Fig. 5A and 7A). In the main olfactory bulb (MOB), the granular cell layer and mitral cell layer exhibited strong GRIN1 mRNA expression, while only weak expression occurred in the glomerular layer. In the accessory olfactory bulb (AOB), the granule cell layer showed strong expression signal, whereas only weak signal was detectable in the external plexiform layer and glomerular layer. On the other hand, GRIN1 immunoreactivity was observed in almost all layers of the AOB and MOB (Fig. 7G). Most layers yielded strong signals, yet there was relatively weak immunostaining in the glomerular layers of the AOB and MOB. Also, weak immunostaining was found in the mitral cell somata of the MOB where strong signal of *GRIN1* mRNA was detected. The comparison of ISH and immunohistochemical results suggested that GRIN1 protein might be localized to axons and/or dendrites of ISH-positive cells. In contrast to GRIN1, Gao immunoreactivity was enriched in the AOB and glomerular layer of the MOB in Fig. 7M. However, Gao was distributed in almost all layers of the olfactory bulb and was localized outside of the mitral cell somata, which was similar to GRIN1 immunostaining pattern. Hence distribution pattern of GRIN1 protein was essentially overlapped with that of Gao protein (Fig. 7S).

Piriform cortex—In the piriform cortex, *GRIN1* antisense RNA probes strongly labeled cells in the pyramidal cell layer (Fig. 7B), while the GRIN1 antibody marked regions surrounding the pyramidal cell somata (Fig. 7H). This suggested that GRIN1 protein might be transported into axons and/or dendrites of the pyramidal cells. Gao protein also appeared to be accumulated in axons and/or dendrites of the pyramidal cells (Fig. 7N).

Amygdala—Weak yet prominent *GRIN1* mRNA expression was found in the amygdala (Fig. 7C). Relatively strong ISH signals were observed in the lateral, basolateral and medial

amygdala (Fig. 5C). Same was the case for GRIN1 proteins in this brain region (Fig. 7I). Gao proteins appeared to be colocalized with GRIN1 proteins (Fig. 7O, U).

Hippocampus—In the hippocampal formation, GRIN1 antibody immunostained most layers in the hippocampus and dentate gyrus (Fig. 7J). Consistent with ISH analysis (Fig. 5C and 7D), the immunoreactivity of GRIN1 was quite intense in the CA1 and dentate gyrus. The stratum moleculare of the dentate gyrus, the stratum oriens and stratum radiatum of the CA1 were strongly labeled. Weaker yet distinct labeling was further observed along the stratum oriens and stratum radiatum of the CA2/3 region. On the other hand, the stratum lacunosum moleculare showed weak protein expressions of GRIN1. Although *GRIN1* mRNA was observed throughout the stratum pyramidale of the CA1-CA3 and the stratum granulosum of the dentate gyrus, relatively weak immunoreactivities for GRIN1 were found in somata of the pyramidal and granule cells, again implicating axonal and dendritic localization of GRIN1 protein.

Those cells expressed GRIN1 protein in the hippocampal regions primarily coexpressed Gao protein. However, we found a subtle difference between the protein distribution patterns of GRIN1 and Gao (Fig. 7V). Compared to other areas in the hippocampal formation, the stratum lacunosum moleculare showed weak GRIN1 immunoreactivity, while Gao immunoreactivity in the area was relatively moderate. Additionally, GRIN1 immunoreactivity in the stratum moleculare of the dentate gyrus and CA1 region was highest in the hippocampal formation, while Gao immunoreactivity in the stratum moleculare of the dentate gyrus was much weaker than that in the CA1 region.

Cerebrum—In the cerebral cortex, considerable number of cells expressed *GRIN1* mRNA without any layer and region specificity at the adult stage (Fig. 7E). This was also supported by our immunostaining results (Fig. 7K). Gao proteins showed a similar distribution pattern to GRIN1 proteins (Fig. 7Q, W).

Cerebellum—The cerebellum was another brain region in which moderate labeling was detected by ISH at the adult stage. *GRIN1* mRNA was concentrated in the granule cell layer in the cerebellum (Fig. 5E-G and 7F). This expression pattern was quite similar to that previously reported for *Gao* mRNA (Brann et al., 1987). Anti-GRIN1 antibody predominantly immunostained the granule cell layer. Moderate staining with this antibody was also found in the molecular layer (Fig. 7L). On the other hand, Gao immunoreactivity was highly concentrated in the molecular layer and moderate immunoreactivity was present in the granule cell layer as previously reported (Worley et al., 1986; Asano et al., 1987). Thus, *GRIN1* and *Gao* mRNAs were expressed in cells along the granule cell layer but their proteins exhibited colocalization with different gradients between the molecular layer and granule cell layer (Fig. 7X).

Colocalization of GRIN1 and Gao at cellular level—To rigorously examine whether the immunoreactivities of GRIN1 colocalize with those of Gao, we captured images by using the confocal microscope (Fig. 8). Consequently, we found apparent colocalization of the GRIN1 and Gao staining signals in neurons. For instance, those neurons in the hippocampal CA1, layer V of the cerebral cortex, and granular layer of the cerebellar cortex coexpressed GRIN1 and Gao. Of all those areas examined in Fig. 7, GRIN1 and Gao immunoreactivities were primarily colocalized at the regions outside of the neuronal cell somata, i.e., in neuropil and on the surface of their soma. Only faint labeling of GRIN1 was observed in the cell somata (Fig. 8B).

Collectively, overall staining patterns in the adult brain sections for *GRIN1* mRNA and protein were parallel. In addition, our data suggest that GRIN1 protein would be transported

into the axons and/or dendrites, since GRIN1 immunoreactivity was mainly observed outside of neuronal cell somata where *GRIN1* mRNA signals were strongly detected. Furthermore, GRIN1 and Gao were essentially colocalized at the cellular level in the adult mouse brain (Fig. 8), suggesting cooperative roles of GRIN1 and Gao in the mature nervous system.

GRIN1 protein is specifically expressed in neurons and localized in the dendrite and axon with Gao protein

Our ISH results demonstrated that *GRIN1* mRNA was expressed in neuron-enriched regions of the developing and adult brains. To ensure the type of cells expressing GRIN1, we prepared monolayer cultures from the cerebellum of mouse P7 brain. The cerebellar cells were then analyzed by double immunostaining using antibodies to GRIN1 and MAP2 (neuronal marker) or GFAP (glial marker). As shown in Fig. 9A and 9B, GRIN1 immunoreactivities were preferentially found in MAP2-immunoreactive cells but not in GFAP-immunoreactive cells. These results demonstrate that GRIN1 is a neuron-specific protein.

Our immunoreactive mapping of GRIN1 protein on brain sections in Fig. 7 and 8 suggested that the protein might localize at the axons and/or dendrites of neuronal cells. In order to further examine the distribution of GRIN1 protein in neuronal cells, we doubly immunostained primary culture from the cerebellum with antibodies to GRIN1 and marker proteins, such as MAP2 for dendrites, neurofilament for axons, or synaptophysin for synapses. Double immunostaining of 10 days-cultured cells with typical neurites clearly demonstrated colocalization of GRIN1 with MAP2 and neurofilament, and partially with synaptophysin (Fig. 9B, C and D). We also found that GRIN1 and Gao shared the same subcellular distribution in these cerebellar cells (Fig. 9E). Similar analyses were performed using the hippocampal cultures with essentially the same results (data not shown). These observations clearly indicate that GRIN1 protein is specifically expressed in neurons and arranged along the dendrites, axon, and synapses with its interacting partner, Gao.

Discussion

We have previously reported the cloning and initial characterization of GRIN1 in *in vitro* contexts (Chen et al., 1999; Nakata et al., 2005). In this study, we examined the expression profiles of GRIN1 *in vivo* and found that *GRIN1* mRNA expression was confined within the differentiating field of the mouse nervous system over the course of embryonic and postnatal development. This enrichment of GRIN1 mRNA in the differentiating field, together with neuronal specific expression and dendritic/axonal localization of GRIN1 protein, suggests that GRIN1 plays important roles in differentiation and arrangements of neuronal cells.

Spatiotemporal expression pattern of *GRIN1* well correlated with differentiation waves of neuronal cells, implicating roles of GRIN1 in neuronal differentiation

Spatiotemporal patterns of neurogenesis in the rat nervous system have been characterized extensively (for review, see Bayer and Altman, 2004). Given that rat development is generally ~1.5 day behind the mouse, *GRIN1* expression profiles determined in the mouse brain well reflect patterns of neurogenesis mapped during rat brain development. For instance, in the rat hippocampal formation, neurogenesis begins at E16 with the generation of pyramidal cells. The granule cells in the dentate gyrus are noted for their late time of origin and most granule cells are generated after birth in rat. Likewise, *GRIN1* mRNA expression was first detected in the mouse hippocampal formation at E14.5, and the distinct expression in the mouse dentate gyrus was observed at P2 and adult stages. *GRIN1*

expression in the mouse nervous system tended to be highest in the developing fields at stage coincident with the peak of neurogenesis.

In addition to the spatial and temporal features of *GRIN1* expression patterns, following observations supported that *GRIN1* is preferentially expressed along the postmitotic yet immature neurons. First, earliest hybridization signals were detectable in the restricted developmental units such as the cranial ganglia and ventral spinal cord at E9.5 when first neurons start to migrate and differentiate (Easter et al., 1993; Mastick and Easter, 1996). Second, the proliferative ventricular zones exhibited no expression throughout the neuroaxis at all developmental stages tested. Third, in the adult brain, prominent ISH signals were detected in the olfactory bulb, piriform cortex, and dentate gyrus where neuronal differentiation takes place (Pekcec et al., 2006; for review, see Abrous et al., 2005). These observations indicated that *GRIN1* probably may not play any roles in developmental decisions of neuroblast cells, since it only appeared in the differentiating field of the nervous system. Instead, *GRIN1* is likely to function in signal transduction pathways involved in neuronal differentiation.

Similar expression profiles between *GRIN1* and *Gao* suggests physiological significance of *Gao-GRIN1* signaling in neuronal cells

Our previous reports have shown that *GRIN1* is specifically expressed in the brain and is enriched in the nerve growth cones (Chen et al., 1999). In this study, we demonstrated that mouse *GRIN1* transcript appeared in the maturing postmitotic neurons from E9.5 and was maintained until the adult stage. Among members of *Gai/o* subfamily, only *Gao* appeared to harbor similar expression profiles to *GRIN1* (Strathmann et al., 1990; Strittmatter et al., 1990; Schmidt et al., 1994; Morishita et al., 1999). We also confirmed that *GRIN1* protein is colocalized with *Gao* protein in the section of the adult brain and that *GRIN1* and *Gao* share the dendritic, axonal and synaptic localization in the primarily cultured neurons. In addition to the *in vivo* and *in vitro* similarity of expression profiles between *GRIN1* and *Gao*, coexpression of *GRIN1* with a constitutively active mutant of *Gao* promotes neurite extension in a neuronal cell line (Chen et al., 1999; Nakata et al., 2005), supporting the notion that *Gao-GRIN1* interaction plays physiologically significant roles in the nervous system.

We have previously shown that *GRIN1* also interact with *Gai* members *in vitro* (Chen et al., 1999). Our data do not exclude interactions of *GRIN1* with *Gai* members *in vivo*. For instance, in the main olfactory bulb where *Gao* exhibits only weak expression, *GRIN1* may function with *Gai* subunits.

Possible functions of *Gai/o-GRIN1* signaling in the mouse developing nervous system

Landmarks of neuronal development include controlled birth of immature neurons from neuronal precursors, their directed migrations from the birthplaces to final positions, neurite outgrowth and guidance of the motile growth cone towards postsynaptic partners, and finally the generation and/or maintenance of synapses along functional neuronal circuits. The enrichment of *GRIN1* in the developing fields of the nervous system along with its time course of expression in the elaborated brain organizations suggested that *GRIN1* could be more or less involved in these processes of neural development.

Although the results need to be confirmed by double staining with markers for migratory neurons, our ISH analysis suggested that *GRIN1* was expressed in a migratory population of neurons: *GRIN1* mRNA expression in the rhombic lip and posterior precerebellar extramural migration implicates that the *GRIN1* might be involved in dorsoventral migration from the caudal rhombic lip to the precerebellar nuclei of the brainstem (Wang et al., 2005).

Furthermore, abundant *GRIN1* expression was observed in the olfactory bulb and dentate gyrus of the adult brain, where migrations and incorporations of immature neurons were continuously processed. Importantly, *Gai/o*-coupled receptor signaling, in which *GRIN1* might participate, is known to regulate directed neuronal migrations. *GABA_B* receptors, for example, harbor a critical modulatory role in the migration of cortical interneurons and affect the length of their leading process (López-Bendito et al., 2003). *SDF-1/CXCR-4* signalings can also control migration of the hippocampal, cerebellar and cortical neurons (Bagri et al., 2002; Zhu et al., 2002; Stumm et al., 2003). Taken together, it is likely that *Gai/o-GRIN1* signaling has regulatory function in directed migration of immature neurons.

Several lines of evidence further argue for a specific role of *Gai/o* signaling in regulating neurite outgrowth and axon guidance. *Go* is a major G protein subclass accumulated in the growth cone membrane (Strittmatter et al., 1990). It was also shown that the transfection of PC12 cells with constitutively active *Gao* mutant doubles the neurite length and increase the number of neurites (Strittmatter et al., 1994). *N-CAM*- and *N-cadherin*-induced neurite outgrowth from PC12 cells can be further blocked by pertussis toxin, a specific inhibitor for *Gai/o* (Doherty et al., 1991). *Gao* also modulates neurite outgrowth by regulating the signaling pathway initiated by cyclic AMP in F11 neuroblastoma cells (Ghil et al., 2000). Many neurotransmitters have also been reported to control neurite growth and axonal guidance: *D2* dopamine, serotonin *1B*, *CB1* cannabinoid, and *CXCR4* receptors regulates neurite growth (Reinoso et al., 1996; Lotto et al., 1999; Jordan et al., 2005); *CXCR4* and *GABA_B* receptors guide nerve growth cones (Xiang et al., 2002); and *CXCR4* and *5-HT_{1B/1D}* receptors modulates axonal responsiveness to guidance cues (Chalasani et al., 2003; Bonnin et al., 2007). A common feature among these receptors is that they couple and function through the *Gai/o* pathway. These findings, together with our observations that *GRIN1* protein was localized along the dendrites, axons and synapses, and that *Gao-GRIN1* signaling pathway could regulate neurite growth/extension (Chen et al., 1999; Nakata et al., 2005), lead us to conclude that *Gai/o-GRIN1* interaction may influence cellular differentiation through *GPCR-Gai/o*-mediated signaling for neurite growth and nerve growth cone guidance, which would bring effects on neuronal circuit formation/maintenance in the developing and adult brains. Such a hypothesis will be further examined by generating *GRIN1* gene knockout mice or using *RNAi* approach for *GRIN1* in the future studies.

Experimental Procedures

Animals

All animal experiments in this study conform to Japanese governmental guidelines and have been approved by the Animal Care and Use Committee of the National Institute of Neuroscience, Japan. *ICR* (CD-1) mice (Charles River, Japan) were mated, and embryos at various stages were collected. The day of appearance of the vaginal plug was taken as *E0* and the day of birth was regarded as *P0*. *C57BL/6* mice (Charles River, Japan) were used for immunohistochemistry and primary culture of cerebellar cells.

Antibodies

For *GRIN1* protein detection, a polyclonal antibody against *GRIN1* specific peptide corresponding to amino acid residues 550 to 567 (Chen et al. 1999) was used. Anti-GFAP (glial fibrillary acidic protein) monoclonal antibodies (clone G-A-5, Sigma, MO, USA), anti-MAP2 (microtubule-associated protein 2) monoclonal antibodies (clone HM-2, Sigma, MO, USA), anti-neurofilament monoclonal antibodies (clone 2H3, Developmental Studies Hybridoma Bank, IA, USA), anti-synaptophysin monoclonal antibodies (clone SVP-38, Sigma, MO, USA) and anti-*Gao* monoclonal antibodies (clone 2A.3, BIOMOL, PA, USA) were applied to mark distinct cell types and cellular compartments.

A Plasmid Construct

A 3.6kb mouse GRIN1 cDNA fragment was released from the pCMV5-GRIN1 expression vector (Chen et al., 1999) by using *KpnI* and *HindIII* restriction enzymes. The 3.6kb fragment was then cloned into *KpnI-HindIII* sites of pBluescriptII KS(-) (Stratagene, CA, USA), yielded pBS-GRIN1.

Synthesis of Hybridization Probes

Sense or antisense GRIN1 probes labeled with digoxigenin (Dig)-11-UTP (Roche Diagnostic GmbH, Penzberg, Germany) were prepared from pBS-GRIN1. For whole-mount *in situ* hybridization of *GRIN1*, pBS-GRIN1 linearized by *Asp718* was transcribed with T7 RNA polymerase for antisense probes (Invitrogen, CA, USA) and that linearized by *SpeI* was transcribed with T3 RNA polymerase for sense probes (Roche Diagnostic GmbH, Penzberg, Germany). These probes cover the full-length coding region of GRIN1. For section *in situ* hybridization of *GRIN1*, pBS-GRIN1 was digested at the *XhoI* site, which cleaved 3 prime untranslated region of *GRIN1* cDNA, generating a template for the antisense RNA probes. T7 RNA polymerase was then used to transcribe the riboprobes only covering the 956bp 3 prime untranslated region of *GRIN1* gene, whose DNA sequences are specific for *GRIN1* among the mouse homologues. These RNA probes were purified by G-50 Sephadex column (Roche Diagnostic GmbH, Penzberg, Germany), and stored at -80°C.

Whole-Mount In Situ Hybridization

Embryos were dissected out to remove extraembryonic membranes in Tyrode's solution and fixed overnight in 4% paraformaldehyde (PFA) in PBS at 4°C. Fixed embryos were washed with phosphate-buffered saline (PBS) containing 0.1% Tween 20 (PBT) and bleached with 5:1 methanol/30% hydrogen peroxide for 6 hours at room temperature. Bleached embryos were dehydrated into 100% methanol and stored at -20°C. For sliced samples, we fixed embryos and brains overnight in PBS containing 4% PFA at 4°C and bleached them for 6 hours at room temperature in 5:1 methanol/30% hydrogen peroxide. After rehydration in a graded series of methanol/PBT, they were embedded in 2% agarose/PBS, and sectioned into 500-μm slices by micro slicer (D.S.K. DTK-3000, Kyoto, Japan). Slices were dehydrated into 100% methanol and stored at -20°C. Whole-mount *in situ* hybridization was performed as described previously (Conlon et al., 1992; Inoue et al., 1997; Inoue et al., 1998).

Western Blot Analysis

The liver and brain were dissected out from a 10 weeks-old C57BL/6 female mouse. Approximately 0.6 mg of tissue was sonicated in 5 volume of RIPA buffer (50 mM Tris-HCl, pH 8.0, 150 mM NaCl, 1% Nonidet P-40, 0.5% sodium deoxycholate, 0.1% SDS). Debris was removed by centrifugation at 20,000 x g for 30 min at 4°C. Supernatants of tissue extracts were subjected to protein analysis using a DC Protein Assay (BioRad, CA, USA). 10 μg protein (per lane) of tissue extracts were separated on 10% SDS-polyacrylamide gel, and transferred onto nitrocellulose membrane. Non-specific binding sites were blocked with 5% skim milk in PBT for 30 min at room temperature. The blot was incubated overnight at 4°C with anti-GRIN1 antibody (at 1:2,000 dilution with 5% skim milk in PBT). After washing with PBT, horseradish-peroxidase-conjugated anti-rabbit IgG was applied as the secondary antibody (at 1:5,000 dilution with 5% skim milk in PBT) for 45 min at room temperature. Finally, the blot was incubated with ECL plus Western blotting detection reagents (Amersham Pharmacia Biotech, IL, USA) and chemiluminescence was detected with an imager (LAS-1000; Fuji Film, Tokyo, Japan).

Section In Situ Hybridization

Section *in situ* hybridization was performed as described previously (Funatsu et al., 2004).

Primary Culture of the Cerebellar Cells

The cerebella were dissected out from 7-day-old mouse pups and minced into 1 mm slices in dissection buffer (10 mM HEPES, 137 mM NaCl, 5.3 mM KCl, 33 mM glucose, 45 mM sucrose). Then, the minced cerebella were incubated in dissection buffer containing 20 U/ml papain (Worthington, NJ, USA) for 20 min at 37°C. The tissues were washed three times with neuronal medium (Minimum Essential Medium (Invitrogen, CA, USA), 6 mM glucose, 2 mM L-glutamine, 0.1% Serum Extender (Becton Dickinson, MA, USA), 10% horse serum (Invitrogen, CA, USA), 20 mM KCl), and dissociated by trituration with a fire-polished Pasteur pipette. The cerebellar cells (500 cells/mm²) were seeded onto glass coverslips coated with poly-L-lysine (Sigma, MO, USA) and collagen (Vitrogen, CA, USA), and were cultured in neuronal medium. After 3 days in culture, 100 nM 5-Fluoro-2'-deoxyuridine (Sigma, MO, USA) and 250 nM uridine (Sigma, MO, USA) were added in cultured medium for suppression of glial cells proliferation, and cells were incubated for 24 hours. After 6 days, the cells were used for the experiments.

Immunohistochemistry

Under anesthesia, adult C57BL/6 mice were perfused through the ascending aorta with 4% PFA in PBS. The whole brains were removed, postfixed in 4% PFA in PBS overnight, and immersed in 30% sucrose in PBS for 48 hours at 4°C. To prepare cryosections, brains were placed in OCT compound and snap-frozen on dry ice. Cryosections (30~40 µm in thickness) were prepared and free-floating immunohistochemistry were performed. The sections were treated with 0.5% Triton X-100 in PBS for 30 min, and blocked with 5% skim milk in PBS for 2 hours. After the blocking, sections were incubated with 5% skim milk/PBS containing anti-GRIN1 antibody (1:300) and anti-Gao antibody (1:200) for 48 hours at 4°C. After washing, the sections were incubated with Alexa 594 goat anti-rabbit IgG antibody (1:1000) and Alexa 488 goat anti-mouse IgG antibody (1:500) in PBS containing 5% skim milk at 4°C overnight. After serial washings, sections were mounted onto adhesive slides and coverslipped with PermaFluor Mountant Medium (Shandon Immunon, PA, USA).

For immunohistochemistry on primarily cultured cells, the cerebellar cells at 10 days *in vitro* (DIV) were fixed with 4% PFA in PBS at 4°C for 20 min. Cells were permeabilized with 0.5% Triton X-100 in PBS for 5 min at room temperature. Cells were then incubated with 5% skim milk in PBS for 1 hour at room temperature, followed by incubation for 48 hours at 4°C in 5% skim milk/PBS containing anti-GRIN1 antibody (1:300) plus antibodies against marker proteins such as anti-GFAP antibody (1:750), anti-MAP2 antibody (1:500), anti-neurofilament antibody (1:300), anti-synaptophysin antibody (1:500) or anti-Gao antibody (1:300). After washing, cells were incubated with Cy5 goat anti-rabbit IgG antibody (1:1000) and Alexa 488 goat anti-mouse IgG antibody (1:500) for 90 min at room temperature. After immunoreaction, cells were mounted with PermaFluor Mountant Medium. Fluorescent images were captured using fluorescence microscopes (Axiophot2, Carl Zeiss Jena GmbH, Jena, Germany; DM5000B, Leica, Wetzlar, Germany) equipped with CCD cameras (ORCA-ER, Hamamatsu, Japan; DFC300FX, Leica, Wetzlar, Germany) or a confocal laser scanning microscope (TCS SP2, Leica, Wetzlar, Germany).

Identification of GRIN1 Positive Regions

For anatomical identification of GRIN1 positive regions, we referred to atlases by Paxinos and Franklin (2001), Kaufmann (1992), and Altman and Bayer (1995).

Acknowledgments

We thank Dr. N. Yokoyama for his support in the initial phase of this study, Dr. M. Yokosuka for helpful discussions and Dr. S. Kohsaka for allowing us to use their stereoscopic microscope. This study was supported in part by grants from the National Institutes of Health (GM61454 and NS4144 to T. Kozasa), and the Program for Promotion of Fundamental Studies in Health Sciences of the National Institute of Biomedical Innovation (05-32 to S. Nakamura & T. Inoue).

Grant sponsor: National Institutes of Health; Grant numbers: GM61454 and NS4144; Grant sponsor: National Institute of Biomedical Innovation; Grant number: 05-32.

References

- Abrons DN, Koehl M, Moal ML. Adult neurogenesis: from precursors to network and physiology. *Physiol Rev.* 2005; 85:523–569. [PubMed: 15788705]
- Altman, J.; Bayer, SA. *Atlas of Prenatal Rat Brain Development.* CRC Press; Boca Raton: 1995. p. 660
- Anderson SA, Eisenstat DD, Shi L, Rubenstein JLR. Interneuron Migration from Basal Forebrain to Neocortex: Dependence on *Dlx* Genes. *Science.* 1997; 278:474–476. [PubMed: 9334308]
- Asano T, Semba R, Ogasawara N, Kato K. Highly sensitive immunoassay for the alpha subunit of the GTP-binding protein Go and its regional distribution in bovine brain. *J Neurochem.* 1987; 48:1617–1623. [PubMed: 3104540]
- Bayer, SA.; Altman, J. Development of the telencephalon: neural stem cells, neurogenesis, neuronal migration. In: Paxinos, G., editor. *The Rat Nervous System.* Academic Press; San Diego: 2004. p. 27-73.
- Bagri A, Gurney T, He X, Zou YR, Littman DR, Tessier-Lavigne M, Pleasure SJ. The chemokine SDF1 regulates migration of dentate granule cells. *Development.* 2002; 129:4249–4260. [PubMed: 12183377]
- Bonnin A, Torii M, Wang L, Rakic P, Levitt P. Serotonin modulates the response of embryonic thalamocortical axons to netrin-1. *Nat Neurosci.* 2007; 10:588–597. [PubMed: 17450135]
- Brann MR, Collins RM, Spiegel A. Localization of mRNAs encoding the alpha-subunits of signal-transducing G-proteins within rat brain and among peripheral tissues. *FEBS Lett.* 1987; 222:191–198. [PubMed: 3115828]
- Chalasani SH, Sabelko KA, Sunshine MJ, Littman DR, Raper JA. A Chemokine, SDF-1, reduces the effectiveness of multiple axonal repellents and is required for normal axon pathfinding. *J Neurosci.* 2003; 23:1360–1371. [PubMed: 12598624]
- Chen LT, Gilman AG, Kozasa T. A candidate target for G protein action in brain. *J Biol Chem.* 1999; 274:26931–26938. [PubMed: 10480904]
- Conlon RA, Rossant J. Exogenous retinoic acid rapidly induces anterior ectopic expression of murine Hox-2 genes in vivo. *Development.* 1992; 116:357–368. [PubMed: 1363087]
- Doherty P, Ashton SV, Moore SE, Walsh FS. Morphoregulatory activities of NCAM and N-cadherin can be accounted for by G protein-dependent activation of L- and N-type neuronal Ca²⁺ channels. *Cell.* 1991; 67:21–33. [PubMed: 1680564]
- Easter SS Jr, Ross LS, Frankfurter A. Initial tract formation in the mouse brain. *J Neurosci.* 1993; 13:285–299. [PubMed: 8423474]
- Funatsu N, Inoue T, Nakamura S. Gene expression analysis of the late embryonic mouse cerebral cortex using DNA microarray: identification of several region- and layer-specific genes. *Cereb Cortex.* 2004; 14:1031–1044. [PubMed: 15142957]
- Ghil SH, Kim BJ, Lee YD, Suh-Kim H. Neurite outgrowth induced by cyclic AMP can be modulated by the alpha subunit of Go. *J Neurochem.* 2000; 74:151–158. [PubMed: 10617116]
- Hepler JR, Gilman AG. G proteins. *Trends Biochem Sci.* 1992; 17:383–387. [PubMed: 1455506]
- Huff RM, Axton JM, Neer EJ. Physical and immunological characterization of a guanine nucleotide-binding protein purified from bovine cerebral cortex. *J Biol Chem.* 1985; 260:10864–10871. [PubMed: 3928624]

- Iida N, Kozasa T. Identification and biochemical analysis of GRIN1 and GRIN2. *Methods Enzymol.* 2004; 390:475–483. [PubMed: 15488195]
- Inoue T, Chisaka O, Matsunami H, Takeichi H. Cadherin-6 expression transiently delineates specific rhombomeres, other neural tube subdivisions, and neural crest subpopulations in mouse embryos. *Dev Biol.* 1997; 183:183–94. [PubMed: 9126293]
- Inoue T, Tanaka T, Suzuki SC, Takeichi M. Cadherin-6 in the developing mouse brain: expression along restricted connection systems and synaptic localization suggest a potential role in neuronal circuitry. *Dev Dyn.* 1998; 211:338–351. [PubMed: 9566953]
- Jordan JD, He JC, Eungdamrong NJ, Gomes I, Ali W, Nguyen T, Bivona TG, Philips MR, Devi LA, Iyengar R. Cannabinoid receptor-induced neurite outgrowth is mediated by Rap1 activation through G α i-triggered proteasomal degradation of Rap1GAP1. *J Biol Chem.* 2005; 280:11413–11421. [PubMed: 15657046]
- Kaufman, MH. *The Atlas of Mouse Development.* Academic Press; San Diego: 1992. p. 512
- Kozasa T, Gilman AG. Protein kinase C phosphorylates G12 α and inhibits its interaction with G $\beta\gamma$. *J Biol Chem.* 1996; 271:12562–12567. [PubMed: 8647866]
- López-Bendito G, Luján R, Shigemoto R, Ganter P, Paulsen O, Molnár Z. Blockade of GABA $_B$ receptors alters the tangential migration of cortical neurons. *Cereb Cortex.* 2003; 13:932–942. [PubMed: 12902392]
- Lotto B, Upton L, Price DJ, Gaspar P. Serotonin receptor activation enhances neurite outgrowth of thalamic neurones in rodents. *Neurosci Lett.* 1999; 269:87–90. [PubMed: 10430511]
- Mastick GS, Easter SS Jr. Initial organization of neurons and tracts in the embryonic mouse fore- and midbrain. *Dev Biol.* 1996; 173:79–94. [PubMed: 8575640]
- Morishita R, Shinohara H, Ueda H, Kato K, Asano T. High expression of the γ 5 isoform of G protein in neuroepithelial cells and its replacement of the γ 2 isoform during neuronal differentiation in the rat brain. *J Neurochem.* 1999; 73:2369–2374. [PubMed: 10582595]
- Nakata H, Kozasa T. Functional characterization of G α o signaling through G protein-regulated inducer of neurite outgrowth 1. *Mol Pharmacol.* 2005; 67:695–702. [PubMed: 15585744]
- Neer EJ. Heterotrimeric G proteins: organizers of transmembrane signals. *Cell.* 1995; 80:249–257. [PubMed: 7834744]
- Nornes HO, Das GD. Temporal Pattern of Neurogenesis in Spinal Cord: Cytoarchitecture and Directed Growth of Axons. *Proc Natl Acad Sci USA.* 1972; 69:1962–1966. [PubMed: 4114859]
- Paxinos, G.; Franklin, KBJ. *The Mouse Brain in Stereotaxic Coordinates.* Academic Press; San Diego: 2001. p. 296
- Pekcec A, Loscher W, Potschka H. Neurogenesis in the adult rat piriform cortex. *Neuroreport.* 2006; 17:571–574. [PubMed: 16603913]
- Reinoso BS, Undie AS, Levitt P. Dopamine receptors mediate differential morphological effects on cerebral cortical neurons in vitro. *J Neurosci Res.* 1996; 43:439–453. [PubMed: 8699530]
- Schmidt CJ, Zubiaur M, Valenzuela D, Neer EJ, Drager UC. Go, a guanine nucleotide binding protein, is expressed during neurite extension in the embryonic mouse. *J Neurosci Res.* 1994; 38:182–187. [PubMed: 8078103]
- Strathmann M, Wilkie TM, Simon MI. Alternative splicing produces transcripts encoding two forms of the α subunit of GTP-binding protein Go. *Proc Natl Acad Sci USA.* 1990; 87:6477–6481. [PubMed: 1697681]
- Strittmatter SM, Valenzuela D, Kennedy TE, Neer EJ, Fishman MC. Go is a major growth cone protein subject to regulation by GAP43. *Nature.* 1990; 344:836–841. [PubMed: 2158629]
- Strittmatter SM, Fishman MC, Zhu XP. Activated mutants of the alpha subunit of G(o) promote an increased number of neurites per cell. *J Neurosci.* 1994; 14:2327–2338. [PubMed: 8158271]
- Stumm RK, Zhou C, Ara T, Lazarini F, Dubois-Dalcq M, Nagasawa T, Höllt V, Schulz S. CXCR4 Regulates Interneuron Migration in the Developing Neocortex. *J Neurosci.* 2003; 23:5123–5130. [PubMed: 12832536]
- Wang VY, Rose MF, Zoghbi HY. Math1 expression redefines the rhombic lip derivatives and reveals novel lineages within the brainstem and cerebellum. *Neuron.* 2005; 48:31–43. [PubMed: 16202707]

- Wettschureck N, Offermanns S. Mammalian G proteins and their cell type specific functions. *Physiol Rev.* 2005; 85:1159–1204. [PubMed: 16183910]
- Worley PF, Baraban JM, Dop CV, Neer EJ, Snyder SH. Go, a guanine nucleotide-binding protein: immunohistochemical localization in rat brain resembles distribution of second messenger systems. *Proc Natl Acad Sci USA.* 1986; 83:4561–4565. [PubMed: 3086888]
- Xiang Y, Li Y, Zhang Z, Cui K, Wang S, Yuan XB, Wu CP, Poo MM, Duan S. Nerve growth cone guidance mediated by G protein-coupled receptors. *Nat Neurosci.* 2002; 5:843–848. [PubMed: 12161754]
- Zhu Y, Yu T, Zhang XC, Nagasawa T, Wu JT, Rao Y. Role of the chemokine SDF-1 as the meningeal attractant for embryonic cerebellar neurons. *Nat Neurosci.* 2002; 5:719–720. [PubMed: 12080344]

Figure 1.

Onsets of *GRIN1* mRNA expression at E9.5. Whole-mount *in situ* hybridization (ISH) was carried out on E8.0-E10.5 embryos with an antisense probe for *GRIN1*. (A) Dorsal view of an E8.0-8.5 embryo. (B) Lateral view of an E8.5-9.0 embryo. *GRIN1* expression is not detected until E8.5-E9.0. (C) Lateral view of an E9.5 embryo. *GRIN1* expression commences in the cranial ganglia, dorsal root ganglia and spinal cord. (D, E) Lateral (D) and dorsolateral (E) views of an E10.5 embryo. *GRIN1* expression in the brain appears in the ventral midbrain and rhombic lip. (F) Transverse section through upper body of an E10.5 embryo shows *GRIN1* positive cells in the ventral horn of the spinal cord and dorsal root ganglia. Developmental stages are indicated in the panels. Scale bars, 0.5 mm for (A, B and F), 1 mm (C-E).

Figure 2.

The distribution of *GRIN1* mRNA is restricted to the nervous system during the mouse embryonic development. E11.5, E12.5, and E14.5 mouse embryonic slices examined by whole-mount ISH for *GRIN1* expression are shown. As indicated, the left, middle, and right columns are for E11.5 (A and B), E12.5 (C and B), and E14.5 (E and F) embryos, respectively. The top row shows the levels of embryonic slices. Note that *GRIN1* mRNA expression is restricted to the central and peripheral nervous system. Scale bars, 1 mm.

Figure 3.

GRIN1 mRNA is widely distributed in the developing embryonic brains. E14.5 (A-H) and E16.5 (I-P) brains coronally sliced are subjected to whole-mount ISH. The sectioning levels of E14.5 and E16.5 brains are indicated in the top schematic drawings. Slices are arranged from the rostral forebrain to cervical spinal cord. (G, K and O) Migratory streams are indicated by arrowheads. Scale bars, 1 mm for (A-G and I-O), 0.5 mm for (H and P).

P2

Figure 4.

GRIN1 mRNA expression in the brain decreases at the postnatal stages. Coronal slices from a P2 brain are subjected to whole-mount ISH and are arranged as explained in the legend for Figure 3. After birth, *GRIN1* expression is greatly reduced in caudal and ventral areas of the brain. (A-D) In the forebrain, *GRIN1* mRNA expression is intense in the mitral cell layer and granular cell layer of the olfactory bulb, cerebral cortex, piriform cortex, hippocampus, thalamus, and amygdala. On the other hand, the ventral areas of the forebrain, the septum, striatum and the ventral hypothalamus, exhibit weak or no signal. (D-H) In the midbrain and hindbrain, only the pontine nucleus displays an intense signal and weak signal is present in

the cerebellum, inferior colliculus, superior colliculus and periaqueductal gray. Scale bars, 1 mm for (A and H), 2 mm for (B-G).

Adult

Figure 5.

Expression of *GRIN1* mRNA in the adult brain is at the highest level in the regions where newborn neurons are continuously integrated. Slice samples from an adult brain processed to whole-mount ISH are arranged as in Figure 3 and 4. In the adult brain, the most intense signal for *GRIN1* mRNA is enriched in the dentate gyrus, piriform cortex and mitral cell layer and granular cell layer of the olfactory bulb. The hippocampus also exhibits the intense signal. Weak signals are present in the cerebral cortex, amygdala, and granular layer of cerebellar cortex. Scale bars, 1 mm for (I and P), 2 mm for (J-O).

Figure 6. An anti-GRIN1 antibody used in this study specifically recognizes GRIN1 protein. (A) Extracts prepared from adult mouse liver and brain are separated on 10% SDS-polyacrylamide gel and stained with Coomassie Brilliant Blue. (B) The proteins are transferred to nitrocellulose and subjected to Western blot analysis using anti-GRIN1 antibody. The specific signal corresponding to GRIN1 is indicated by an arrow.

Figure 7.

Distribution patterns of *GRIN1* mRNA, GRIN1 protein and $G\alpha o$ protein in the adult mouse brain. *GRIN1* mRNA expression examined by ISH are compared to the result of immunohistochemistry for GRIN1 and $G\alpha o$ proteins on sections. (A-F) Adult mouse brain sections examined by ISH with the *GRIN1* antisense probe are arranged. (G-R) Sections at similar level with panels A-F doubly immunostained with anti-GRIN1 antibody (G-L) and anti- $G\alpha o$ antibody (M-R) are assembled. (S-X) Merged images of GRIN1 and $G\alpha o$ immunostaining are shown. Characters in merged images indicate regions labeled by immunohistochemistry. Layers in the cerebral cortex are indicated by Roman numerals. Scale bars, 200 μm .

Figure 8.

Confocal images of GRIN1 and G α o immunoreactivities in the adult mouse brain. Adult mouse brain sections are doubly immunostained with anti-GRIN1 antibody (A-C) and anti-G α o antibody (D-F) to examine colocalization of GRIN1 and G α o immunoreactivities under the confocal microscope. (G-H) Merged images of GRIN1 and G α o immunostaining are shown. The hippocampal CA1 region (A, D, and G), layer V of the cerebral cortex (B, E, and H), and granular layer of the cerebellar cortex (C, F, and I) are shown as examples. Neuronal somata are marked with asterisks. Arrowheads indicate the colocalization of GRIN1 and G α o on the surface of neuronal cell somata. Arrows point to the cytoplasmic signals of GRIN1. Scale bars, 30 μ m.

Figure 9.

Identification of GRIN1 immunoreactive cell types and subcellular localization of GRIN1 and *Gαo* proteins in the cerebellar primary culture. (A-E) The double immunofluorescent images of the cerebellar primary culture cells. Immunoreactivity of GRIN1 (upper panels, red fluorescence) is compared to that of glial fibrillary acidic protein (GFAP; A), microtubule-associated protein 2 (MAP2; B), neurofilament (NF; C), synaptophysin (SYP; D), or *Gαo* (E) (lower panels, green fluorescence). The bottom panels show the merged images of both fluorescence signals. Note that GRIN1 expression colocalizes with neuronal markers and *Gαo*. Scale bars, 20 μm for (B-E), 50 μm for (A).

Table 1

List of anatomical abbreviations

V	trigeminal ganglion	Me	medial amygdala
VII	facial ganglion	Mi	mitral cell layer of olfactory bulb
VIII	acoustic ganglion	MoV	trigeminal motor nucleus
IX	glossopharyngeal ganglion	OB	olfactory bulb
X	vagal ganglion	PAG	periaqueductal gray
XI	vagal ganglion	PIM	precerebellar intramural migration
Am	amygdala	Pir	piriform cortex
APEM	anterior precerebellar extramural migration	Pn	pontine nucleus
BLA	basolateral amygdala	PPEM	posterior precerebellar extramural migration
BMA	basomedial amygdala	PVN	paraventricular nucleus
Cb	cerebellum	Pyr	pyramidal cell layer of piriform cortex
CbG	granular layer of cerebellar cortex	R	retina
CbM	molecular layer of cerebellar cortex	RH	rhombencephalon
CbW	white matter of cerebellar cortex	RN	red nucleus
Ce	central amygdala	rRL	rostral rhombic lip
CN	cochlear nucleus	SC	superior colliculus
cRL	caudal rhombic lip	SCG	sympathetic ganglion
Cx	cerebral cortex	SG	stratum granulosum
DG	dentate gyrus	SLM	stratum lacunosum
Di	diencephalon	SM	stratum moleculare
DRG	dorsal root ganglion	SO	stratum oriens
EPI	external plexiform layer of olfactory bulb	SON	superior olivary nucleus
EPIA	external plexiform layer of accessory olfactory bulb	Sp	septum
FMN	facial motor nucleus	SP	stratum pyramidale
Gl	glomerular layer of olfactory bulb	SpC	spinal cord
GIA	glomerular layer of accessory olfactory bulb	SR	stratum radiatum
Gr	granular cell layer of olfactory bulb	St	striatum
GrA	granular cell layer of accessory olfactory bulb	Te	telencephalon
H	hypothalamus	TEG	tegmentum
Hip	hippocampus	Tg	tegmental nucleus
IC	inferior colliculus	Th	thalamus
LA	lateral amygdala	TN	trigeminal nucleus complex
LC	locus coeruleus	VH	ventral horn of spinal cord
LCx	lateral cerebral cortex	VM	ventral midbrain
LMS	lateral migratory stream	ZI	zona incerta
LOT	nucleus of the lateral olfactory tract		