Published in final edited form as:

Schizophr Res. 2008 December ; 106(2-3): 265–267. doi:10.1016/j.schres.2008.08.021.

Sequence analysis of *P21-activated kinase 3 (PAK3)* in chronic schizophrenia with cognitive impairment

Eric M. Morrow, MD, PhD^{a,b}, Anna Kane^a, Donald C. Goff, MD^b, and Christopher A. Walsh, MD, PhD^{a,*}

a Division of Neurogenetics and Howard Hughes Medical Institute, Department of Neurology, Beth Israel Deaconess Medical Center, and Division of Genetics, Children's Hospital Boston, Harvard Medical School, New Research Building 266, 77 Avenue Louis Pasteur, Boston, MA 02115, USA

b Schizophrenia Research Program, Department of Psychiatry, Massachusetts General Hospital, Harvard Medical School, Boston, MA 02114, USA

Abstract

The P21-activated kinase PAK3 is critical for cognitive development and truncating mutations cause non-syndromic mental retardation (MR). Missense mutations are also associated with psychotic disorders, most commonly with schizophrenia involving premorbid MR, namely “pfpopschizophrenie”. We set out to measure the frequency of sequence variants in *PAK3* in schizophrenia without premorbid MR. We conducted complete gene resequencing of all coding exons and exon-intron boundaries in patients with schizophrenia with cognitive impairment but without premorbid MR. Deleterious variants in schizophrenia alone were rare (<1/159 or 0.6%). Thereby, while PAK3 remains a strong biological candidate in psychosis, evidence from human genetics provides strongest support for a link to pfpopschizophrenie and not to schizophrenia without premorbid intellectual disability.

Keywords

Schizophrenia; Mental retardation; Pfpopschizophrenie; P21-activated kinase PAK3; cognitive development

1. Introduction

Abnormalities in GABAergic interneurons (Lewis et al., 2005) and loss of dendritic spines (Kolluri et al., 2005, Rosoklija et al., 2000) are consistent cellular abnormalities in schizophrenia. The p21 (CDKN1A)-activated kinase PAK3 is involved in both interneuron development (Cobos et al., 2007) as well as actin remodeling in dendrites (Hayashi et al., 2004, Kreis et al., 2007). In addition to being a plausible biological candidate in schizophrenia

*Corresponding Author: email: cwals@bidmc.harvard.edu, 617-667-0813 (tel), 617-667-0815 (fax).

Contributors: D.C.G. was involved in recruitment of patient subjects. E.M.M., A.K. and C.A.W. were involved in preparation of DNA, gene sequencing and data analysis. All were involved in hypothesis generation and preparation of this manuscript.

Conflicts of interest: E.M.M. has received grant funding from Pfizer and Merck. Dr. Goff has received research funding or honoraria from, or has been on an advisory board of AstraZeneca, BMS, Cephalon, Cortex, Eli Lilly, GSK, Janssen, Merck, Organon, Pfizer, Solvay and Wyeth. All other authors declare that they have no conflicts of interest.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

susceptibility, there are strong human genetic data that support a role for the X-linked *PAK3* gene in psychotic disorders. In particular, multiple families with *PAK3* mutations have members with psychosis and premorbid mild mental retardation (Gedeon *et al.*, 2003, Peippo *et al.*, 2007, Rejeb *et al.*, 2008), known as “pfpopschizophrenie” (defined in (Doody *et al.*, 1998)). Indeed, we contend that while *DISC1* has received a great deal of attention based largely on the discovery of a highly penetrant translocation in a single extended pedigree (St Clair *et al.*, 1990), highly penetrant mutations in *PAK3* in psychosis have been largely overlooked. *PAK3* is critical for cognitive development and truncating mutations cause moderate, non-syndromic mental retardation (MR) (Allen *et al.*, 1998); however, missense mutations are associated with neurobehavioral disorders (and milder MR) in all pedigrees described (Donnelly *et al.*, 1996, Gedeon *et al.*, 2003, Peippo *et al.*, 2007, Rejeb *et al.*, 2008). Three of four such families have associated psychotic illness (Gedeon *et al.*, 2003, Peippo *et al.*, 2007, Rejeb *et al.*, 2008). Indeed, in an extended pedigree with an A365E mutation, mental retardation was generally mild and at least two members had psychotic illness, one with chronic paranoid schizophrenia and an IQ of 80 (Gedeon *et al.*, 2003).

In this experiment, we set out to measure the frequency of sequence variants in *PAK3* in patients ascertained for schizophrenia without premorbid intellectual disability. We specifically chose a schizophrenia patient population with profound cognitive impairment with the notion that this population may be relatively enriched for *PAK3* mutations.

2. Materials and methods

2.1. Patient ascertainment

Study procedures were approved by the Partners HealthCare and Massachusetts Department of Mental Health institutional review boards. After providing written informed consent, 159 outpatients with schizophrenia (43 females and 116 males) who receive treatment at an urban community mental health center in Boston were enrolled consecutively. The diagnosis of schizophrenia was confirmed by a research psychiatrist using DSM-IV criteria. Inclusion criteria included DSM-IV diagnosis of schizophrenia, age 18–70, ability to complete symptom rating scales and cognitive tests. Exclusion criteria included psychosis secondary to a substance use disorder or a general medical condition. Of enrolled subjects, 120 were Caucasian (75%), 37 were of African descent (23%), one was of East/Southeast Asian descent (1%), and one was of Latino descent (1%). All patients were medicated and stable in terms of psychotic symptoms.

2.2. Cognitive assessment

The Wechsler Adult Intelligence Scale (WAIS-III) was also administered to assess global cognitive status. Subnormal IQ was generally construed to be due to cognitive abnormalities of schizophrenia; however, psychotic symptoms and medication effects cannot be excluded as contributory factors.

2.3. Sequence analysis

All coding exons and exon-intron boundaries in *PAK3* were sequenced. Each amplicon was amplified by PCR and sequenced using capillary methods. Sequence reads were made first manually and secondarily by the *SNP-Detector* software package.

3. Results

In female subjects, IQ ranged from 66 to 137 with six females under IQ of 70 and an overall mean female IQ of 86.3 $\pm$ 16.7. In male subjects, IQ ranged from 59 to 130 with eight males under IQ of 70 and mean male IQ of 89.1 $\pm$ 14.0. While no patients were known to have premorbid MR, cognitive status was significantly impaired and frequently in the mild to

borderline MR range. Coding sequence across *PAK3* is known to be highly invariant (<http://www.ncbi.nlm.nih.gov/projects/SNP/>). There are a total of 242 known SNPs across the 124561 bp genomic interval and all but 1 outside the 1635 open reading frame. One coding non-synonymous SNP (rs1805077) has been reported (but not validated), A489C, which putatively changes Arginine 31 to Serine. On re-sequencing in the schizophrenia patient sample, three total SNPs were discovered. We did not find rs1805077. Two of the SNPs had been previously discovered and were in intronic regions, namely rs16986385 and rs10521535 (Table 1). The single novel SNP discovered in the patient sample occurred in exon 12, substituting C at mRNA position 1786 for a T, and resulting in a synonymous SNP at Leucine 395. This novel SNP was found in a Caucasian female patient with normal IQ who was heterozygous for this sequence change.

4. Discussion

While four mutations in *PAK3* have been reported to cause MR and also to have neurobehavioral disorders in affected members of the pedigree (Donnelly *et al.*, 1996, Gedeon *et al.*, 2003, Peippo *et al.*, 2007, Rejeb *et al.*, 2008), including three with psychotic illness (Gedeon *et al.*, 2003, Peippo *et al.*, 2007, Rejeb *et al.*, 2008), the frequency of potentially causative rare sequence variants in *PAK3* is less than one in 159 cases of schizophrenia alone (less than 0.6%). Variants do not appear to be enriched by selecting for patients with impaired cognitive function. Kraepelin named a schizophrenia subtype in 1907 “pfpopschizophrenie” for schizophrenia engrafted on top of intellectual disability (Doody *et al.*, 1998). Each pedigree thus far reported with *PAK3* mutations has been ascertained for mild to moderate mental retardation as a primary diagnosis. The risk of psychotic illness is elevated in the presence of premorbid intellectual disability (Zammit *et al.*, 2004) but generally quoted as less than 8% (Reid, 1980). As with mental retardation, in schizophrenia, cognitive impairment is a core feature and neurodevelopment abnormalities are prevalent. *PAK3* mutations may represent a genetic susceptibility for pfpopschizophrenie and an important shared biological component between MR and schizophrenia, although our data here indicate that sequence mutations or variants are rare in schizophrenia without premorbid MR. Additional experiments are now needed to define the frequency of *PAK3* sequence variants in schizophrenia with premorbid intellectual disability.

Acknowledgements

We are grateful to the patients who participated in this study. We are grateful for support from the NIMH (1K23MH080954-01 to E.M.M.). E.M.M. holds a Career Award for Medical Scientists from the Burroughs Wellcome Fund and is also grateful for support from the Rappaport Research Scholarship in Neuroscience at MGH. This work was supported by a NARSAD Young Investigator award to E.M.M. who is the Sidney R. Baer Jr. Foundation Investigator. D.C.G. is supported by a mid-career development award from the NIMH (K24 MH002025-06). C.A.W. is an Investigator of the Howard Hughes Medical Institute.

Role of funding source: The funding sources had no further role in study design; in the collection, analysis and interpretation of data; in the writing of the report; and in the decision to submit the paper for publication.

References

- Allen KM, Gleeson JG, Bagrodia S, Partington MW, MacMillan JC, Cerione RA, Mulley JC, Walsh CA. *PAK3* mutation in nonsyndromic X-linked mental retardation. *Nat Genet* 1998;20(1):25–30. [PubMed: 9731525]
- Cobos I, Borello U, Rubenstein JL. Dlx transcription factors promote migration through repression of axon and dendrite growth. *Neuron* 2007;54(6):873–88. [PubMed: 17582329]
- Donnelly AJ, Partington MW, Ryan AK, Mulley JC. Regional localisation of two non-specific X-linked mental retardation genes (MRX30 and MRX31). *Am J Med Genet* 1996;64(1):113–20. [PubMed: 8826460]

- Doody GA, Johnstone EC, Sanderson TL, Owens DG, Muir WJ. 'Pffropfschizophrenie' revisited. Schizophrenia in people with mild learning disability. *Br J Psychiatry* 1998;173:145–53. [PubMed: 9850227]
- Gedeon AK, Nelson J, Gecz J, Mulley JC. X-linked mild non-syndromic mental retardation with neuropsychiatric problems and the missense mutation A365E in PAK3. *Am J Med Genet A* 2003;120(4):509–17. [PubMed: 12884430]
- Hayashi ML, Choi SY, Rao BS, Jung HY, Lee HK, Zhang D, Chattarji S, Kirkwood A, Tonegawa S. Altered cortical synaptic morphology and impaired memory consolidation in forebrain- specific dominant-negative PAK transgenic mice. *Neuron* 2004;42(5):773–87. [PubMed: 15182717]
- Kolluri N, Sun Z, Sampson AR, Lewis DA. Lamina-specific reductions in dendritic spine density in the prefrontal cortex of subjects with schizophrenia. *Am J Psychiatry* 2005;162(6):1200–2. [PubMed: 15930070]
- Kreis P, Thevenot E, Rousseau V, Boda B, Muller D, Barnier JV. The p21-activated kinase 3 implicated in mental retardation regulates spine morphogenesis through a Cdc42-dependent pathway. *J Biol Chem* 2007;282(29):21497–506. [PubMed: 17537723]
- Lewis DA, Hashimoto T, Volk DW. Cortical inhibitory neurons and schizophrenia. *Nat Rev Neurosci* 2005;6(4):312–24. [PubMed: 15803162]
- Peippo M, Koivisto AM, Sarkamo T, Sipponen M, von Koskull H, Ylisaukkooja T, Rehnstrom K, Froyen G, Ignatius J, Jarvela I. PAK3 related mental disability: further characterization of the phenotype. *Am J Med Genet A* 2007;143(20):2406–16. [PubMed: 17853471]
- Reid AH. Psychiatric disorders in mentally handicapped children: a clinical and follow-up study. *J Ment Defic Res* 1980;24(Pt 4):287–98. [PubMed: 7218340]
- Rejeb I, Saillour Y, Castelnau L, Julien C, Bienvenu T, Taga P, Chaabouni H, Chelly J, Jemaa LB, Bahi-Buisson N. A novel splice mutation in PAK3 gene underlying mental retardation with neuropsychiatric features. *Eur J Hum Genet*. 2008
- Rosoklija G, Toomayan G, Ellis SP, Keilp J, Mann JJ, Latov N, Hays AP, Dwork AJ. Structural abnormalities of subicular dendrites in subjects with schizophrenia and mood disorders: preliminary findings. *Arch Gen Psychiatry* 2000;57(4):349–56. [PubMed: 10768696]
- St Clair D, Blackwood D, Muir W, Carothers A, Walker M, Spowart G, Gosden C, Evans HJ. Association within a family of a balanced autosomal translocation with major mental illness. *Lancet* 1990;336(8706):13–6. [PubMed: 1973210]
- Zammit S, Allebeck P, David AS, Dalman C, Hemmingsson T, Lundberg I, Lewis G. A longitudinal study of premorbid IQ Score and risk of developing schizophrenia, bipolar disorder, severe depression, and other nonaffective psychoses. *Arch Gen Psychiatry* 2004;61(4):354–60. [PubMed: 15066893]

Table 1

Exon	Amplicon (hg18 genomic assembly)	Sequence Variation
1	chrX:110,252,938-110,253,212	No variation
2	chrX:110,271,930-110,272,130	No variation
3	chrX:110,276,355-110,276,499	No variation
4	chrX:110,277,571-110,277,824	No variation
5	chrX:110,282,244-110,282,381	rs16986385 (18 males, 10 females - heterozygous), intronic SNP, 25 bp from SA site.
6	chrX:110,292,749-110,292,980	No variation
7	chrX:110,293,396-110,293,661	No variation
8	chrX:110,302,852-110,303,015	No variation
9	chrX:110,321,961-110,322,109	No variation
10	chrX:110,322,340-110,322,552	No variation
11	chrX:110,324,135-110,324,352	No variation
12	chrX:110,325,676-110,325,875	chrX:110,325,800, C->T, Leu395->Leu395 (1 female - heterozygous), novel SNP
13	chrX:110,326,278-110,326,574	No variation
14	chrX:110,346,255-110,346,492	rs10521535 (6 males, 6 females - heterozygous), intronic SNP, 40 bp from the SD site.
15	chrX:110,350,192-110,350,381	No variation