Published in final edited form as:

Hippocampus. 2009 February ; 19(2): 118–123. doi:10.1002/hipo.20492.

MR DETERMINED HIPPOCAMPAL ASYMMETRY IN FULL TERM AND PRETERM NEONATES

Deanne K. Thompson^{1,2}, Stephen J. Wood³, Lex W. Doyle⁴, Simon K. Warfield⁵, Gary F. Egan¹, and Terrie E. Inder²

¹ Neuroimaging and Neuroinformatics, Howard Florey Institute, and Centre for Neuroscience, University of Melbourne, Australia

² Department of Pediatrics, St Louis Children's Hospital, Washington University in St Louis, USA

³ Department of Psychiatry, Melbourne Neuropsychiatry Centre, University of Melbourne, Australia

⁴ Department of Obstetrics and Gynecology, Royal Women's Hospital, University of Melbourne, and Murdoch Childrens Research Institute, Melbourne, Australia

⁵ Department of Radiology, Children's Hospital, Harvard Medical School, Boston USA

Abstract

Hippocampi are asymmetrical in children and adults, where the right hippocampus is larger. To date, no literature has confirmed that hippocampal asymmetry is evident at birth. Furthermore, gender differences have been observed in normal hippocampal asymmetry, but this has not been examined in neonates. Stress, injury and lower IQ have been associated with alterations to hippocampal asymmetry. These same factors often accompany preterm birth. Therefore, prematurity is possibly associated with altered hippocampal asymmetry. There were three aims of this study: Firstly we assessed whether hippocampi were asymmetrical at birth, secondly whether there was a gender effect on hippocampal asymmetry, and thirdly whether the stress of preterm birth altered hippocampal asymmetry. This study utilized volumetric MRI to compare left and right hippocampal volumes in 32 full term and 184 preterm infants at term. Full term infants demonstrated rightward hippocampal asymmetry, as did preterm infants. In the case of preterm infants, hippocampal asymmetry was proportional to total hemispheric asymmetry. This study is the first to demonstrate that the normal pattern of hippocampal asymmetry is present this early in development. We did not find gender differences in hippocampal asymmetry at term. Preterm infants tended to have less asymmetrical hippocampi than full term infants, a difference which became significant after correcting for hemispheric brain tissue volumes. This study may suggest that hippocampal asymmetry develops in-utero and is maintained into adulthood in infants with a normal neurological course.

Keywords

infant, premature; magnetic resonance imaging; volumetry, hippocampus

INTRODUCTION

The main role of the hippocampus is in learning and memory (Burgess, 1999). It is also implicated in neuropsychiatric disorders (Geuze et al., 2005), and memory impairments such as Alzheimer's disease and amnesia (Jack et al., 2000). Hemispheric changes in the hippocampus may differentially affect function. Surgical damage to the left hippocampus reveals deficits in memory for verbal material (i.e., words), whereas damage to the right reveals deficits in non-verbal memory (i.e., faces or nonsense figures) (Zaidel, 1995).

Neurologically normal adults (Jack et al., 2000; Watson et al., 1992; Weis et al., 1989) and children (Giedd et al., 1996; Pfluger et al., 1999; Utsunomiya et al., 1999) have significantly larger right hippocampi. However, there is no literature confirming the presence or absence of hippocampal asymmetry at earlier stages in brain development. Hence it is uncertain how early in brain development asymmetry is acquired: in utero, during infancy or childhood. We therefore utilized quantitative MR Imaging technologies to determine whether hippocampal asymmetry was present in infants at term.

Gender differences have been associated with hippocampal asymmetry in older neurologically normal populations (Giedd et al., 1996; Pfluger et al., 1999). To determine if gender affects are evident at term, we compared hippocampal asymmetry between newborn males and females.

The hippocampus is vulnerable to stress (Perlman, 2001) and a wide variety of neurological insults including hypoxic-ischemic injury and metabolic insults (Schmidt-Kastner and Freund, 1991). Abnormal hippocampal asymmetry is associated with lower IQ (Abernethy et al., 2002) and temporal lobe epilepsy (Cook et al., 1992). The vulnerabilities and impairments associated with the hippocampus are reminiscent of the difficulties faced by preterm infants. Specifically, preterm infants often experience stressors such as hypoxic-ischemic injury and metabolic disorders (Fuller et al., 1983). They often have cognitive and learning deficits (Hack and Fanaroff, 1999), memory impairments (Briscoe et al., 1998), and increased risk of neuropsychological disorders such as attention deficit hyperactivity disorder (Warner-Rogers, 2002). These impairments may be related, at least in part, to hippocampal changes, and specifically hippocampal asymmetry. Our group has previously shown that the hippocampi of preterm infants are not disproportionately smaller than full term infants' relative to overall brain size (Thompson et al., 2008). However, uncorrected preterm hippocampal volumes were significantly smaller for the right hemisphere only (Thompson et al., 2008). Preterm birth is a stressful experience, and its effect on hippocampal asymmetry has never been specifically investigated. Therefore, we compared preterm and full term hippocampal asymmetry at term.

Two hundred and thirty three preterm infants (birth weight <1250 g or gestational age <30 weeks) from the Royal Women's Hospital in Melbourne and 51 healthy full term controls were recruited between July 2001 and December 2003. This study population has been previously described (Thompson et al., 2008).

All infants were scanned at term equivalent age (TEA: 38 - 42 weeks' postmenstrual age) without sedation. Images were acquired with a 1.5 Tesla General Electric Signa MRI scanner (Milwaukee, Wisconsin). T2 dual echo fast spin echo sequences were performed with interleaved acquisition (1.7 to 3.0 mm coronal slice thickness; in-plane resolution 0.35 to 0.43 mm²; Repetition Time 4000msec; Echo Time 60 / 160 msec; Field Of View 22 × 16 cm; matrix 256 × 192, interpolated 512 × 512). One hundred and eighty four preterm and 32 full term MRI scans (76% of those recruited) were able to be analyzed. The remaining scans were not performed within the TEA range (9%), or experienced movement or imaging artifacts (15%). Images were considered inappropriate for volumetric analysis if the brain

tissue contrast was too poor for tissues to be classified by automated segmentation (Warfield et al., 2000). The perinatal and demographic characteristics of the cohort are described in Table 1. Preterm infant gestational age ranged from 22 to 32 weeks, and birthweight ranged from 414 to 1395 g. There were no significant differences between the preterm and full term infants for either the gestational age at the time of MRI ($p=0.7$) or the male: female ratio ($p=0.6$) (Table 1).

Volumetric MRI analyses were undertaken on Sun Microsystems workstations (Palo Alto, California). The hippocampus was manually outlined as previously described (Thompson et al., 2008) by a single operator blinded to all perinatal data. The hippocampus was outlined on the coronal view of a combined raw T2-weighted and proton density weighted image volume with '3D slicer 2.5' software (<http://slicer.org/>). Combining these two images improved tissue contrast for optimized visualization of the hippocampal boundaries. Considering the neonatal brain is largely unmyelinated, T2 images display improved gray-white matter contrast over the T1 images routinely used in children and adults (Abernethy et al., 2004; Bernasconi et al., 2003). Generally, anatomical boundaries followed the approach of Watson et al. (Watson et al., 1992). Firstly the posterior boundary at the hippocampal tail was defined as the most anterior slice where the fornix was seen in its entirety. The inferior boundary of the hippocampal tail consisted of the white matter of the parahippocampal gyrus (excluded). The fimbria, subiculum and alveus were included in the measurement. In-plane boundaries were traced sequentially on each slice. The uncus was included once it appeared fused to the hippocampal body. The anterior boundary at the head of the hippocampus was defined as the most anterior slice where cerebrospinal fluid was still seen temporally to the hippocampus. The head of the hippocampus was distinguished from the overlying amygdala by following the path of the top of the temporal horn of the lateral ventricles, where a line of intermingling CSF and white matter was often visible. This resulted in a 3D hippocampal volume for both the right and left hippocampi (Figure 1). The average (SD) number of slices from which the full term hippocampi were obtained was 10.4 (2.0) for a total hippocampal length (measured between the first and last slice) of 21.4 (3.0) mm. Intraclass correlation coefficients for intra-observer reliability were 0.97 for the right and 0.96 for the left hippocampi.

Each image was segmented twice, once in original orientation and again after being flipped in the left/right direction. This minimized possible hemispheric bias by the operator. Tracing mirrored and non-mirrored scans were alternated and always traced at least one day apart. There were no significant differences between the hippocampal volumes traced in the original space and those that had been flipped, for either the right [Original: 1.14 (0.17) cm^3 ; Flipped: 1.14 (0.17) cm^3 , $p=0.8$], or the left hippocampi [Original: 1.11 (0.18) cm^3 ; Flipped: 1.13 (0.18) cm^3 , $p=0.2$]. These results suggest there was no hemispheric bias in the operator. The two volumes (original and flipped) were averaged for each hemisphere.

Statistical analyses were performed using SPSS version 15.0. In order to assess hippocampal asymmetry, the left and right hippocampal volumes were compared by paired samples t -tests. The paired variables were the left and right hippocampal volume. The entire cohort was analysed as well as preterm and full term infants separately. Differences in asymmetry based on gender were examined by repeated measures ANOVA, with hemisphere as the within-subjects variable, and gender as the between-subject factor. Differences in asymmetry based on group were also examined by repeated measures ANOVA, again with hemisphere as the within subject variable, but group as the between subject factor. Correction for total head size (intracranial volume) was performed as previously described (Thompson et al., 2008). Results did not differ, and therefore these analyses are not included. The same method was used to correct hippocampal volumes for the total brain tissue of each hemisphere, including cortical and deep nuclear gray matter, myelinated and

unmyelinated white matter (excluding cerebrospinal fluid). The hemispheres were divided along the mid-sagittal plane after aligning the brain image along the anterior and posterior commissure line (Thompson et al., 2007). Correcting for each hemisphere ensured that hippocampal asymmetry was not simply a reflection of whole brain laterality differences.

Hippocampal asymmetry at term: Newborn infant's hippocampi were asymmetrical at term. The right (R) was 0.019 cm³ larger than the left (L) hemisphere [R 1.14 (0.156) cm³, L 1.12 (0.160) cm³; 95% Confidence Interval (CI) 0.014, 0.025; $p < 0.001$]. This was significant for both the full term infants, with 0.031 cm³ larger right hippocampi [R 1.19 (0.121) cm³, L 1.16 (0.132) cm³; 95% CI 0.013, 0.049; $p = 0.001$], and the preterm infants with 0.017 cm³ larger right hippocampi [R 1.13 (0.160) cm³, L 1.12 (0.164) cm³; 95% CI 0.012, 0.023; $p < 0.001$] (Figure 2a). Hippocampal asymmetry was significant only for the full term infants after correcting for total brain tissue of the hemisphere. The difference between left and right hippocampi was no longer significant for preterm infants after correcting for total hemispheric asymmetry [All infants: R 1.17 (0.127), L 1.16 (0.128), $p = 0.2$; FT: R 1.19 (0.104), L 1.16 (0.112), $p = 0.01$, PT: R 1.16 (0.130), L 1.16 (0.131), $p = 0.9$].

Gender effect on hippocampal asymmetry: Gender did not substantially affect hippocampal asymmetry in neonates ($F_{1, 214} = 0.32$, $p = 0.6$), either full term ($F_{1, 30} = 0.10$, $p = 0.8$) or preterm ($F_{1, 182} = 0.77$, $p = 0.4$), even after correction for total hemispheric tissue (All infants: $p = 0.2$, FT: $p = 0.1$, PT: $p = 0.6$).

Effect of preterm birth on hippocampal asymmetry: Preterm infants tended to have reduced hippocampal asymmetry when compared with full term infants, although this did not reach statistical significance ($F_{1, 214} = 2.7$, $p = 0.09$). However after correction for total brain tissue within the hemisphere, the difference in asymmetry between preterm and full term infants was significant ($F_{1, 214} = 7.4$, $p = 0.007$) (Figure 2b).

Hippocampal asymmetry is present at term. Furthermore, at term males and females do not differ in their pattern of asymmetry. Preterm hippocampi tend to be less asymmetrical than full term infants'.

The maximum growth rate of the hippocampus occurs between the first and second month after birth (Kretschmann et al., 1986; Utsunomiya et al., 1999), and continues up to 2 years (Utsunomiya et al., 1999). Although the hippocampus is still at an early stage of development, this study confirmed that hippocampal asymmetry is present at birth. The right hippocampus was significantly larger than the left for both full term and preterm infants. In contrast, during adolescence, left and right hippocampi do not appear to be different in either preterm or full term subjects (Isaacs et al., 2000). The current study extends previous volumetric findings that have shown that the cortical and deep nuclear gray matter, as well as total brain tissue volume (excluding cerebrospinal fluid) demonstrate 'rightward' asymmetry in neonates (Thompson et al., 2007). When taking into account the asymmetry of the total brain, preterm right and left hippocampi were no longer different, suggesting hippocampal asymmetry in the preterm infants is proportional to total brain asymmetry. In contrast, full term infant hippocampi were more asymmetrical than the rest of the brain. This suggests that under normal circumstances, rightward hippocampal asymmetry is greater than total brain asymmetry. This disproportional growth of the right hippocampus may prime the infant for greater visuospatial (right brain) as opposed to linguistic ability (left brain).

The underlying mechanisms for the right-larger-than-left asymmetry in the hippocampus are largely unknown. Originally it was thought that asymmetry was functionally based (Geschwind and Levitsky, 1968). However the fact that asymmetry is present at birth may suggest otherwise. Twin studies suggest moderate heritability of hippocampal volumes in healthy subjects (Peper et al., 2007). However, hippocampal volume loss in psychiatric

conditions appears to be influenced by environmental factors (de Geus et al., 2007; van Erp et al., 2004) such as pregnancy and birth complications (Stefanis et al., 1999) or stress (Sapolsky, 2000). We have previously shown that white matter injury, postnatal steroid exposure and indomethacin treatment negatively influence hippocampal growth in preterm infants (Thompson et al., 2008). Environmental influences may affect the hippocampus asymmetrically (Stefanis et al., 1999; Vythilingam et al., 2002).

This study could not detect a gender difference in hippocampal asymmetry for either the full term or preterm infants, contrary to the findings of Pfluger et al., who determined that male children had greater asymmetry (Pfluger et al., 1999). Giedd et al., suggested that sex-specific maturational changes in the hippocampus of healthy children were potentially due to hormonal responsivity (Giedd et al., 1996). Other studies have been unable to find a gender effect on hippocampal asymmetry (Jack et al., 1989; Szabo et al., 2001). Gender differences may become apparent later in development. Therefore hormonal influences on hippocampal development are worthy of further investigation, particularly longitudinally through adolescence.

Raw hippocampal volumes showed a tendency to be more symmetrical in preterm infants compared to full term infants, but this was not statistically significant. Similarly, a previous study examining adolescents born preterm did not find asymmetrical differences between preterm and full term hippocampi (Isaacs et al., 2000). One interpretation of these results is that preterm hippocampal asymmetry develops similarly to that of the full term infant, and is therefore 'normal'. It should be noted, however, that our relatively small sample of full term infants may have contributed to this non-significant finding in both the current study and our previous study (Thompson et al., 2008). Furthermore, correction for total brain volume within each hemisphere revealed that preterm hippocampal asymmetry *was* significantly smaller than for full-terms. Therefore, an alternative interpretation is that preterm hippocampal asymmetry is *not* developing normally. Overall, these results suggest that hippocampal asymmetry normally develops in-utero, but in preterm infants born at least 10 weeks early, this disproportionate growth of the right hippocampus may not proceed normally.

The hippocampi were manually segmented based on solid neuroanatomical landmarks, ensuring a representative and reproducible measurement. The same landmarks were used in each hemisphere, and hippocampal segmentation was performed in both original and flipped orientation. This should have precluded hemispheric bias of the segmentation method or operator. Hippocampal volumetry is subject to artifacts arising from MRI acquisition and post-processing. Nonetheless, the high intra-observer reliability estimates (>0.95) indicate that acquisition and processing were adequate to obtain reliable hippocampal volume measurements. The in-plane resolution was high for these scans. However, 46% of infant images (86 preterm and 13 full term) were acquired with large slice thicknesses (3.0 mm) relative to their hippocampal size. This may generate partial volume error, particularly at the borders of the hippocampal tail and head. Cook and colleagues suggested that slices of 3 mm or thinner allow accuracy within 5% for adult hippocampal volumes (Cook et al., 1992). Presumably this level of accuracy would require thinner slices for smaller neonatal hippocampi. We recognise that it would have been desirable to obtain more slices of reduced thickness. However, this would require an increased imaging time, which was impractical for infants of this age.

This study is the first to demonstrate that healthy newborn infants have 'rightward' hippocampal asymmetry. This normal pattern of asymmetry is similar for both male and female full term infants. There is evidence to suggest that preterm infant hippocampal asymmetry is altered. The implications of these findings in regards to future neurological

health of these infants require further investigation. Consequently, neurodevelopmental follow up of this cohort is planned.

Acknowledgments

We gratefully thank Marilyn Bear, Tamara-Leigh E Brawn, Gregory A Lodygensky, Hong X Wang, Michael J Farrell, Peter J Anderson and Rodney W Hunt, as well as the families and infants who participated in this study.

Grant sponsors: This study was supported by the National Medical and Health Research Council of Australia (Grant number: 237117) and the United Cerebral Palsy Foundation (USA). This study was supported in part by the NIH (Grant numbers: R01 RR021885, R01 GM074068, R01 EB008015 and P30 HD018655), and by the Mather Foundation (USA) and the Brown Foundation (USA). SJW is supported by an NHMRC Clinical Career Development Award and a NARSAD Young Investigator Award. GFE is supported by a NHMRC Research Fellowship (Grant number: 400317).

REFERENCES

- Abernethy LJ, Cooke RWI, Foulder-Hughes L. Caudate and hippocampal volumes, intelligence, and motor impairment in 7-year-old children who were born preterm. *Pediatr Res*. 2004; 55(5):884–93. [PubMed: 14764914]
- Abernethy LJ, Palaniappan M, Cooke RW. Quantitative magnetic resonance imaging of the brain in survivors of very low birth weight. *Archives of Disease in Childhood*. 2002; 87(4):279–83. [PubMed: 12243993]
- Bernasconi N, Bernasconi A, Caramanos Z, Antel SB, Andermann F, Arnold DL. Mesial temporal damage in temporal lobe epilepsy: a volumetric MRI study of the hippocampus, amygdala and parahippocampal region. *Brain*. 2003; 126(Pt 2):462–9. [PubMed: 12538412]
- Briscoe J, Gathercole SE, Marlow N. Short-term memory and language outcomes after extreme prematurity at birth. *Journal of Speech Language and Hearing Research*. 1998; 41(3):654–666.
- Burgess, N.; Jeffery, KJ.; O'Keefe, J. Integrating hippocampal and parietal functions: a spatial point of view. In: Burgess, N.; Jeffery, KJ.; O'Keefe, J., editors. *The hippocampal and parietal foundations of spatial cognition*. Oxford University Press, Inc.; New York: 1999. p. 1-29.
- Cook MJ, Fish DR, Shorvon SD, Straughan K, Stevens JM. Hippocampal volumetric and morphometric studies in frontal and temporal lobe epilepsy. *Brain*. 1992; 115(Pt 4):1001–15. [PubMed: 1393499]
- de Geus EJC, van't Ent D, Wolfensberger SPA, Heutink P, Hoogendijk WJG, Boomsma DI, Veltman DJ. Intrapair differences in hippocampal volume in monozygotic twins discordant for the risk for anxiety and depression. *Biol Psychiatry*. 2007; 61(9):1062–71. [PubMed: 17137562]
- Fuller PW, Guthrie RD, Alvord EC Jr. A proposed neuropathological basis for learning disabilities in children born prematurely. *Developmental Medicine & Child Neurology*. 1983; 25(2):214–31. [PubMed: 6852386]
- Geschwind N, Levitsky W. Human brain: left-right asymmetries in temporal speech region. *Science*. 1968; 161(837):186–7. [PubMed: 5657070]
- Geuze E, Vermetten E, Bremner JD. MR-based in vivo hippocampal volumetrics: 2. Findings in neuropsychiatric disorders. *Molecular Psychiatry*. 2005; 10(2):160–184. [PubMed: 15356639]
- Giedd JN, Vaituzis AC, Hamburger SD, Lange N, Rajapakse JC, Kaysen D, Vauss YC, Rapoport JL. Quantitative MRI of the temporal lobe, amygdala, and hippocampus in normal human development: ages 4–18 years. *J Comp Neurol*. 1996; 366(2):223–30. [PubMed: 8698883]
- Hack M, Fanaroff AA. Outcomes of children of extremely low birthweight and gestational age in the 1990's. *Early Hum Dev*. 1999; 53(3):193–218. [PubMed: 10088988]
- Isaacs EB, Lucas A, Chong WK, Wood SJ, Johnson CL, Marshall C, Vargha-Khadem F, Gadian DG. Hippocampal volume and everyday memory in children of very low birth weight. *Pediatric Research*. 2000; 47(6):713–20. [PubMed: 10832727]
- Jack CR Jr, Petersen RC, Xu Y, O'Brien PC, Smith GE, Ivnik RJ, Boeve BF, Tangalos EG, Kokmen E. Rates of hippocampal atrophy correlate with change in clinical status in aging and AD. *Neurology*. 2000; 55(4):484–89. [PubMed: 10953178]

- Jack CR Jr, Twomey CK, Zinsmeister AR, Sharbrough FW, Petersen RC, Cascino GD. Anterior temporal lobes and hippocampal formations: normative volumetric measurements from MR images in young adults. *Radiology*. 1989; 172(2):549–54. [PubMed: 2748838]
- Kretschmann HJ, Kammradt G, Krauthausen I, Sauer B, Wingert F. Growth of the hippocampal formation in man. *Bibl Anat*. 1986; (28):27–52. [PubMed: 3707511]
- Peper JS, Brouwer RM, Boomsma DI, Kahn RS, Hulshoff Pol HE. Genetic influences on human brain structure: a review of brain imaging studies in twins. *Hum Brain Mapp*. 2007; 28(6):464–73. [PubMed: 17415783]
- Perlman JM. Neurobehavioral deficits in premature graduates of intensive care - Potential medical and neonatal environmental risk factors. *Pediatrics*. 2001; 108(6):1339–1348. [PubMed: 11731657]
- Pfluger T, Weil S, Weis S, Vollmar C, Heiss D, Egger J, Scheck R, Hahn K. Normative volumetric data of the developing hippocampus in children based on magnetic resonance imaging. *Epilepsia*. 1999; 40(4):414–23. [PubMed: 10219266]
- Sapolsky RM. Glucocorticoids and hippocampal atrophy in neuropsychiatric disorders. *Arch Gen Psychiatry*. 2000; 57(10):925–35. [PubMed: 11015810]
- Schmidt-Kastner R, Freund TF. Selective vulnerability of the hippocampus in brain ischemia. *Neuroscience*. 1991; 40(3):599–636. [PubMed: 1676492]
- Stefanis N, Frangou S, Yakeley J, Sharma T, O'Connell P, Morgan K, Sigmudsson T, Taylor M, Murray R. Hippocampal volume reduction in schizophrenia: effects of genetic risk and pregnancy and birth complications. *Biol Psychiatry*. 1999; 46(5):697–702. [PubMed: 10472422]
- Szabo CA, Xiong J, Lancaster JL, Rainey L, Fox P. Amygdalar and hippocampal volumetry in control participants: differences regarding handedness. *Ajnr: American Journal of Neuroradiology*. 2001; 22(7):1342–5. [PubMed: 11498423]
- Thompson DK, Warfield SK, Carlin JB, Pavlovic M, Wang HX, Bear M, Kean MJ, Doyle LW, Egan GF, Inder TE. Perinatal risk factors altering regional brain structure in the preterm infant. *Brain*. 2007; 130(Pt 3):667–77. [PubMed: 17008333]
- Thompson DK, Wood SJ, Doyle LW, Warfield SK, Lodygensky GA, Anderson PJ, Egan GF, Inder TE. Neonate hippocampal volumes: Prematurity, perinatal predictors, and 2-year outcome. *Ann Neurol*. 2008; 63(5):642–651. [PubMed: 18384167]
- Utsunomiya H, Takano K, Okazaki M, Mitsudome A. Development of the temporal lobe in infants and children: Analysis by MR-based volumetry. *American Journal of Neuroradiology*. 1999; 20(4):717–723. [PubMed: 10319988]
- van Erp TGM, Saleh PA, Huttunen M, Lonnqvist J, Kaprio J, Salonen O, Valanne L, Poutanen V-P, Standertskjold-Nordenstam C-G, Cannon TD. Hippocampal volumes in schizophrenic twins. *Arch Gen Psychiatry*. 2004; 61(4):346–53. [PubMed: 15066892]
- Vythilingam M, Heim C, Newport J, Miller AH, Anderson E, Bronen R, Brummer M, Staib L, Vermetten E, Charney DS. Childhood trauma associated with smaller hippocampal volume in women with major depression. *American Journal of Psychiatry*. 2002; 159(12):2072–80. others. [PubMed: 12450959]
- Warfield SK, Kaus M, Jolesz FA, Kilinis R. Adaptive, Template Moderated, Spatially Varying Statistical Classification. *Medical Image Analysis*. 2000; 4(1):43–55. [PubMed: 10972320]
- Warner-Rogers, J. Attention deficit hyperactivity disorder. In: Howlin, P.; Udwin, O., editors. *Outcomes in neurodevelopmental and genetic disorders*. Cambridge University Press; Cambridge: 2002.
- Watson C, Andermann F, Gloor P, Jonesgotman M, Peters T, Evans A, Olivier A, Melanson D, Leroux G. Anatomic Basis of Amygdaloid and Hippocampal Volume Measurement by Magnetic-Resonance-Imaging. *Neurology*. 1992; 42(9):1743–1750. [PubMed: 1513464]
- Weis S, Haug H, Holoubek B, Orun H. The cerebral dominances: quantitative morphology of the human cerebral cortex. *Int J Neurosci*. 1989; 47(1-2):165–8. [PubMed: 2793342]
- Zaidel DW. The case for a relationship between human memory, hippocampus and corpus callosum. *Biol Res*. 1995; 28(1):51–7. [PubMed: 8728820]

FIGURE 1.

(a) Segmentation boundaries of four coronal sections through the hippocampal tail, body and head. (b) The hippocampal segmentation boundaries shown in all three orthogonal views: axial, sagittal and coronal, as well as a three dimensional representation of the left (blue) and right (pink) hippocampal volumes.

FIGURE 2.

(a) Right and left hippocampal volumes for preterm and full term infants. Box plots display the median, 25th and 75th centiles, range of the data and outliers. (b) Estimated differences in mean hippocampal volumes (corrected for asymmetry in total brain tissue) between right and left hemispheres (Right minus Left, 95% CI), for both preterm and full term infants.

TABLE 1

Perinatal and demographic characteristics of the cohorts.

Characteristic	Full term n=32	Preterm n=184
Gestational age at birth (wk) – mean (SD)	39.0 (1.2)	27.6 (1.9)
Gestational age at MRI – mean (SD)	40.5 (1.0)	40.1 (1.1)
Birthweight, g – mean (SD)	3289 (504)	964 (219)
Weight at MRI, g – mean (SD)	3484 (490)	2971 (531)
Male – n (%)	18 (56)	93 (51)
Multiple births – n (%)	1 (3)	79 (43)
Bronchopulmonary dysplasia ¹ – n (%)	0 (0)	68 (35)
Inotropic Support – n (%)	0 (0)	71 (39)
Postnatal corticosteroids ² – n (%)	0 (0)	14 (8)
Intrauterine growth restriction ³ – n (%)	2 (6)	22 (12)
Necrotizing enterocolitis – n (%)	0 (0)	19 (10)
Proven sepsis – n (%)	1 (3)	80 (44)
Antenatal corticosteroids – n (%)	0 (0)	164 (89)
Positive pressure ventilation, hr – mean (SD)	0.59 (3.4)	223 (365)
Parenteral nutrition, days – mean (SD)	0.19 (1.1)	14 (17)
Patent ductus arteriosus (indomethacin administered), n (%)	0 (0)	63 (34)
WM injury (any grade), n (%)	1 (3)	118 (64)
WM injury (grade III/IV), n (%)	0 (0)	31 (17)
Intraventricular Hemorrhage, n (%)	0 (0)	24 (13)

¹ Required Oxygen at 36 weeks gestational age.² Postnatal dexamethasone, 0.15 mg/kg per day, reducing over 10 days.³ Z score > 2 SD below mean weight for gestational age.