Published in final edited form as:

Tetrahedron Lett. 2008 November 3; 49(45): 6383–6385. doi:10.1016/j.tetlet.2008.07.184.

Biomimetic total synthesis of tricycloillicinone and mechanistic studies toward the rearrangement of prenyl phenyl ethers

Xiaoguang Lei^a, Mingji Dai^a, Zihao Hua^b, and Samuel J. Danishefsky^{a,b,*}

^aDepartment of Chemistry, Columbia University, 3000 Broadway, New York, NY 10027, USA

^bLaboratory for Bioorganic Chemistry, Sloan-Kettering Institute for Cancer Research, 1275 York Avenue, New York, NY 10065, USA

Abstract

The paper describes a short and biomimetic synthesis of tricycloillicinone, which was found to enhance the action of choline acetyltransferase (ChAT). The synthetic route has two critical reactions: bulky, oxygenophilic methylaluminum bis(4-bromo-2,6-di-*tert*-butylphenoxide) (MABR) promoted rearrangement of prenyl phenyl ether and photochemical cyclization. Furthermore, experiments were designed to explore the process of MABR promoted rearrangement. It was found that stereochemistry of deuterium labeled prenyl group was only partially scrambled, which suggests that there may be two possible reaction pathways involved in this process. It also suggests that the direct migration of prenyl group to *para*-position under these conditions is slightly favored over the Claisen-Cope process. The highly efficient synthetic route also provides important new opportunities to explore the biological behavior of tricycloillicinone.

Tricycloillicinone (**1**, Scheme 1) is a member of an intriguing class of small molecule natural products that have been demonstrated to possess neurotrophic activity. Isolated from *Illicium tashiroi* by Fukuyama and coworkers in 1995,¹ tricycloillicinone was found to enhance the activity of choline acetyltransferase (ChAT), an enzyme required for the biosynthesis of the important neurotransmitter, acetylcholine.² Deficiencies in acetylcholine levels have been shown to be associated with the progression of neurodegenerative disorders, such as Alzheimer's disease.³ It is therefore conceivable that a ChAT inducer, such as tricycloillicinone, could serve as a valuable lead compound in the development of novel therapeutic agents for the treatment of neurodegenerative diseases.⁴

In 1998, our laboratory disclosed the first total synthesis of **1**.⁵ More recently, Terashima and coworkers reported the enantioselective total synthesis of tricycloillicinone.⁶ Unfortunately, both of these synthetic routes are quite long, and the difficulties involved in gaining access to sufficient quantities of synthetic material have thus far limited our ability to perform all of the needed biological evaluations. We report herein the development of a rather more efficient, biomimetically inspired second generation total synthesis of tricycloillicinone. A key feature of this total synthesis involves rearrangement of a prenyl phenyl ether (see **5** → **2**). In addition to exploiting this reaction for our total synthesis, we conducted experiments hoping to gain a

© 2008 Elsevier Ltd. All rights reserved.

*Corresponding author. Tel.: +1-212-639-5501; fax: +1-212-772-8691; e-mail: danishes@mskcc.org.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

deeper understanding of the migration process. Though the results did not provide a clear-cut definitive answer to our queries, they are related herein for their intrinsic interest.

It has been shown that illicinone A (**2**) is a potential biosynthetic precursor of tricycloillicinone. Furukawa and co-workers further demonstrated that illicinole (**5**) could be converted to illicinone A (**2**) and thence to tricycloillicinone (**1**) by photo-irradiation, although the reaction yield is very low and several side products (**3**, **4**, and **6**) are also generated (Scheme 1).⁷

In order to achieve a more efficient biomimetic total synthesis of tricycloillicinone, we would first need to develop a concise route to the key precursor, illicinone **2**. Accordingly, our route to **2** began with alkylation of commercially available sesamol **7**, thus affording allyl ether **8**. Claisen rearrangement of **8** under thermal conditions smoothly provided **9** in 91% yield. Prenylation of phenol **9** afforded illicinole **5**. Initial attempts to generate illicinone **2** under thermal conditions were unsuccessful. We therefore decided to investigate a Lewis acid-catalyzed rearrangement of **5**. Upon screening a number of different catalysts, we were pleased to find that the bulky Lewis acid MABR (**10**), developed by Yamamoto and coworkers,⁸ showed the most promising results. As shown, treatment of illicinole **5** with MABR at -78°C for 2 hours cleanly afforded illicinone **2** in 70 % yield. With intermediate **2** in hand, we were able to investigate the key photochemical reaction. Upon optimization of the photochemistry conditions, we were able to obtain tricycloillicinone **1** in 35% yield, along with tetracycloillicinone **3** in 26% yield.

We next hoped to answer some questions regarding the mechanism of the key MABR-mediated rearrangement. As outlined in Figure 1, two possible reaction pathways can readily be envisioned: in the first (path a), substrate **5** would undergo Claisen rearrangement (to generate **6**), followed by Cope rearrangement, to afford product **2**. Alternatively, the transformation might involve direct prenyl group migration to the *para* position to generate illicinone **2** (path b).

We hoped that the mechanistic nature of this key rearrangement step could be probed through a well-designed deuterium isotope study. Along these lines, we prepared the deuterium-labeled illicinole **12**, as shown in Scheme 3. Thus, treatment of phenol **9** with alcohol **14**⁹ under Mitsunobu conditions afforded vinyl iodide **13** in 86% yield. Cuprate coupling¹⁰ of vinyl iodide **13** using CD_3Li cleanly provided the desired deuterium labeled illicinole **12**. Importantly, compound **12** has a well-defined *cis*-olefin geometry. Our thinking was that, if rearrangement path a were to predominate, then the rearranged product would be produced as a 1:1 mixture of *cis:trans* isomers. By contrast, if path b were operational, the product could well retain the *cis*-geometry of its rearrangement precursor.

In the event, following exposure to MABR, as shown, illicinole isotope **12** rearranged to afford adduct **15** as a 3:1 mixture of *cis* and *trans* isomers (Figure 2). This result suggests that, in fact, both reaction pathways may be involved; however, the direct prenyl migration is apparently favored. Yamamoto and coworkers have proposed a transition state for this Lewis acid mediated rearrangement process.^{8b} As shown in Figure 2, the experimental results imply that the initial coordination of bulky aluminum catalyst (MABR) to prenyl ether **12** generates the sterically more favored complex **A** rather than the alternate complex **B**. This presumably favored conformation of the complex would indeed tend to rationalize the observed preference for the direct prenyl migration pathway. Unfortunately, our data do not definitively speak to the question as to whether there may be some erosion in the stereointegrity of the prenyl double bond, even in a direct prenyl migration pathway. If some scrambling occurred in such a "direct" walkover, there would be no need to invoke any level of participation from the two-stage Claisen-Cope scenario (see compound **6**) to account for the loss of geometric "memory". Of course, avoidance of formation of an intermediate of the type **6** would explain the absence of

competing allyl migration. However, it is not at all unlikely that the migration aptitude of the dimethylallyl group in the hypothetical **6** would be far greater than the allyl group in the migration of the allyl. Hence, the question of the intermediacy of **6** as a competing process is not fully answered.

In summary, an efficient biomimetic total synthesis of the neurotrophic agent tricycloillicinone has been accomplished from commercially available sesamol in only 6 steps. The synthesis relies on a key MABR-promoted rearrangement of prenyl phenyl ether and a subsequent photochemical cyclization sequence. Preliminary mechanistic studies toward the rearrangement process suggest that two different reaction pathways (Claisen–Cope and direct prenyl migration) may be operational in the key rearrangement step. With the development of this concise and scalable route, we are now able to readily access large quantities of synthetic tricycloillicinone, and its analogs, for further biological and SAR studies.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We are grateful to the National Institutes of Health (HL25848 to SJD) for financial support of this research. We thank Ms. Rebecca Wilson for valuable help in the editing of this manuscript.

References

1. Fukuyama Y, Shida N, Kodama M, Chaki H, Yugami T. Chem. Pharm. Bull 1995;43:2270. [PubMed: 8582027]
2. For a review, see: Rosenberg, S.; Section, I. Annual Reports in Medicinal Chemistry-27. New York: Academic Press; 1992. CNS Reagents; p. 41
3. Seiger A, Nordberg A, vonHolst H, Backman L, Ebendal L, Alafuzoff I, Ambria K, Herlitz A, Lilja A. Behav. Brain Res 1993;57:255. [PubMed: 8117429]
4. Hefti F. J. Neurobiol 1994;25:1418. [PubMed: 7852995]
5. (a) Pettus TRR, Chen X-T, Danishefsky SJ. J. Am. Chem. Soc 1998;120:12684. (b) Pettus TRR, Inoue M, Chen X-T, Danishefsky SJ. J. Am. Chem. Soc 2000;122:6160.
6. Furuya S, Terashima S. Tetrahedron Lett 2003;44:6875–6878.
7. Yakushijin K, Fukukawa H, McPhail AT. Chem. Pharm. Bull 1984;32:23.
8. (a) Nonoshita K, Banno H, Maruoka K, Yamamoto H. J. Am. Chem. Soc 1990;112:316. (b) Maruoka K, Sato J, Banno H, Yamamoto H. Tetrahedron Lett 1990;31:377.
9. Cowell A, Stille JK. J. Am. Chem. Soc 1980;102:4193.
10. Salinaro RF, Berson JA. J. Am. Chem. Soc 1982;104:2228.

Figure 1.
Possible reaction pathways for the MABR-mediated rearrangement

Figure 2.
MABR-mediated rearrangement.

Scheme 1.
Photochemical Reactions of the Illicinone Series.

Scheme 2.
Efficient synthesis of tricycloillicinone (1)

Scheme 3.
Synthesis of deuterium labeled illicinole **12**.