Published in final edited form as:

J Med Chem. 2009 January 22; 52(2): 456–468. doi:10.1021/jm801128g.

Nonpeptide Macrocyclic Histone Deacetylase Inhibitors

Adegboyega K. Oyelere^{*,¶,§}, Po C. Chen[¶], William Guerrant[¶], Sandra C. Mwakwari[¶], Rebecca Hood^{#,¶}, Yunzhe Zhang[±], and Yuhong Fan^{±,§}

[¶]*School of Chemistry and Biochemistry, Georgia Institute of Technology, Atlanta, GA 30332-0400 USA*

[±]*School of Biology, Georgia Institute of Technology, Atlanta, GA 30332-0400 USA*

[§]*Parker H. Petit Institute for Bioengineering and Bioscience, Georgia Institute of Technology, Atlanta, GA 30332-0400 USA*

Abstract

Inhibition of Histone Deacetylases inhibitors (HDACi) hold great promise in cancer therapy due to their demonstrated ability to arrest proliferation of nearly all transformed cell types. Of the several structurally distinct small molecules HDACi reported, macrocyclic depsipeptides have the most complex recognition cap-group moieties and present an excellent opportunity for the modulation of the biological activities of HDACi. Unfortunately, the structure–activity relationship (SAR) studies for this class of compounds have been impaired largely because most macrocyclic HDACi known to date are comprised of complex peptide macrocycles. In addition to retaining the pharmacologically disadvantaged peptidyl-backbone, they offer only limited opportunity for side-chain modifications. Here we report the discovery of a new class of macrocyclic HDACi based on the macrolide antibiotics skeletons. SAR studies revealed that these compounds displayed both linker-length and macrolide-type dependent HDAC inhibition activities with IC₅₀ in low nanomolar range. In addition, these nonpeptide macrocyclic HDACi are more selective against HDAC 1 and 2 relative to HDAC 8, another class I HDAC isoform, hence have sub-class HDAC isoform selectivity.

Introduction

Inhibition of Histone Deacetylases (HDACs) has recently been clinically validated as a novel therapeutic strategy for cancer treatment.¹ Because of their demonstrated ability to arrest proliferation of nearly all transformed cell types,² HDAC inhibitors (HDACi) hold great promise as agents of choice, either as stand alone therapeutics or in combination with others, in the fight against the cancer scourge. To date, several structurally distinct small molecule HDACi have been reported including aryl hydroxamates, benzamides, short-chain fatty acids, electrophilic ketones and macrocyclic-peptides (Scheme 1).^{3–6} All HDACi so far reported fit a three-motif pharmacophoric model namely, a zinc-binding group (ZBG), a hydrophobic linker and a recognition cap-group.³ The X-ray crystal structures of a bacterial HDAC homolog, histone deacetylase-like protein (HDLP) bound to suberoylanilide hydroxamic acid (SAHA) and trichostatin A (TSA), and recently human HDAC8 and HDAC7, have validated this model.^{7, 8} Of these HDACi, macrocyclic-peptides have the most complex recognition cap-group moieties and present an excellent opportunity for the modulation of the biological activities of HDACi. Although cyclic-peptide HDACi possess potent HDAC inhibition activity (nanomolar range), their broad application in cancer therapy currently remains largely unproven.³ One promising exception, **FK-228** (Scheme 1), is currently in phase II study for the treatment of cutaneous T-Cell lymphoma.⁹

* To whom the correspondence should be addressed. Phone: 404-894-4047; fax: 404-894-2291; e-mail: aoyelere@gatech.edu.

#Beckman Undergraduate Research Fellow

The dearth of clinically effective cyclic-peptide HDACi may be in part due to development problems characteristic of large peptides, most especially poor oral bioavailability. In addition to retaining the pharmacologically disadvantaged peptidyl-backbone, they offer only limited opportunity for side-chain modifications.¹⁰ Identification of non-peptide macrocyclic HDACi will offer a new class of macrocyclic HDACi with potentially more favorable drug-like properties. Furthermore, this will aid comprehensive SAR studies and further enhance our understanding of the roles of specific interactions between the enzyme outer rim and inhibitor cap-groups in HDACi activity and selectivity. Herein we report the discovery of a new class of potent, non-peptide macrocyclic HDACi derived from the macrolide macrocyclic ring structures.

Results and Discussion

Macrolides are glycosylated polyketide antibiotics that have been in use for over 50 years for the treatment of respiratory tract infections. Additionally, macrolides have elicited other non-antibiotic effects, including anti-inflammatory and immunomodulatory effects that make them promising candidates for the management of diseases of chronic airway inflammation.^{11, 12} More recently, macrolides derived from the 6-*O*-methylerythromycin A ring have been reported to serve as nonpeptidic surrogates for the peptide backbone of macrocyclic peptide luteinizing hormone-releasing hormone (LHRH) receptor antagonists.¹³ Drawing inspiration from the peptidomimetic property of macrolides, we hypothesize that an appropriate substitution of the cyclic peptide moiety of a prototypical cyclic-peptide HDACi with macrolide skeletons will generate a new class of potent non-peptide macrocyclic HDACi.

To test our hypothesis, we first sought a SAHA-macrolide conjugate which incorporated the 15-membered azalide ring of azithromycin as the macrolide template (Scheme 1). Azithromycin skeleton is an attractive choice due to its excellent pharmacokinetic (PK) profile and ease of chemical transformation of key moieties on the skeleton.¹⁴ Our design approach is to attach the HDAC inhibiting group to a macrolide moiety that is remote from the macrocyclic ring. We anticipate that this will minimize the potential steric clash at the HDAC active site that might result from the introduction of the azithromycin skeleton, a macrocyclic ring not optimized to bind the HDAC enzyme. NMR structure¹⁵ and 3D model revealed that the 3'-tertiary amine of the desosamine sugar and the 4''-OH of the cladinose sugar met our design requirement. We opted for the former group because of the well established facile transformations of the tertiary amine moiety.^{13, 14} Almost all modifications of this moiety have resulted in the attenuation of the antibacterial activity of this class of compound.^{14b} Furthermore, coupling of such desosamine modifications with cladinose sugar removal will result in compounds devoid of antibacterial activity,¹⁶ a property of the parent macrolides undesirable to our goals. Toward this end, we synthesized compound **8**, which incorporates a SAHA-like moiety (Fig. 1) into the 3'-tertiary amine of azithromycin, as a prototype molecule. Compound **8** was synthesized from azithromycin **5** and methyl 8-chloro-8-oxooctanoate **1** through a five-step synthetic route as shown in Fig. 1. To gain some preliminary insights into the roles of key macrolide moieties in anti-HDAC activity of **8**, we synthesized compound **10**, an analog of **8** lacking the cladinose sugar. Acid promoted removal of the cladinose sugar according to a published protocol¹³ quantitatively yielded compound **9**, which was converted to the hydroxamate **10** using the same protocol for the synthesis of **8** from **7** (Fig. 1). We then tested for the HDAC inhibition activity of **8** and **10** using a cell free kit assay,¹⁷ and found that both compounds caused a concentration dependent inhibition of HDACs 1 and 2 from HeLa cell nuclear extract (Table 1). In fact, both compounds have identical anti-HDAC activity, with IC₅₀ values in low nanomolar range. Conversely, compound **7**, the methyl ester precursor of **8**, is completely inactive in this assay. This result suggests that the 15-membered azalide ring of azithromycin macrolide is a suitable non-peptidyl surrogate for the macrocyclic peptide moiety of a typical cyclic-peptide HDACi. Additionally, the binding orientations of these

compounds at the HDAC active site may be such that the hydroxamate group is efficiently presented to chelate the active site Zn^{2+} ion while the cladinoso moiety is oriented away from the enzyme's outer rim.

We then initiated structure activity relationship studies on **8** and **10** to optimize the HDAC binding affinity of these compounds. Preliminarily, we focused on the effects of the modification of the linker-cap group connection moiety, macrolide skeleton, and linker length on anti-HDAC activity. We used the AutoDock program¹⁸ and the crystal structure of a histone deacetylase-like protein (HDLP)⁷ to guide our structural optimization. In an earlier study, we demonstrated the suitability of a 1,2,3-triazole ring as an alternative linker-cap group connection moiety in SAHA-like HDAC inhibitors.¹⁹ In addition to serving as an alternative connection moiety with a potentially favorable pharmacokinetic properties, the triazole ring is a more synthetically tractable group,²⁰ and has been adopted in the construction of active-site directed chemical probes for profiling HDAC activities in proteomes and live cells.²¹ Its incorporation into our design could facilitate a facile, high yielding synthesis of the triazole-based SAHA-macrolide conjugates and thus provide solution to the problems of low yields that currently plagued the synthesis of the amide-based compounds **8** and **10** (Fig. 1).

Accordingly, we proceeded to test the compatibility of the triazole ring with the anti-HDAC activity of **8** and **10**. We prepared compounds **16a** and **16b**, analogs of **8** and **10** respectively, having the amide moiety connecting the cap group and the linker region substituted with a 1, 2, 3-triazole ring (Fig. 2a). The reaction of 4-ethynylbenzyl mesylate with **11** led to the desired desosamine alkylation product **12** in good yields. Copper (I) catalyzed cycloaddition reaction of alkyne **12** with O-silyl azido-hydroxamate **14**^{19, 20} gave the desired cycloadduct **15** in good yields. Similarly, the reaction of alkynes **12** and **13** with azido ester **18** gave the desired cycloadducts **19a** and **19b** respectively. The deprotection of the silyl group of compound **15** is accomplished with TBAF treatment, leading to desired hydroxamate **16a** in good yields. However, difficulties were encountered in separating **16a** from the TBAF reagent. In much the same manner as in the synthesis of **8** or **10** (Fig. 1), the conversion of methyl ester **19a** and **19b** to the corresponding hydroxamate is dogged by low yields. Subsequently, we found that a direct Cu(I) catalyzed cycloaddition between unprotected azido-hydroxamate **17** and alkyne **12** or **13** in anhydrous, degassed THF under exclusion of oxygen uneventfully gave the desired hydroxamates (Fig. 2a).

HDAC inhibition studies revealed that the triazolyl-compounds **16a** and **16b** have virtually identical anti-HDAC activity as the amide-compounds **8** and **10** (Table 1). This result is in contrast with our observation on simple aliphatic hydroxamates where the introduction of the triazole ring led to an enhancement of anti-HDAC activity.¹⁹ To gain insights on the molecular interactions between these non-peptide macrocyclic HDACi and HDAC active site, we performed molecular docking analysis of **16b** on HDLP, using the AutoDock program, as previously described.^{18, 19} We chose to use the HDLP structure because it shared conserved active site residues with class I HDACs. Additionally, direct docking experiments using this structure or HDAC1 homology model built from the same HDLP structure have given docking results that are essentially of the same quality and also agreed with experimentally obtained data. Either of these approaches has been extensively used in the literature to interrogate the binding interactions of HDAC inhibitors at the protein active site.^{18b, c} The structures we obtained from our docking experiments indicated interesting molecular surface complementarities between the macrolide skeleton of **16b** and the HDAC outer rim. Previous investigations have shown that there are four possible binding pockets on the HDLP surface whose interactions with the HDACi cap groups could enhance the inhibitor binding ability.^{18b} Compound **16b** adopts a docked structure that placed the macrolide macrocyclic ring in binding pockets 1 and 3 (Fig 3a). In addition, the hydrophobic components of the macrolide ring optimally interact with the hydrophobic residues in pockets 1 and 3 while the hydrophilic

hydroxyl groups are oriented away from the pocket's hydrophobic residues. Compared to the structure of SAHA, the hydroxamate moiety of **16b** is farther oriented from the active site Zn^{2+} ion (Fig 3b). Based on the preceding observation, we inferred that optimization of the linker region could result in compounds with enhanced HDAC affinity due to a better presentation of the hydroxamate moiety to the catalytic Zn^{2+} ion. Subsequent docking analyses with analogs of **16b** having varied methylene-linker lengths revealed that **16d**, a C_7 -linker compound, optimally interacts with the Zn^{2+} ion (Fig 3c). Interestingly, compound **24d**, an analog of **16d** in which the 15-membered azithromycin ring has been substituted with the 14-membered clarithromycin ring, has a slight preference for the enzyme. A closer analysis of the docked structures of **16d** and **24d** revealed that the C12 - C14 region of the larger 15-membered ring is about 0.5 Å closer to the phenyl ring of Phe338 that defines one of the hydrophobic pockets at the enzyme outer rim compared to that of the 14-membered compound (Fig 3d). This might compromise the binding affinity of the 15-membered compounds relative to the 14-membered analogs.

To experimentally test these *in silico* observations, we synthesized compounds **16c-h** and **24a-h**, the 14- and 15-membered non-peptide macrocyclic hydroxamates respectively (Fig. 2a and 2b). Results from HDAC inhibition assay on these compounds revealed HDAC inhibition activities that essentially paralleled the *in silico* prediction (Table 1). The compounds displayed both linker-length and macrolide-type dependent HDAC inhibition activities. For compounds derived from the same macrolide ring, an increase in the linker length from C_6 to C_7 conferred a better anti HDAC activity. Further linker length increase did not improve HDAC inhibition activity; in fact such an increase is detrimental to function in some cases. For compounds with C_6 and C_7 linkers, a head-to-head comparison between 14- and 15-membered macrolides revealed that the 14-membered compounds are about 2-5 folds better HDACi than their 15-membered counterparts (Table 1, see **16c** and **24c** for example). However, this preference dissipated with increase in linker length. This is presumably due to a relief of steric clash between the macrocyclic ring and the phenyl ring of Phe338 at the enzyme's outer rim, conferred by the longer linkers.

To obtain evidence for the HDAC isoform selectivity of the macrocyclic HDACi described herein, we tested their HDAC8 inhibition activity. We chose HDAC8 because it is in the same sub-class as HDACs 1 and 2, the principal HDACs contained in the HeLa cell nuclear extract used in assay kit employed in this study. There are very few examples of HDAC inhibitors that are selective for HDAC isoforms within the same class; hence this choice should permit a quick, yet rigorous assessment of HDAC isoform selectivity of our compounds. Compared to SAHA, all non-peptide macrocyclic hydroxamates tested are more selective for HDAC1/2. In particular, C_7 -linked, 14-membered compounds **24c** and **24d** are several folds more selective than their 15-membered counterparts (Table 1). Although HDLP and HDAC8 shared similar amino acid sequences and topology at the active site, observation from the analysis of X-ray data however revealed significant inhibitor specific changes in the enzyme active site topology of HDAC8.^{8b} Our docking analysis operates in the rigid receptor mode and it is incapable of capturing such crystallographically observed ligand induced conformational changes. Nevertheless, we performed molecular docking analysis of **16b** on HDAC8 structure reported by Somoza *et al.*^{8b} In contrast to its docked structure on HDLP, the orientation of **16b** which has a chance of making any interaction with the active site Zn^{2+} ion is that which adopted a closed conformation nestled atop of the entrance to the enzyme active site. In this conformation, the linker group wrapped around the macrocyclic ring to orient the hydroxamate moiety toward the Zn^{2+} ion, albeit much farther away to make any stabilizing interaction (see Fig. S1, supporting information). An alternative lower energy confirmation of **16b** oriented the hydroxamate moiety away from the Zn^{2+} ion. It is therefore possible that the observed isoform selectivity may be due to the inability of the nonpeptide macrocyclic hydroxamates herein described to efficiently induce active site conformational changes that facilitate HDAC8

specific inhibitor association with the enzyme active site.^{22a} Alternatively, HDAC8 activity has been observed to depend on the sequence of its peptide substrate,^{22b} hence it is also conceivable that the extent of enzyme inhibition, and consequently isoform selectivity, may depend on the substrate used in the inhibition study.

To further test for HDAC isoform selectivity, we investigated the effect of selected macrocyclic HDACi on the deacetylase activity of HDAC6, a representative member of class II HDAC. For this preliminary study, we chose compounds **16c** and **16d**, and **24c** and **24d**, the most potent representative 15- and 14-membered macrocyclic HDACi respectively (Table 1). Cell free HDAC6 inhibition assay was performed as recommended by the supplier.¹⁷ Unlike SAHA which equally inhibited HDAC1/2 and HDAC6, we observed that the macrocyclic HDACi still displayed significant preference for HDAC1/2 (see Table S1, supporting information). This observation is in agreement with the literature reports which suggest that complex head groups tend to promote isoform selectivity.^{3, 22c}

To screen for the whole cell activity of compounds described in this study, we studied their effect on the viability of SK-MES-1 (human NSCLC cell line), NCI-H69 (human SCLC cell line), DU-145 (a human prostate cancer cell line,) and non-transformed human primary lung fibroblasts and mammary epithelial cell lines. The non-transformed cell lines were investigated to obtain evidence for compound selective toxicity. Drug concentrations necessary for 50 % inhibition of cell viability (EC₅₀) were quantitatively measured using trypan blue exclusion,²³ as previously described.¹⁹ Table 2 shows the EC₅₀ values of each compound. The EC₅₀ values obtained for SAHA are in close agreement with the reported values under similar experimental conditions.²⁴ Macrocyclic methyl ester **7**, the precursor to compound **8** (Fig. 1), has no effect on cell viability. This result may not be unexpected since compound **7** has no HDAC inhibition activity (Table 1). However, all macrocyclic hydroxamates inhibit the proliferation of all transformed cells studied. Most importantly, compounds **16b**, **24c**, and **24d** are at least twice as potent as SAHA in DU-145 cells (Table 2). Gratifyingly, none of the macrocyclic hydroxamates tested shows any growth inhibitory effects on the normal human primary lung fibroblast and mammary epithelial cell lines at concentrations in excess of 10 μ M. This data showed that the macrocyclic compounds are selectively toxic to the transformed cells, a trait that tracks with that of many HDACi.

An important biomarker that is primarily associated with intracellular HDAC inhibition is the expression levels of p21^{WAF1/CIP1} gene. Up-regulation of p21^{WAF1/CIP1} gene has been generally observed with cellular HDAC inhibition.^{25, 26} As part of study aimed at cellular target and mechanistic validation, we investigated the effect of **24d**, a representative macrocyclic HDACi, on the intracellular status of p21^{WAF1/CIP1} protein in SK-MES-1 cells. We used SAHA as a positive control for HDAC inhibition. Cultured cells were exposed to various concentrations of our test agents for 8 hours and the cellular p21^{WAF1/CIP1} expression levels were determined by Western blotting according to literature protocol.²⁷ We found that both **24d** and SAHA induced a dose dependent increase in p21^{WAF1/CIP1} expression (Fig. 4). This result provided additional evidence that the likely mechanism of the antiproliferative activities of the non-peptide macrocyclic hydroxamates described here is through intracellular HDACs inhibition.

Conclusion

We have identified a new class of non-peptide HDACi derived from the macrocyclic skeletons of clinically useful macrolides. These compounds will enable a molecular description of the interaction between the HDAC enzyme's outer rim and the inhibitors' macrocyclic cap-groups thereby further aid our understanding of the roles of this interaction in inhibitors binding affinity and possibly HDAC isoform selectivity. In addition, due to the selective tissue

distribution that may be conferred by the appended macrolide moiety, some of these HDACi are anticipated to have targeted anti-cancer activity. Specifically, compounds incorporating azithromycin skeleton could be selectively accumulated in the lungs,²⁸ thereby possess lung-selective anti-cancer activity. The prospect of tissue-specific HDACi delivery is a particularly enticing alternative to isoform selective HDACi and could lead to the identification of new chemotherapeutic agents for use in targeted cancer therapy applications. Efforts are underway in our lab to profile the tissue distribution of the new class of HDACi described here.

Experimental Section

Clarithromycin and azithromycin were purchased from Greenfield Chemical and Pfizer, respectively. 4-Ethynylbenzyl alcohol, ethyl 6-bromohexanoate and ethyl 7-bromoheptanoate, 8-bromooctanoic acid and methyl 10-bromodecanoate were purchased from Sigma Aldrich. 9-bromononanoic acid was purchased from Karl Industries Inc. Common reaction solvents were either high performance liquid chromatography (HPLC) grade or American Chemical Society (ACS) grade, and used without further purification. HDAC Fluorimetric Assay kit and recombinant HDACs were procured from BIOMOL® International, PA. Analtech silica gel plates (60 F₂₅₄) were used for analytical TLC, and Analtech preparative TLC plates (UV 254, 2000 µm) were used for purification. UV light was used to examine the spots. 200-400 Mesh silica gel was used in column chromatography. Nuclear magnetic resonance (NMR) spectra were recorded on a Varian-Gemini 400 magnetic resonance spectrometer. ¹H NMR spectra were recorded in parts per million (ppm) relative to the peak of CDCl₃, (7.24 ppm), CD₃OD (3.31 ppm), DMSO-d₆ (2.49 ppm) or acetone-d₆ (2.04 ppm). ¹³C spectra were recorded relative to the central peak of the CDCl₃ triplet (77.0 ppm), CD₃OD (49.0 ppm), DMSO-d₆ septet (39.7 ppm), or acetone-d₆ (2.04 ppm) and were recorded with complete hetero-decoupling. High-resolution mass spectra were recorded at the Georgia Institute of Technology mass spectrometry facility in Atlanta. 4-Ethynylbenzyl methylsulfonate **11** and azido alkylesters were synthesized by adapting literature protocol.²⁹⁻³¹ 6-Azido-*O*-silyl hexahydroxamate **14** was prepared from the corresponding azido carboxylic acid, t-BuPh₂SiCl and NaH, according to the procedure described by Muri *et al.*³²

Synthesis of Methyl 8-(4-(hydroxymethyl)phenylamino)-8-oxooctanoate (3)

To a solution of (4-aminophenyl)methanol **2** (0.314 g, 2.500 mmol) in anhydrous pyridine (5 mL) was added chlorotrimethylsilane (0.32 mL, 2.500 mmol) at room temperature and stirring continued for 2 h. The reaction was cooled in ice bath to 0°C and to the mixture was added methyl 8-chloro-8-oxooctanoate **1** (0.32 mL, 2.200 mmol) and a catalytic amount of DMAP. The reaction was allowed to warm to room temperature and stirring continued overnight. Water (5 mL) and 1 M TBAF in tetrahydrofuran (THF) (0.25 mL, 0.250 mmol) were added and stirring continued for additional 30 min. EtOAc (50 mL) and 1N HCl (30 mL) were added, the two layers were separated, the organic layer was washed with 1N HCl (30 mL) and saturated brine (30 mL) and dried over Na₂SO₄. Solvent was evaporated off and the crude was purified by preparative TLC, eluting with EtOAc/hexanes 2:1 to give 275 mg (43 %) compound **3** as a yellow-white solid. ¹H-NMR (DMSO-d₆, 400 MHz) δ 1.26 (4H, m), 1.47-1.56 (4H, m), 2.26 (4H, m), 3.55 (3H, s), 4.40 (2H, d, *J* = 5.6 Hz), 5.06 (1H, t, *J* = 5.6 Hz), 7.19 (2H, d, *J* = 8.8 Hz), 7.51 (2H, d, *J* = 8.8 Hz), 9.79 (1H, s); ¹³C NMR (DMSO-d₆, 100MHz) δ 24.3, 24.9, 28.2, 28.3, 33.2, 36.3, 51.2, 62.6, 118.8, 126.9, 136.9, 138.0, 171.2, 173.4; HRMS (FAB, thioglycerol) calc for [C₁₆H₂₃NO₄ + H]⁺ 294.1705, found 294.1652

4'-Desmethyl azithromycin (6)

To a solution of azithromycin (2.000 g, 2.548 mmol) and sodium acetate (1.78 g, 21.500 mmol) in 80% aqueous methanol (30 mL) at 90 °C was added iodine (0.700 g, 2.756 mmol) in three batches within 5 min. The reaction was maintained at pH 8-9 by addition of 1M NaOH (2 mL,

once at 10 min of reaction time) and stirring continued for 3 h. The reaction was poured into cold water containing 5% sodium thiosulfate (80 mL) and extracted with CH_2Cl_2 (2×40 mL). The aqueous layer was basified with concentrated NH_4OH , extracted with 10 % MeOH in CH_2Cl_2 (3×40 mL) and the organic layer was dried over Na_2SO_4 . Solvent was evaporated off to give 1.57 g of compound **6** as an off white solid (>90 % purity, TLC: $\text{CHCl}_3/\text{MeOH}/\text{NH}_4\text{OH}$ 12:1:0.1). The crude **6** was used without further purification.

Azithromycin-arylalkyl methyl ester (7)

To a solution of compound **3** (0.175 g, 0.597 mmol) in CH_2Cl_2 (7 mL) and triethylamine (Et_3N) (0.24 mL, 1.800 mmol) was added mesyl chloride (0.10 mL, 1.200 mmol) at 0°C and the reaction was allowed to warm to room temperature. Stirring continued for 1 h during which TLC revealed a quantitative conversion into a higher R_f product. CH_2Cl_2 (40 mL) and saturated sodium bicarbonate (30 mL) were added, the two layers were separated. The organic layer was washed with sodium bicarbonate (2×30 mL), saturated brine (30 mL) and dried over Na_2SO_4 . Solvent was evaporated off to give compound **4** as white solid.

A mixture of 4'-desmethylazithromycin **6** (0.315 g, 0.430 mmol) and crude compound **4**, in anhydrous DMSO (7 mL) and Hunig's base (0.7 mL) was stirred at 85°C for 1.5 h. The reaction was cooled and diluted with EtOAc (60 mL) and washed with saturated NaHCO_3 (40 mL) and saturated brine (40 mL). The organic layer was dried over Na_2SO_4 , concentrated *in vacuo* and the crude was purified by preparative TLC, eluting with $\text{EtOAc}/\text{hexanes}/\text{Et}_3\text{N}$ 3:2:0.1 to give 152 mg (35 %) of compound **7** as a white solid. $^1\text{H-NMR}$ (CDCl_3 , 400MHz) δ 0.78-0.82 (7H, m), 0.94-1.03 (11H, m), 1.09-1.32 (22H, m), 1.34-1.58 (5H, m), 1.59- 1.73 (4H, m), 1.77-2.01 (5H, m), 2.12 (3H, s), 2.20-2.28 (8H, m), 2.42-2.53 (2H, m), 2.54-2.71 (3H, m), 2.87-3.01 (2H, m), 3.11 (3H, s), 3.25-3.33 (3H, m), 3.42 (2H, m), 3.54-3.65 (6H, m), 3.97 (1H, m), 4.18 (1H, m), 4.36 (1H, d, $J = 6.8$ Hz), 4.61 (2H, m), 5.03 (1H, d, $J = 4.4$ Hz), 7.14 (2H, d, $J = 8.4$ Hz), 7.42 (2H, d, $J = 8.4$ Hz), 7.73 (1H, s), 8.97 (1H, bs); $^{13}\text{C-NMR}$ (CDCl_3 , 100MHz) δ 7.5, 9.1, 11.3, 14.8, 16.2, 16.9, 18.2, 20.5, 21.2, 21.4, 21.5, 21.9, 24.6, 25.3, 26.7, 27.5, 28.7, 29.6, 33.9, 34.7, 36.2, 36.7, 37.3, 39.1, 41.9, 42.2, 45.1, 48.5, 49.3, 51.4, 57.3, 62.2, 64.3, 65.4, 68.5, 69.9, 70.5, 72.7, 73.5, 73.8, 74.1, 77.7, 77.9, 83.4, 94.4, 102.6, 119.5, 129.0, 134.2, 137.0, 171.0, 173.8, 178.4 MS; HRMS (FAB, *mnba*) calc for $[\text{C}_{53}\text{H}_{91}\text{N}_3\text{O}_{15} + \text{H}]^+$ 1010.6529, found 1010.6450

Azithromycin-arylalkyl hydroxamic acid (8)

To a solution of compound **7** (0.050 g, 0.050 mmol) in 1:1 THF/MeOH (2 mL) was added hydroxylamine (50 % in H_2O) (0.07 mL, 1.260 mmol) and a catalytic amount of KCN. The mixture was stirred at room temperature for 24 h. The reaction was partitioned between 5 % MeOH in CH_2Cl_2 (30 mL) and saturated sodium bicarbonate (25 mL), the two layers were separated and the aqueous layer was extracted with 5 % MeOH in CH_2Cl_2 (2×20 mL). The combined organic layer was washed with saturated brine (40 mL) and dried over Na_2SO_4 . Solvent was evaporated off and the crude was purified by preparative TLC, eluting with $\text{CH}_2\text{Cl}_2/\text{MeOH}/\text{NH}_4\text{OH}$ 10:1:0.1 to give 26 mg (52 %) compound **8** as brown-white solid. $^1\text{H-NMR}$ (CD_3OD , 400MHz) δ 0.87-0.92 (6H, m), 1.03-1.12 (12H, m), 1.17-1.37 (m), 1.43-1.69 (m), 1.75-1.88 (6H, m), 1.99 (4H, m), 2.08 (3H, m), 2.13-2.19 (3H, m), 2.24-2.41 (13H, m), 2.54 (1H, d, $J = 11.2$ Hz), 2.75-2.80 (4H, m), 3.00 (1H, d, $J = 9.6$ Hz), 3.19 (4H, m), 3.47-3.51 (1H, m), 3.60-3.78 (6H, m), 4.14-4.22 (3H, m), 4.50 (1H, d, $J = 7.2$ Hz), 5.02 (1H, d, $J = 4.8$ Hz), 7.29 (2H, d, $J = 8.0$ Hz), 7.49 (2H, d, $J = 8.4$ Hz); $^{13}\text{C-NMR}$ (Acetone- d_6 , 100MHz) δ 7.6, 9.7, 11.6, 14.3, 15.3, 17.4, 18.9, 21.7, 21.9, 22.2, 22.5, 26.0, 26.1, 27.3, 27.9, 30.9, 32.6, 33.2, 35.5, 36.5, 36.8, 37.6, 43.0, 43.1, 46.1, 49.6, 58.6, 63.1, 64.1, 64.5, 66.2, 68.9, 70.0, 71.6, 73.3, 73.6, 74.2, 74.9, 75.3, 78.0, 78.9, 84.3, 95.6, 103.7, 119.8, 130.0, 135.0, 139.5, 171.9, 178.8; HRMS (EI) calc for $[\text{C}_{52}\text{H}_{90}\text{N}_4\text{O}_{15} + \text{H}]^+$ 1011.6481, found 1011.6580

Desclasinose azithromycin-arylalkyl hydroxamate (10)

A mixture of compound **7** (0.050 g, 0.050 mmol) in 0.25 N HCl (15 mL) was stirred at room temperature for 20 h and poured into EtOAc (20 mL). The two layers were separated and the aqueous layer was washed with EtOAc (2 × 20 mL), basified with concentrated NH₄OH and then extracted with 5 % MeOH in CH₂Cl₂ (2 × 30 mL). The combined organic layer was washed with saturated brine (30 mL) and dried over Na₂SO₄. Solvent was evaporated off to give compound **9** which was used for the next reaction without further purification.

To a solution of compound **9** in 1:1 THF/MeOH (2 mL) was added hydroxylamine (50 % in H₂O) (0.05 mL, 0.790 mmol) and a catalytic amount of KCN. The mixture was stirred at room temperature for 24 h. The reaction was partitioned between 5 % MeOH in CH₂Cl₂ (30 mL) and saturated brine (20 mL), the two layers were separated and the organic layer was dried over Na₂SO₄. Solvent was evaporated off and the crude was purified by preparative TLC, eluting with CH₂Cl₂/MeOH/NH₄OH 9:1:0.1 to give 7 mg (16 %) compound **10** as brown-white solid. ¹H-NMR (CD₃OD, 400MHz) δ 0.78 (3H, m), 0.85 (3H, d, *J* = 7.2 Hz), 0.91 (3H, d, *J* = 8.0 Hz), 0.99 (3H, s), 1.08-1.14 (10H, m), 1.18-1.79 (20H, m), 1.87 (2H, m), 1.98 (2H, t, *J* = 7.4 Hz), 2.05-2.13 (2H, m), 2.17 (3H, s), 2.25 (2H, t, *J* = 7.4 Hz), 2.52-2.65 (4H, m), 2.95 (1H, bs), 3.24 (m), 3.37-3.56 (6H, m), 3.67 (1H, d, *J* = 13.2 Hz), 4.53 (2H, d, *J* = 7.6 Hz), 7.19 (2H, d, *J* = 8.4 Hz), 7.39 (2H, d, *J* = 8.4 Hz); HRMS (EI) calc for [C₄₄H₇₆N₄O₁₂ + H]⁺ 853.5538, found 853.5488

4'-Ethynylbenzyl azithromycin (12)

To a solution of 4'-desmethylazithromycin **6** (0.940 g, 1.280 mmol) in anhydrous DMSO (15 ml) was added Hunig's base (1.5 ml) and 4-ethynylbenzyl methanesulfonate **11** (0.380 g, 1.800 mmol). The reaction mixture was heated with stirring under argon at 85°C for 2 h. The reaction was cooled and diluted with EtOAc (100 mL) and washed with saturated NaHCO₃ (3 × 60 mL) and saturated brine (60 mL). The organic layer was dried over Na₂SO₄ and concentrated *in vacuo*. The crude product was purified by flash chromatography (silica, 12:1:0.05 CH₂Cl₂/MeOH/conc. NH₄OH) to give 850 mg (78%) of **12** as a brown-white solid. ¹H-NMR (CDCl₃, 400MHz) δ 0.86 (6H, m), 0.99 (3H, d, *J* = 7.6 Hz), 1.05 (7H, m), 1.11-1.32 (19H, m), 1.36-1.55 (2H, m), 1.68-1.77 (2H, m), 1.81-2.07 (4H, m), 2.22 (3H, s), 2.27-2.31 (4H, m), 2.51 (2H, m), 2.65-2.75 (2H, m), 2.87 (1H, bs), 2.98 (1H, t, *J* = 9.8 Hz), 3.04 (1H, s), 3.10 (3H, s), 3.29-3.34 (2H, m), 3.40-3.47 (2H, m), 3.58 (1H, d, *J* = 6.8 Hz), 3.65 (1H, s), 3.74 (1H, d, *J* = 13.2 Hz), 3.99 (1H, m), 4.21 (1H, dd, *J* = 2 Hz, 4.4 Hz), 4.38 (1H, d, *J* = 7.2 Hz), 4.65 (1H, dd, *J* = 2.8 Hz, 10 Hz), 5.06 (1H, d, *J* = 4.4 Hz), 7.23 (2H, d, *J* = 8 Hz), 7.41 (2H, d, *J* = 8 Hz); ¹³C-NMR (CDCl₃, 100MHz) δ 7.5, 9.2, 11.3, 14.9, 16.3, 18.3, 21.3, 21.4, 21.6, 22.0, 26.8, 27.6, 29.6, 34.7, 36.3, 36.9, 42.0, 42.3, 45.2, 49.2, 57.7, 62.3, 63.7, 65.4, 68.5, 70.0, 70.6, 72.7, 73.5, 73.8, 74.2, 77.1, 77.9, 78.0, 83.4, 83.7, 94.5, 102.6, 120.8, 128.5, 132.0, 139.6, 178.3; HRMS (FAB, mnb) calc for [C₄₆H₇₆N₂O₁₂ + H]⁺ 849.5476, found 849.5411

Descladinose-4'-ethynylbenzyl azithromycin (13)

A solution of 4'-ethynylbenzyl azithromycin **12** (0.12 g, 0.14 mmol) in 0.25 N HCl (15 mL) was stirred at room temperature for 20 h and poured into EtOAc (20 mL). The two layers were separated and the aqueous layer was washed with EtOAc (2 × 20 mL), basified with concentrated NH₄OH and then extracted with 5 % MeOH in CH₂Cl₂ (2 × 30 mL). The combined organic layer was washed with saturated brine (30 mL) and dried over Na₂SO₄. Solvent was evaporated off to give 89 mg (91%) of descladinose compound **12** as a white solid. ¹H-NMR (CDCl₃, 400MHz) δ 0.80-1.52 (16H, m), 1.65-1.90 (2H, m), 1.97 (3H, s), 2.20-2.30 (6H, m), 2.42-2.69 (3H, m), 2.77 (1H, s), 2.89 (1H, s), 3.02 (1H, s), 3.18 (1H, s), 3.30-3.37 (3H, m), 3.49-3.63 (6H, m), 3.72 (2H, d, *J* = 10.6 Hz), 3.84 (1H, s), 3.89 (2H, s), 4.04 (3H, q, *J* = 14.4, 7.2 Hz), 4.41 (1H, d, *J* = 7.2 Hz), 4.64 (1H, d, *J* = 10.8 Hz), 7.15 (2H, d, *J* = 8.0 Hz), 7.37 (2H, d, *J* = 8.4 Hz); ¹³C NMR (CDCl₃, 100 MHz) δ 7.7, 7.9, 10.9, 14.2,

16.1, 20.9, 21.0, 21.2, 25.8, 26.6, 29.2, 35.9, 36.4, 37.0, 42.0, 44.5, 57.6, 60.3, 62.4, 65.3, 69.8, 70.4, 70.9, 73.0, 74.1, 75.4, 79.4, 83.3, 94.9, 106.4, 120.8, 128.2, 131.9, 139.2, 171.0, 177.2; HRMS (FAB, thioglycerol) calc for $[C_{38}H_{62}N_2O_9 + H]^+$ 691.4533, found 691.4513

Azithromycin-*N*-benzyltriazolyl-*O*-silyl-hexahydroxamate (15)

4'-Ethynylbenzyl azithromycin **12** (0.045 g, 0.050 mmol) and 6-azido-*O*-silyl hexahydroxamate **14** (0.060 g, 0.146 mmol) were dissolved in anhydrous THF (5 mL) and stirred under argon at room temperature. Copper (I) iodide (0.010 g, 0.050 mmol), Hunig's base (0.5 mL) and tris[(1-benzyl-1*H*-1,2,3-triazol-4-yl)methyl]amine, (TBTA) (0.016 g, 0.030 mmol) were then added to the reaction mixture, and stirring continued for 2 h. The reaction mixture was diluted with CH_2Cl_2 (40 mL) and washed with 1:4 NH_4OH /saturated NH_4Cl (2 × 30 mL) and saturated NH_4Cl (30 mL). The organic layer was dried over Na_2SO_4 and concentrated *in vacuo*. The crude product was purified by prep TLC (silica, 12:1:0.1 CH_2Cl_2 /MeOH/conc. NH_4OH) to give 38 mg (60%) of silyl protected compound **15** as a brown-white solid. 1H -NMR ($CDCl_3$, 400MHz) δ 0.82-1.52 (39H, m), 1.71- 2.04 (13H, m), 2.21 (3H, s), 2.28 (6H, s), 2.44-2.57 (2H, m), 2.65-2.70 (2H, m), 2.96 (1H, br s), 3.09 (2H, s), 3.31-3.35 (2H, m), 3.40-3.48 (6H, m), 3.56-3.60 (1H, m), 3.76 (1H, d, J = 13.2 Hz), 4.00 (1H, br s), 4.20 (3H, br s), 4.39 (1H, d, J = 6.8 Hz), 4.64 (1H, d, J = 9.2 Hz), 5.09 (1H, br s) 7.29-7.41 (10H, m), 7.64-7.83 (5H, m); ^{13}C NMR ($CDCl_3$, 100 MHz) δ 7.4, 9.1, 11.3, 14.7, 16.3, 18.2, 21.3, 21.4, 21.5, 22.0, 26.7, 27.5, 29.6, 29.7, 29.9, 34.6, 36.2, 36.9, 42.2, 45.2, 49.3, 49.9, 57.6, 62.4, 64.1, 65.4, 68.5, 69.9, 70.5, 72.7, 73.5, 74.1, 77.7, 77.9, 83.4, 94.3, 102.7, 119.2, 125.5, 127.6, 129.0, 135.5, 147.1, 178.4; HMRS (ESI) calcd for $[C_{68}H_{106}N_6O_{14}Si + H]^+$ 1259.7609, found 1259.7570

Methyl azithromycin-*N*-benzyltriazolylhexanoate (19a)

4'-Ethynylbenzyl azithromycin **12** (0.045 g, 0.053 mmol) and azido-ester **18** (0.014 g, 0.080 mmol) were dissolved in anhydrous THF (5 mL) and stirred under argon at room temperature. Copper (I) iodide (0.010 g, 0.053 mmol), and Hunig's base (0.05 mL) were then added to the reaction mixture, and stirring continued for 12 h. The reaction mixture was diluted with CH_2Cl_2 (30 mL) and washed with 1:4 NH_4OH /saturated NH_4Cl (3 × 25 mL) and again with saturated NH_4Cl (25 mL). The organic layer was dried over Na_2SO_4 and concentrated under vacuum. The crude product was purified by preparative TLC, eluting with Hexane/EtOAc/ Et_3N 3:2:0.1 to give 50 mg (92%) of **19a** as a white-brown solid. 1H -NMR ($CDCl_3$, 400MHz) δ 0.82-0.90 (3H, m), 0.98 (3H, d, J = 7.6 Hz), 1.05-1.52 (24H, m), 1.60-1.74 (6H, m), 1.80-2.06 (9H, m), 2.22-2.37 (9H, m), 2.56 (3H, m), 2.67 (3H, m), 2.95 (2H, t, J = 9.8 Hz), 3.07 (3H, m), 3.29-3.34 (1H, m), 3.46 (3H, m), 3.54 (1H, d, J = 6.8 Hz), 3.61 (3H, s), 3.68 (1H, bs), 3.77 (1H, m), 3.97 (1H, m), 4.18 (1H, m), 4.34-4.38 (3H, m), 4.69 (1H, m), 5.06 (1H, d, J = 4 Hz), 7.32 (2H, d, J = 6.4 Hz), 7.73-7.75 (3H, m); ^{13}C -NMR ($CDCl_3$, 100MHz) δ 8.7, 9.2, 11.3, 14.2, 14.7, 16.5, 18.2, 21.4, 21.5, 22.2, 24.2, 25.9, 26.6, 27.3, 29.7, 30.0, 33.6, 34.6, 36.4, 36.9, 42.4, 45.3, 45.8, 49.3, 50.0, 51.5, 57.7, 63.9, 65.5, 68.6, 69.4, 70.5, 72.7, 73.8, 74.2, 77.2, 77.6, 78.0, 83.4, 94.4, 102.7, 119.3, 125.5, 129.1, 129.4, 147.2, 173.4, 178.1. HRMS (FAB, mnba) calc for $[C_{53}H_{89}N_5O_{14} + H]^+$ 1020.6484, found 1020.6430

Methyl descladinoseazithromycin-*N*-benzyltriazolylhexanoate (19b)

Compound **13** (0.080 g, 0.115 mmol) and azido-ester **18** (0.030 g, 0.173 mmol) were dissolved in anhydrous THF (5 mL) and stirred under argon at room temperature. Copper (I) iodide (0.010 g, 0.053 mmol), and Hunig's base (0.05 mL) were then added to the reaction mixture, and stirring continued for 12 h. The reaction mixture was diluted with CH_2Cl_2 (30 mL) and washed with 1:4 NH_4OH /saturated NH_4Cl (3 × 25 mL) and again with saturated NH_4Cl (25 mL). The organic layer was dried over Na_2SO_4 and concentrated *in vacuo*. The crude product was purified by preparative TLC, eluting with Hexane/EtOAc/ Et_3N 3:2:0.1 to give 65 mg (65%)

of **19b** as a white-brown solid. $^1\text{H-NMR}$ (CDCl_3 , 400MHz) δ 0.79-0.86 (6H, m), 1.00-1.07 (6H, m), 1.17-1.26 (m), 1.42-1.51 (m), 1.55-1.72 (6H, m), 1.80-1.94 (6H, m), 2.00-2.05 (2H, m), 2.1 (3H, s), 2.23-2.27 (4H, m), 2.33 (3H, s), 2.47 (1H, d, $J = 10.4$ Hz), 2.58-2.72 (5H, m), 3.32-3.41 (3H, m), 3.52-3.73 (6H, m), 3.92-4.00 (2H, m), 4.34 (2H, t, $J = 7.0$ Hz), 4.41 (1H, d, $J = 7.6$ Hz), 4.69 (1H, d, $J = 10.8$ Hz), 7.24 (2H, d, $J = 8.4$ Hz), 7.71 (2H, d, $J = 8$ Hz), 7.73 (1H, s); $^{13}\text{C NMR}$ (CDCl_3 , 100MHz) δ 7.7, 7.9, 8.7, 10.9, 16.1, 16.1, 20.9, 21.2, 24.2, 25.8, 25.9, 26.6, 29.2, 29.6, 30.0, 33.6, 36.0, 36.3, 37.1, 42.0, 44.5, 45.8, 50.0, 51.5, 57.7, 62.6, 65.1, 69.9, 70.4, 73.1, 74.1, 75.3, 79.4, 94.8, 106.4, 119.3, 125.5, 128.9, 129.6, 138.2, 147.1, 173.4, 177.2. HRMS (EI) calc for $[\text{C}_{45}\text{H}_{75}\text{N}_5\text{O}_{11} + \text{H}]^+$ 862.5541, found 862.5566

Representative Procedure for Conversion of Methyl or Ethyl Ester to Hydroxamic Acid. 6-Azidohexahydroxamic acid (**17a**)

To a solution of ethyl 6-azidohexanoate (1.00 g, 5.840 mmol) in 1:1 THF (5 mL) and anhydrous methanol (5 mL) was added aqueous hydroxylamine (4 mL, 70.100 mmol) and KCN (0.070 g, 1.170 mmol), and the stirring continued for 24 h. The reaction was diluted with EtOAc (30 mL) and washed with saturated NaHCO_3 (2×30 mL) and saturated brine (30 mL). The organic layer was dried over Na_2SO_4 and concentrated *in vacuo* to give 797 mg (80%) of **17a** as white solid. $^1\text{H-NMR}$ ($\text{DMSO}-d_6$, 400MHz) δ 1.24-1.30 (2H, m), 1.45-1.53 (4H, m), 1.93 (2H, t, $J = 7.2$ Hz), 3.29 (2H, t, $J = 6.8$ Hz), 8.65 (1H, s), 10.3 (1H, s); $^{13}\text{C NMR}$ (CDCl_3 , 100 MHz) δ 24.8, 26.1, 28.4, 32.6, 51.1, 171.4; HRMS (ESI) calc for $[\text{C}_6\text{H}_{12}\text{N}_4\text{O}_2 + \text{H}]^+$ 173.0947, found 173.0983

7-Azidoheptahydroxamic acid (**17b**)

Reaction of ethyl 7-azidoheptanoate (1.00 g, 5.400 mmol) and aqueous hydroxylamine (4 mL, 70.100 mmol) within 24 h as described for the synthesis of **17a**, gave 820 mg (82%) of **17b** as white solid. $^1\text{H NMR}$ ($\text{DMSO}-d_6$, 400 MHz) δ 1.23-1.32 (4H, m), 1.45-1.52 (4H, m), 1.92 (2H, t, $J = 7.6$ Hz), 3.29 (2H, t, $J = 7.2$ Hz), 8.63 (1H, br s), 10.3 (1H, br s); $^{13}\text{C NMR}$ (CDCl_3 , 100 MHz) δ 25.1, 26.2, 28.5, 28.6, 32.7, 51.2, 171.6; HRMS (FAB, thioglycerol) calc for $[\text{C}_7\text{H}_{14}\text{N}_4\text{O}_2 + \text{H}]^+$ 187.1195, found 187.1163

8-Azidooctahydroxamic acid (**17c**)

Reaction of methyl 8-azidooctanoate (0.580 g, 2.910 mmol) and aqueous hydroxylamine (2.49 mL, 37.800 mmol) within 24 h as described for the synthesis of **17a**, gave 432 mg (74%) of **17c** as white solid. $^1\text{H NMR}$ (CDCl_3 , 400 MHz) δ 1.33-1.38 (6H, m), 1.54-1.63 (4H, m), 2.13 (2H, t, $J = 7.6$ Hz), 3.24 (2H, t, $J = 6.8$ Hz), 8.85 (1H, br); $^{13}\text{C NMR}$ (CDCl_3 , 100MHz) δ 25.5, 26.7, 29.0, 29.1, 29.2, 33.1, 51.6, 172.0; HRMS (FAB, thioglycerol) calc for $[\text{C}_8\text{H}_{16}\text{N}_4\text{O}_2 + \text{H}]^+$ 201.1351, found 201.1352

9-Azidononahydroxamic acid (**17d**)

Reaction of methyl 9-azidononanoate (0.290 g, 1.370 mmol) and aqueous hydroxylamine (1.18 mL, 17.900 mmol) within 24 h as described for the synthesis of **17a**, gave 225 mg (77%) of **17d** as white solid. $^1\text{H NMR}$ (CDCl_3 , 400 MHz) δ 1.30-1.40 (8H, m), 1.56-1.61 (4H, m), 2.13 (2H, t, $J = 7.2$ Hz), 3.25 (2H, t, $J = 8$); $^{13}\text{C NMR}$ (CDCl_3 , 100MHz) δ 25.6, 26.8, 29.0, 29.2, 29.3, 33.1, 51.6, 172.2; HRMS (FAB, thioglycerol) calc for $[\text{C}_9\text{H}_{18}\text{N}_4\text{O}_2 + \text{H}]^+$ 215.1508, found 215.1529

10-Azidodecahydroxamic acid (**17e**)

Reaction of methyl 10-azidodecanoate (0.300 g, 1.310 mmol) and aqueous hydroxylamine (1.13 mL, 17.100 mmol) within 24 h as described for the synthesis of **17a**, gave 254 mg (84 %) of **17e** as white solid. $^1\text{H NMR}$ (CDCl_3 , 400 MHz) δ 1.24-1.39 (8H, m), 1.54-1.65 (4H, m), 2.12 (2H, t, $J = 7.6$ Hz), 3.24 (2H, t, $J = 7.2$ Hz), 9.0 (1H, br); $^{13}\text{C NMR}$ (CDCl_3 , 100MHz)

δ 25.6, 26.9, 29.0, 29.2, 29.3, 29.4, 29.5, 33.2, 51.7, 172.2 ; HRMS (FAB, thioglycerol) calc for $[C_{10}H_{20}N_4O_2 + H]^+$ 229.1665, found 229.1666

Azithromycin-*N*-benzyltriazolyhexahydroxamic acid (**16a**)

Method A—To a solution of compound **19a** (0.040 g, 0.040 mmol) in 1:1 THF/MeOH (3 mL) was added hydroxylamine (50 % in H₂O) (0.03 mL, 0.540 mmol) and a catalytic amount of KCN. The mixture was stirred at room temperature for 24 h. The reaction was partitioned between 5 % MeOH in CH₂Cl₂ (30 mL) and saturated sodium bicarbonate (25 mL), the two layers were separated and the aqueous layer was extracted with 5 % MeOH in CH₂Cl₂ (2 × 20 mL). The combined organic layer was washed with saturated brine (40 mL) and dried over Na₂SO₄. Solvent was evaporated off and the crude was purified by preparative TLC, eluting with CH₂Cl₂/MeOH/NH₄OH 10:1:0.1 to give 6.5 mg (16 %) of compound **16a** as a brown-white solid.

Method B—4'-Ethynylbenzyl azithromycin **12** (0.100 g, 0.109 mmol) and 6-azido-hexahydroxamic acid **17a** (0.081 g, 0.117 mmol) were dissolved in anhydrous THF (5 mL) and stirred under argon at room temperature. Copper (I) iodide (0.011 g, 0.070 mmol) and Hunig's base (0.5 mL) were then added to the reaction mixture, and stirring continued for 4 h. The reaction mixture was diluted with CH₂Cl₂ (40 mL) and washed with 1:4 NH₄OH/saturated NH₄Cl (3 × 30 mL) and saturated NH₄Cl (30 mL). The organic layer was dried over Na₂SO₄ and concentrated *in vacuo*. The crude product was purified by prep TLC (silica, 12:1:0.1 CH₂Cl₂/MeOH/conc. NH₄OH) to give 71 mg (59%) of **16a** as a brown-white solid.

Method C—To a solution of silyl protected compound **15** (0.025 g, 0.020 mmol) in THF (1 mL) was added 1 M TBAF in THF (0.030 mL, 0.030 mmol) and the mixture was stirred at room temperature for 2 h during which TLC revealed a near quantitative conversion to a lower R_f product. The reaction was partitioned between CH₂Cl₂ (30 mL) and saturated NH₄Cl (25 mL), the two layers were separated and the organic layer dried over Na₂SO₄ and concentrated *in vacuo*. The crude product was purified by prep TLC (silica, 12:1:0.1 CH₂Cl₂/MeOH/Et₃N) to give 15 mg (73%) of **16a** as a brown-white solid. ¹H-NMR (Acetone-d₆, 400MHz) δ 0.83-0.92 (6H, m), 1.02 (3H, d, *J* = 7.6 Hz), 1.08-1.11 (8H, m), 1.14 (3H, d, *J* = 7.6 Hz), 1.18 (3H, d, *J* = 6 Hz), 1.24-1.29 (15H, m), 1.33-1.47 (3H, m), 1.54 (1H, dd, *J* = 4.8 Hz, 15.2 Hz), 1.60-1.69 (5H, m), 1.80-2.01 (m), 2.06-2.12 (1H, m), 2.18-2.24 (1H, m), 2.26 (3H, s), 2.28-2.31 (1H, m), 2.35-2.41 (4H, m), 2.51 (1H, d, *J* = 10 Hz), 2.65-2.96 (m), 3.12 (3H, s), 3.22-3.29 (1H, m), 3.47 (1H, m), 3.54-3.69 (6H, m), 3.81 (1H, d, *J* = 13.2 Hz), 4.11 (1H, m), 4.24 (1H, m), 4.45 (3H, t, *J* = 7.0 Hz), 4.50 (1H, d, *J* = 6.8 Hz), 4.75 (1H, d, *J* = 7.2 Hz), 4.97 (1H, d, *J* = 5.2 Hz), 7.42 (2H, d, *J* = 8.0 Hz), 7.84 (2H, d, *J* = 8.0 Hz), 8.35 (1H, s). ¹³C NMR (CDCl₃, 100 MHz) δ 6.5, 8.7, 11.5, 14.4, 16.7, 17.7, 21.3, 21.6, 21.8, 24.6, 25.7, 26.7, 27.0, 29.2, 29.6, 29.9, 33.0, 34.5, 35.6, 36.7, 41.8, 42.7, 45.3, 49.3, 50.0, 53.4, 57.8, 62.6, 63.4, 65.9, 68.6, 69.3, 70.4, 72.6, 73.3, 73.8, 77.8, 78.3, 78.4, 83.4, 94.5, 102.8, 119.6, 125.7, 129.3, 129.7, 138.6, 147.4, 171.3, 178.4; HMRS (ESI) calcd for $[C_{52}H_{88}N_6O_{14} + H]^+$ 1021.6437, found 1021.6409

Desclasinose azithromycin-*N*-benzyltriazolyhexahydroxamic acid (**16b**)

Method A—To a solution of compound **19b** (0.040 g, 0.050 mmol) in 1:1 THF/MeOH (3 mL) was added hydroxylamine (50 % in H₂O) (0.04 mL, 0.540 mmol) and a catalytic amount of KCN. The mixture was stirred at room temperature for 24 h. The reaction was partitioned between 5 % MeOH in CH₂Cl₂ (30 mL) and saturated sodium bicarbonate (25 mL), the two layers were separated and the aqueous layer was extracted with 5 % MeOH in CH₂Cl₂ (2 × 20 mL). The combined organic layer was washed with saturated brine (40 mL) and dried over Na₂SO₄. Solvent was evaporated off and the crude was purified by preparative TLC, eluting with CH₂Cl₂/MeOH/NH₄OH 10:1:0.1 to give 9.0 mg (23 %) of compound **16b** as brown-white solid.

Method B—Reaction of descladinose compound **13** (0.134 g, 0.188 mmol) and 6-azidoheptahydroxamic acid **17a** (0.130 g, 0.755 mmol) within 8 h, according to the protocols of **Method B** described for the synthesis of compound **16a**, followed by prep TLC (silica, 10:1:0.1 CH₂Cl₂/MeOH/conc. NH₄OH) gave 73 mg (43%) of **16b** as a brown-white solid. ¹H-NMR (Acetone-d₆, 400MHz) δ 0.82-0.90 (9H, m), 1.02 (3H, d, *J* = 7.2 Hz), 1.07 (3H, s), 1.09 (3H, d, *J* = 6.8 Hz), 1.18-1.23 (m), 1.28 (3H, s), 1.31-1.39 (m), 1.46-1.56 (4H, m), 1.61-1.65 (3H, m), 1.80-1.99 (7H, m), 2.05-2.11 (2H, m), 2.18-2.21 (1H, m), 2.24 (3H, s), 2.25-2.29 (1H, m), 2.35 (3H, s), 4.47 (1H, d, *J* = 9.2 Hz), 2.61-2.67 (1H, m), 2.70-2.77 (1H, m), 3.30-3.34 (1H, m), 3.41 (1H, m), 3.52-3.65 (5H, m), 3.81 (1H, d, *J* = 13.2 Hz), 4.44 (2H, t, *J* = 7.0 Hz), 4.59 (1H, d, *J* = 7.6 Hz), 4.87 (1H, dd, *J* = 1.8 Hz, 11.0 Hz), 7.43 (2H, d, *J* = 8.4 Hz), 7.83 (2H, d, *J* = 8.4 Hz), 8.34 (1H, s); ¹³C NMR (CDCl₃, 100 MHz) δ 7.3, 7.9, 10.7, 16.0, 16.2, 20.9, 21.1, 24.4, 25.6, 25.7, 26.5, 29.0, 29.6, 29.7, 35.8, 36.3, 36.8, 42.1, 44.4, 50.0, 57.9, 62.7, 64.0, 69.8, 70.5, 70.8, 73.3, 74.1, 75.0, 79.5, 94.8, 106.5, 120.0, 125.8, 129.2, 129.5, 138.4, 147.3, 170.5, 177.5; HMRS (ESI) calcd for [C₄₄H₇₄N₆O₁₁ + H]⁺ 863.5494, found 863.5544

Azithromycin-*N*-benzyltriazolylheptahydroxamic acid (**16c**)

Reaction of 4'-ethynylbenzyl azithromycin **12** (0.134 g, 0.158 mmol) and 7-azidoheptahydroxamic acid **17b** (0.125 g, 0.672 mmol) within 4 h, according to the protocols of **Method B** described for the synthesis of compound **16a**, followed by prep TLC (silica, 12:1:0.1 CH₂Cl₂/MeOH/conc. NH₄OH) gave 93 mg (56%) of **16c** as a brown-white solid. ¹H-NMR (CDCl₃, 400MHz) δ 0.81-1.51 (30H, m), 1.54-1.65 (6H, m), 1.70-2.14 (9H, m), 2.20-2.38 (9H, m), 2.46-2.56 (2H, m), 2.60-2.70 (2H, m), 3.00 (3H, s), 3.31 (2H, t, *J* = 8.8 Hz), 3.38-3.54 (6H, m), 3.60 (1H, s), 3.78 (1H, d, *J* = 12.8 Hz), 3.98-4.20 (2H, m), 4.36 (3H, d, *J* = 7.2 Hz), 4.49 (1H, d, *J* = 7.2 Hz), 5.11 (1H, d, *J* = 4.0 Hz), 7.32 (2H, d, *J* = 7.6 Hz), 7.73 (1H, s), 7.75 (2H, d, *J* = 7.6 Hz); ¹³C NMR (CDCl₃, 100 MHz) δ 6.6, 8.8, 11.5, 14.4, 16.6, 17.7, 21.3, 21.6, 21.8, 25.1, 26.0, 26.7, 27.1, 28.2, 29.2, 29.6, 30.0, 33.1, 34.5, 35.7, 36.7, 41.8, 42.7, 45.3, 49.3, 50.3, 50.7, 57.9, 62.7, 63.0, 65.8, 68.6, 69.4, 70.4, 72.6, 73.2, 73.8, 77.8, 78.1, 78.2, 83.5, 94.4, 102.8, 119.3, 125.7, 129.4, 129.7, 138.4, 147.4, 171.3, 178.4; HMRS (ESI) calcd for [C₅₃H₉₀N₆O₁₄ + H]⁺ 1035.6587, found 1035.6628

Descladinose azithromycin-*N*-benzyltriazolylheptahydroxamic acid (**16d**)

Reaction of descladinose-4'-ethynylbenzyl azithromycin **13** (0.130 g, 0.188 mmol) and 7-azidoheptahydroxamic acid **17b** (0.130 g, 0.755 mmol) within 8 h, according to the protocols of **Method B** described for the synthesis of compound **16a**, followed by prep TLC (silica, 10:1:0.1 CH₂Cl₂/MeOH/conc. NH₄OH) gave 78 mg (47%) of **16d** as a brown-white solid. ¹H-NMR (CDCl₃, 400MHz) δ 0.66-2.32 (42H, m), 2.47 (2H, d, *J* = 10.8 Hz), 2.63-2.70 (4H, m), 3.34-3.51 (6H, m), 3.62-3.69 (5H, m), 4.20-4.40 (5H, m), 4.74 (2H, br s), 7.26 (2H, br s), 7.73 (3H, br s); ¹³C NMR (CDCl₃, 100 MHz) δ 7.4, 7.9, 10.7, 16.0, 16.1, 20.9, 21.1, 25.0, 25.7, 26.5, 28.0, 28.9, 29.8, 35.8, 36.3, 36.9, 42.0, 44.4, 50.1, 57.9, 62.7, 63.9, 69.9, 70.4, 70.8, 73.3, 74.1, 75.2, 79.5, 94.9, 106.6, 119.7, 125.7, 129.2, 129.6, 138.4, 147.3, 177.5; HMRS (ESI) calcd for [C₄₅H₇₆N₆O₁₁ + H]⁺ 877.5645, found 877.5665

Azithromycin-*N*-benzyltriazolylloctahydroxamic acid (**16e**)

Reaction of 4'-ethynylbenzyl azithromycin **12** (0.100 g, 0.120 mmol) and 8-azido-octahydroxamic acid **17c** (0.047 g, 0.240 mmol) within 2.5 h, according to the protocols of **Method B** described for the synthesis of compound **16a**, followed by prep TLC (silica, 12:1:0.1 CH₂Cl₂/MeOH/conc. NH₄OH) gave 72 mg (58 %) of **16e** as brown-white solid. ¹H NMR (CDCl₃, 400 MHz) δ 0.85 (3H, t, *J* = 4.0 Hz), 0.87-1.22 (18H, m), 1.29 (9H, s), 1.30-2.28 (18H, m), 2.29 (6H, s), 2.30-3.00 (8H, m), 3.10 (3H, s), 3.20-3.79 (9H, m), 3.99-4.03 (1H, m), 4.35-4.40 (3H, m), 4.65 (1H, d, *J* = 8.0 Hz), 5.11 (1H, d, *J* = 4.8 Hz), 7.34 (2H, d, *J* = 8.0 Hz),

7.72 (1H, s), 7.77 (2H, d, $J = 8.0$ Hz); ^{13}C NMR (DMSO- d_6 , 100MHz) δ 7.5, 9.8, 11.6, 15.4, 18.2, 19.1, 21.5, 22.1, 22.7, 25.6, 26.3, 26.6, 28.7, 29.0, 29.6, 30.2, 32.0, 32.8, 35.2, 36.4, 37.2, 42.2, 45.3, 49.2, 50.1, 58.3, 63.2, 65.4, 67.7, 70.8, 73.3, 74.2, 77.0, 78.4, 83.4, 102.8, 121.6, 125.6, 129.7, 130.0, 135.0, 147.0, 177.8; HRMS (FAB, thioglycerol) calc for $[\text{C}_{54}\text{H}_{92}\text{N}_6\text{O}_{14} + \text{H}]^+$ 1049.6749, found 1049.6648

Descladinose azithromycin-*N*-benzyltriazolyloctahydroxamic acid (16f)

Reaction of descladinose-4'-ethynylbenzyl azithromycin **13** (0.100 g, 0.144 mmol) and 8-azidoctahydroxamic acid **17c** (0.049 g, 0.246 mmol) within 2.5 h, according to the protocols of **Method B** described for the synthesis of compound **16a**, followed by prep TLC (silica, 10:1:0.1 $\text{CH}_2\text{Cl}_2/\text{MeOH}/\text{conc. NH}_4\text{OH}$) gave 94 mg (73 %) of **16f** as a brown-white solid. ^1H NMR (CDCl_3 , 400 MHz) δ 0.84-0.91 (9H, m), 1.04-1.11 (9H, m), 1.23-1.30 (12H, m), 1.37-2.16 (14H, m), 2.2 (3H, s), 2.35 (3H, s), 2.49-2.74 (5H, m), 3.22-3.75 (6H, m), 4.10-4.12 (1H, m), 4.34-4.42 (3H, m), 4.77 (1H, d, $J = 12$ Hz), 7.30 (2H, d, $J = 7.6$ Hz), 7.79 (3H, m); ^{13}C NMR (CDCl_3 , 100MHz) δ 7.6, 8.2, 11.0, 14.4, 16.2, 16.4, 21.1, 21.4, 25.2, 25.9, 26.1, 26.7, 28.4, 28.8, 28.9, 29.1, 29.9, 30.2, 36.1, 36.6, 42.3, 44.6, 50.5, 51.6, 58.2, 60.6, 63.0, 64.0, 70.1, 70.7, 73.6, 74.4, 75.4, 79.7, 95.1, 106.8, 120.0, 126.0, 129.5, 129.9, 138.6, 147.6, 170.6, 177.8; HRMS (FAB, thioglycerol) calc for $[\text{C}_{46}\text{H}_{78}\text{N}_6\text{O}_{11} + \text{H}]^+$ 891.5806, found 891.5776

Azithromycin-*N*-benzyltriazolylnonahydroxamic acid (16g)

Reaction of 4'-ethynylbenzyl azithromycin **12** (0.100 g, 0.120 mmol) and 9-azidononahydroxamic acid **17d** (0.043 g, 0.200 mmol) within 2.5 h, according to the protocols of **Method B** described for the synthesis of compound **16a**, followed by prep TLC (silica, 12:1:0.1 $\text{CH}_2\text{Cl}_2/\text{MeOH}/\text{conc. NH}_4\text{OH}$) gave 64 mg (51 %) of **16g** as brown-white solid. ^1H NMR (CDCl_3 , 400 MHz) δ 0.84-1.30 (30H, m), 1.33-2.26 (20H, m), 2.30 (6H, s), 2.38-2.68 (8H, m), 2.99 (3H, s), 3.32-3.84 (9H, m), 4.03-4.08 (1H, m), 4.35-4.41 (3H, m), 4.53 (1H, d, $J = 8.0$ Hz), 5.13 (1H, d, $J = 4.0$ Hz), 7.35 (2H, d, $J = 8.0$ Hz), 7.75 (1H, s), 7.78 (2H, d, $J = 8.0$ Hz); ^{13}C NMR (DMSO- d_6 , 100MHz) δ 6.9, 9.0, 11.6, 14.7, 16.9, 18.0, 21.6, 21.8, 22.1, 25.6, 26.4, 26.9, 27.3, 28.8, 29.0, 29.1, 29.5, 29.9, 30.4, 34.8, 36.0, 37.0, 42.1, 43.0, 45.6, 49.5, 50.5, 58.1, 63.5, 66.1, 68.8, 70.7, 72.9, 74.1, 78.1, 78.3, 78.5, 83.7, 94.4, 94.7, 103.1, 119.6, 126.0, 129.7, 130.0, 147.6, 178.7; HRMS (ESI) calc for $[\text{C}_{55}\text{H}_{94}\text{N}_6\text{O}_{14} + \text{H}]^+$ 1063.6900, found 1063.6861

Azithromycin-*N*-benzyltriazolyldecahydroxamic acid (16h)

Reaction of 4'-ethynylbenzyl azithromycin **12** (0.100 g, 0.120 mmol) and 10-azidodecahydroxamic acid **17e** (0.045 g, 0.20 mmol) within 4.5 h, according to the protocols of **Method B** described for the synthesis of compound **16a**, followed by prep TLC (silica, 12:1:0.1 $\text{CH}_2\text{Cl}_2/\text{MeOH}/\text{conc. NH}_4\text{OH}$) gave 70 mg (56 %) of **16h** as a brown-white solid. ^1H NMR (CDCl_3 , 400 MHz) δ 0.85-1.36 (30H, m), 1.41-2.24 (22H, m), 2.28, 2.36 (6H, s), 2.33-3.10 (8H, m), 3.05 (3H, s), 3.23-3.82 (9H, m), 4.06-4.10 (1H, m), 4.36-4.41 (3H, m), 4.49 (1H, d, $J = 8.0$ Hz), 5.15 (1H, d, $J = 4.0$ Hz), 7.34 (2H, d, $J = 8$ Hz), 7.75 (1H, s), 7.78 (2H, d, $J = 8.0$ Hz); ^{13}C NMR (DMSO- d_6 , 100MHz) δ 6.7, 9.0, 11.7, 14.6, 16.9, 17.9, 21.6, 21.9, 22.0, 25.6, 26.4, 26.8, 27.0, 27.3, 28.8, 29.0, 21.9, 29.3, 29.5, 29.9, 30.3, 33.6, 34.9, 35.8, 37.0, 42.1, 43.0, 45.6, 49.6, 50.6, 51.6, 58.0, 62.8, 63.9, 66.2, 68.9, 69.6, 70.7, 72.9, 73.5, 74.0, 74.1, 78.2, 78.6, 83.6, 94.6, 103.0, 119.6, 126.0, 129.6, 129.9, 138.9, 147.6, 178.6; HRMS (MALDI) calc for $[\text{C}_{56}\text{H}_{96}\text{N}_6\text{O}_{14} + \text{H}]^+$ 1077.7057, found 1077.6971

4'-Desmethyl clarithromycin (20)

To a solution of clarithromycin (3.320 g, 4.440 mmol) and sodium acetate (3.280 g, 39.900 mmol) in 80% aqueous methanol (50 mL) at 85 °C was added iodine (1.240 g, 4.890 mmol)

in three batches within 5 min. The reaction was maintained at pH 8-9 by additions of 1M NaOH (2×3 ml, once at 10 min and 45 min of reaction time). Stirring was continued at 85 °C for 3 h and TLC analysis indicated about 90% consumption of the starting material. A solution of 5% sodium thiosulfate (120 mL) and dichloromethane (80 mL) were added and the two layers were separated. The aqueous layer was extracted with CH₂Cl₂ (60 mL), the combined organic layers was washed with saturated brine, dried over Na₂SO₄, and concentrated *in vacuo* to give 2.4 g of **20**, which was used without further purification.

4'-Ethynylbenzyl clarithromycin (21)

To a solution of 4'-desmethyl clarithromycin **20** (2.400 g, 3.340 mmol) in anhydrous DMSO (30 ml) was added Hunig's base (3 ml) and 4-ethynylbenzyl methanesulfonate **11** (0.920 g, 4.340 mmol). The reaction mixture was then heated with stirring under argon at 85°C for 2.5 h. The reaction was cooled and diluted with EtOAc (100 mL) and washed with saturated NaHCO₃ (3×60 mL) and saturated brine (60 mL). The organic layer was dried over Na₂SO₄ and concentrated *in vacuo*. The crude product was purified by flash chromatography (silica, gradient 12:1; 10:1; 8:1; CH₂Cl₂/acetone) to give 1.8 g (63%) of **21** as a brown-white solid. ¹H-NMR (CDCl₃, 400MHz) δ 0.82 (3H, t, $J = 7.2$ Hz), 1.03-1.28 (18H, m), 1.37 (3H, s), 1.40-1.55 (3H, m), 1.65-1.90 (6H, m), 2.03 (1H, d, $J = 10.0$ Hz), 2.22 (3H, s), 2.30 (1H, d, $J = 15.2$ Hz), 2.40-2.60 (2H, m), 2.80-2.90 (2H, m), 2.94-3.00 (6H, m), 3.04 (1H, s), 3.09 (3H, s), 3.16 (1H, s), 3.24-3.29 (1H, m), 3.38-3.46 (3H, m), 3.59 (1H, d, $J = 6.8$ Hz), 3.70-3.75 (3H, m), 3.88-3.95 (1H, m), 4.37 (1H, d, $J = 7.2$ Hz), 4.88 (1H, d, $J = 4.4$ Hz), 5.02 (1H, dd, $J = 11.6, 2.4$ Hz), 7.23 (2H, d, $J = 12.0$ Hz), 7.42 (2H, d, $J = 8.0$ Hz); ¹³C NMR (CDCl₃, 100 MHz) δ 9.2, 10.7, 12.4, 16.1, 18.1, 18.7, 19.9, 21.1, 21.5, 29.3, 32.4, 34.8, 36.9, 37.2, 39.2, 45.0, 45.2, 49.3, 50.6, 53.4, 57.6, 63.3, 65.6, 68.5, 69.0, 70.6, 72.5, 74.2, 76.5, 77.8, 78.1, 78.2, 80.8, 95.8, 102.5, 120.9, 128.6, 132.0, 133.5, 139.4, 175.4; HRMS (ESI) calc for [C₄₆H₇₃NO₁₃ + H]⁺ 848.5155, found 848.5181

Descladinose-4'-ethynylbenzyl clarithromycin (22)

To a solution of 4'-ethynylbenzyl clarithromycin **21** (0.500 g, mmol) in ethanol (20 mL) was added 1N HCl (20 mL), and stirring continued for 22 h at room temperature. The reaction mixture was basified with concentrated NH₄OH to about pH = 9. The reaction mixture was diluted with distilled water (40 mL) and extracted with EtOAc (3×60 mL). The combined organic layers were washed with saturated brine (40 mL), dried over Na₂SO₄, and concentrated *in vacuo*. The crude product was purified by flash chromatography (silica, 8:1 CH₂Cl₂/acetone) to give 320 mg (79%) of **22** as a brown-white solid. ¹H-NMR (CDCl₃, 400MHz) δ 0.82 (3H, t, $J = 7.6$ Hz), 1.09-1.28 (12H, m), 1.34 (3H, s), 1.40-1.55 (3H, m), 1.70-1.74 (2H, m), 1.87-1.94 (3H, m), 2.08-2.15 (6H, m), 2.54-2.66 (2H, m), 2.94-2.98 (3H, m), 3.05 (1H, s), 3.25 (1H, s), 3.31-3.42 (2H, m), 3.48-3.56 (2H, m), 3.66 (2H, d, $J = 10.0$ Hz), 3.82 (1H, s), 3.90 (1H, s), 4.35 (1H, d, $J = 7.6$ Hz), 5.14 (1H, dd, $J = 10.8, 2.0$ Hz), 7.18 (2H, d, $J = 8.0$ Hz), 7.42 (2H, d, $J = 8.4$ Hz); ¹³C NMR (CDCl₃, 100 MHz) δ 8.4, 10.5, 12.7, 15.3, 16.3, 17.8, 18.8, 21.4, 29.2, 35.9, 36.6, 37.5, 38.7, 44.5, 45.5, 49.6, 57.8, 65.0, 69.7, 70.1, 70.6, 74.1, 77.9, 78.9, 83.3, 88.5, 106.5, 121.0, 128.4, 132.1, 139.1, 174.7; HRMS (ESI) calc for [C₃₈H₅₉NO₁₀ + H]⁺ 690.4212, found 690.4259

Clarithromycin-*N*-benzyltriazolyl-*O*-silyl-hexahydroxamate (23)

4'-Ethynylbenzyl clarithromycin **21** (0.100 g, 0.118 mmol) and 6-azido-*O*-silyl hexahydroxamate **14** (0.103 g, 0.251 mmol) were dissolved in anhydrous THF (6 mL) and stirred under argon at room temperature. Copper (I) iodide (0.011 g, 0.057 mmol), TBTA (0.060 g, 0.094 mmol), and Hunig's base (0.1 mL) were then added to the reaction mixture, and stirring continued for 24 h. The reaction mixture was diluted with CH₂Cl₂ (30 mL) and washed with 1:4 NH₄OH/saturated NH₄Cl (3×30 mL) and saturated NH₄Cl (30 mL). The

organic layer was dried over Na_2SO_4 and concentrated *in vacuo*. The crude product was purified by prep TLC (silica, 20:1 $\text{CH}_2\text{Cl}_2/\text{MeOH}$) to give 147 mg (99%) of compound **23** as a brown-white solid. $^1\text{H-NMR}$ (CDCl_3 , 400MHz) δ 0.81 (3H, t, $J = 7.2$ Hz), 1.03-1.51 (30H, m), 1.67-1.88 (12H, m), 2.10-2.29 (8H, m), 2.53-2.58 (3H, m), 2.82-2.90 (2H, m), 2.93-3.01 (6H, m), 3.10 (3H, s), 3.17 (2H, s), 3.28-3.32 (1H, m), 3.40-3.48 (3H, m), 3.60-3.78 (4H, m), 3.91-3.97 (1H, m), 4.22 (1H, br s), 4.41 (2H, d, $J = 8.0$ Hz), 4.87 (1H, d, $J = 4.8$ Hz), 5.03 (1H, d, $J = 10.8$ Hz), 7.21-7.23 (2H, m), 7.30-7.42 (10H, m), 7.64-7.76 (3H, m); $^{13}\text{C NMR}$ (CDCl_3 , 100 MHz) δ 9.1, 10.6, 12.3, 16.0, 18.0, 18.6, 19.8, 21.0, 21.3, 21.5, 26.7, 29.3, 29.6, 29.9, 34.7, 36.8, 37.1, 39.0, 39.2, 45.0, 45.2, 47.0, 49.3, 49.9, 50.5, 53.4, 54.0, 65.5, 68.5, 68.9, 70.6, 72.3, 74.1, 76.4, 77.7, 78.1, 78.2, 80.7, 95.7, 102.5, 119.3, 123.5, 125.4, 127.7, 128.4, 128.8, 129.1, 133.5, 134.4, 135.5, 143.9, 147.0, 175.4; HMRS (ESI) calcd for $[\text{C}_{68}\text{H}_{103}\text{N}_5\text{O}_{15}\text{Si} + \text{H}]^+$ 1258.7292, found 1258.7313

Clarithromycin-*N*-benzyltriazolylhexahydroxamic acid (**24a**)

Method A—4'-Ethynylbenzyl clarithromycin **21** (0.100 g, 0.120 mmol) and 6-azidoheptahydroxamic acid **17a** (0.080 g, 0.470 mmol) were dissolved in anhydrous THF (5 mL) and stirred under argon at room temperature. Copper (I) iodide (0.011 g, 0.057 mmol) and Hunig's base (0.5 mL) were added to the reaction mixture, and stirring continued for 2.5 h. The reaction mixture was diluted with CH_2Cl_2 (50 mL) and washed with 1:4 NH_4OH /saturated NH_4Cl (3×30 mL) and saturated NH_4Cl (30 mL). The organic layer was dried over Na_2SO_4 and concentrated *in vacuo*. The crude product was purified by prep TLC (silica, 12:1:0.1 $\text{CH}_2\text{Cl}_2/\text{MeOH}/\text{conc. NH}_4\text{OH}$) to give 70 mg (58%) of **24a** as a brown-white solid.

Method B—To a solution of compound **23** (0.085g, 0.067 mmol) in anhydrous THF (1.5 mL) was added 1M TBAF in THF (0.1 mL, 0.100 mmol). The reaction was stirred under argon for 2 h. The reaction mixture was diluted with 5% MeOH in dichloromethane (10 mL) and washed with saturated NH_4Cl (15mL). The organic layer was dried over Na_2SO_4 and concentrated *in vacuo*. The crude product was purified on prep TLC (silica, 12:1:0.1 $\text{CH}_2\text{Cl}_2/\text{MeOH}/\text{Et}_3\text{N}$). The purified product was dissolved in 5% MeOH in CH_2Cl_2 (10 mL) and washed with distilled water (10 mL) and saturated brine (10 mL) to remove the last trace of associated TBAF. The organic layer was dried over Na_2SO_4 and concentrated *in vacuo* to give 26 mg (38%) of **24a** as a brown-white product.

$^1\text{H-NMR}$ (CDCl_3 , 400MHz) δ 0.81 (3H, t, $J = 7.6$ Hz), 1.03-1.52 (21H, m), 1.62-1.92 (12H, m), 2.04-2.29 (8H, m), 2.48-2.60 (3H, m), 2.82-2.90 (2H, m), 2.93-2.99 (6H, m), 3.09 (3H, s), 3.19 (2H, s), 3.28-3.33 (1H, m), 3.42-3.46 (3H, m), 3.60 (1H, d, $J = 7.6$ Hz), 3.70-3.80 (3H, m), 3.90-3.98 (1H, m), 4.37-4.40 (3H, m), 4.87 (1H, d, $J = 4.8$ Hz), 5.03 (1H, dd, $J = 11.6, 2.4$ Hz), 7.34 (2H, d, $J = 7.6$ Hz), 7.77 (2H, d, $J = 7.6$ Hz), 7.82 (1H, s); $^{13}\text{C NMR}$ (CDCl_3 , 100 MHz) δ 9.1, 10.5, 12.2, 15.9, 17.9, 18.5, 19.7, 20.9, 21.2, 21.4, 24.3, 25.5, 29.4, 29.6, 34.7, 36.8, 37.1, 39.0, 39.1, 45.0, 45.1, 49.3, 49.9, 50.5, 53.3, 57.5, 63.6, 65.5, 68.5, 69.0, 70.7, 72.4, 74.2, 77.8, 78.2, 80.9, 95.9, 102.6, 119.8, 125.6, 129.4, 147.4, 175.8; HMRS (ESI) calcd for $[\text{C}_{52}\text{H}_{85}\text{N}_5\text{O}_{15} + \text{H}]^+$ 1020.6114, found 1020.6121

Descladinose clarithromycin-*N*-benzyltriazolylhexahydroxamic acid (**24b**)

Reaction of descladinose-4'-ethynylbenzyl clarithromycin **22** (0.075 g, 0.109 mmol) and 6-azidoheptahydroxamic acid **17a** (0.040 g, 0.233 mmol) within 4 h, according to the protocols of **Method A** described for the synthesis of compound **24a**, followed by prep TLC (silica, 10:1:0.1 $\text{CH}_2\text{Cl}_2/\text{MeOH}/\text{conc. NH}_4\text{OH}$) gave 47 mg (51%) of **24b** as a brown-white solid. $^1\text{H-NMR}$ (CDCl_3 , 400MHz) δ 0.79 (3H, t, $J = 7.2$ Hz), 1.08-1.32 (22H, m), 1.39-1.64 (3H, m), 1.71-1.81 (2H, m), 1.82-1.96 (3H, m), 2.04-2.18 (6H, m), 2.51-2.70 (4H, m), 2.92-2.98 (3H, m), 3.18-3.38 (3H, m), 3.45-3.55 (2H, m), 3.60-3.74 (3H, m), 3.81 (3H, s), 3.90 (1H, s), 4.33 (3H, br s), 5.13 (1H, d, $J = 10.4$ Hz), 7.29 (2H, br s), 7.74 (2H, br s); $^{13}\text{C NMR}$ (CDCl_3 ,

100 MHz) δ 8.6, 10.7, 12.9, 15.5, 16.4, 18.0, 19.0, 21.5, 21.6, 29.5, 29.9, 36.1, 36.7, 37.7, 39.0, 44.7, 45.7, 49.8, 50.3, 58.2, 64.6, 70.0, 70.3, 70.9, 74.4, 78.3, 79.1, 88.5, 106.7, 120.3, 126.1, 129.6, 129.9, 147.7, 175.4; HMRS (ESI) calcd for $[C_{44}H_{71}N_5O_{12} + H]^+$ 862.5172, found 862.5155

Clarithromycin-*N*-benzyltriazolylheptahydroxamic acid (**24c**)

Reaction of 4'-ethynylbenzyl clarithromycin **21** (0.130 g, 0.153 mmol) and 7-azidoheptahydroxamic acid **17b** (0.105 g, 0.565 mmol) within 2.5 h, according to the protocols of **Method A** described for the synthesis of compound **24a**, followed by prep TLC (silica, 12:1:0.1 CH_2Cl_2 /MeOH/conc. NH_4OH) gave 105 mg (67%) of **24c** as a brown-white solid. 1H -NMR ($CDCl_3$, 400MHz) δ 0.82 (3H, t, $J = 8.0$ Hz), 1.04-1.52 (21H, m), 1.67-1.92 (14H, m), 2.14-2.29 (8H, m), 2.52-2.60 (3H, m), 2.82-2.90 (2H, m), 2.95-3.00 (6H, m), 3.10 (3H, s), 3.16 (2H, s), 3.27-3.32 (1H, m), 3.41-3.46 (3H, m), 3.59 (1H, d, $J = 6.8$ Hz), 3.69-3.79 (3H, m), 3.90-3.95 (1H, m), 4.34-4.39 (3H, m), 4.87 (1H, d, $J = 4.4$ Hz), 5.02 (1H, d, $J = 9.2$ Hz), 7.33 (2H, d, $J = 6.4$ Hz), 7.77 (3H, d, $J = 8.4$ Hz); ^{13}C NMR ($CDCl_3$, 100 MHz) δ 9.1, 10.5, 12.2, 15.9, 17.9, 18.5, 19.8, 20.9, 21.2, 21.4, 24.7, 25.5, 27.7, 29.8, 34.7, 36.8, 37.1, 39.1, 45.0, 45.2, 49.3, 50.0, 50.5, 57.6, 63.6, 65.6, 68.6, 69.0, 70.7, 72.4, 74.2, 77.8, 78.3, 80.9, 95.9, 102.7, 119.5, 125.7, 129.4, 147.5, 175.8; HMRS (ESI) calcd for $[C_{53}H_{87}N_5O_{15} + H]^+$ 1034.6271, found 1034.6246

Descladinose clarithromycin-*N*-benzyltriazolylheptahydroxamic acid (**24d**)

Reaction of descladinose-4'-ethynylbenzyl clarithromycin **22** (0.075 g, 0.109 mmol) and 7-azidoheptahydroxamic acid **17b** (0.040 g, 0.233 mmol) within 4 h, according to the protocols of **Method A** described for the synthesis of compound **24a**, followed by prep TLC (silica, 10:1:0.1 CH_2Cl_2 /MeOH/conc. NH_4OH) gave 80 mg (84%) of **24d** as a brown-white solid. 1H -NMR ($CDCl_3$, 400MHz) δ 0.78 (3H, t, $J = 7.2$ Hz), 1.06-1.31 (22H, m), 1.40-1.53 (5H, m), 1.71 (2H, d, $J = 11.6$ Hz), 1.80-1.91 (3H, m), 2.01-2.20 (6H, m), 2.50-2.65 (4H, m), 2.91-2.97 (3H, m), 3.16 (1H, t, $J = 6.4$ Hz), 3.26-3.35 (2H, m), 3.42-3.54 (2H, m), 3.64-3.71 (3H, m), 3.80 (1H, br s), 3.90 (1H, br s), 4.30-4.34 (3H, m), 5.12 (1H, dd, $J = 11.6, 2.4$ Hz), 7.27 (2H, d, $J = 8.0$ Hz), 7.72 (2H, d, $J = 7.2$ Hz), 7.80 (1H, s); ^{13}C NMR ($CDCl_3$, 100 MHz) δ 8.3, 10.3, 12.5, 15.2, 16.1, 17.6, 18.6, 21.1, 21.3, 24.8, 25.1, 25.5, 26.2, 27.8, 28.5, 29.1, 29.6, 29.7, 32.3, 35.8, 36.3, 37.4, 38.6, 44.3, 45.4, 49.5, 50.0, 51.2, 57.8, 64.2, 69.7, 69.9, 70.6, 74.1, 77.9, 78.7, 88.0, 106.3, 119.9, 125.7, 129.3, 129.5, 138.3, 147.3, 175.1; HMRS (ESI) calcd for $[C_{45}H_{73}N_5O_{12} + H]^+$ 876.5329, found 876.5301

Clarithromycin-*N*-benzyltriazolylloctahydroxamic acid (**24e**)

Reaction of 4'-ethynylbenzyl clarithromycin **21** (0.101 g, 0.120 mmol) and 8-azidoctahydroxamic acid **17c** (0.047 g, 0.240 mmol) within 2.5 h, according to the protocols of **Method A** described for the synthesis of compound **24a**, followed by prep TLC (silica, 12:1:0.1 CH_2Cl_2 /MeOH/conc. NH_4OH) gave 92 mg (74%) of **24e** as a brown-white solid. 1H NMR ($CDCl_3$, 400 MHz) δ 0.81 (3H, t, $J = 7.2$ Hz), 1.04-2.05 (45H, m), 2.22 (3H, s), 2.19-2.82 (m, 7H), 3.00 (3H, s), 3.08 (3H, s), 2.91-3.80 (6H, m), 3.95 (3H, m), 4.38 (3H, m), 4.88 (1H, d, $J = 4.4$ Hz), 5.04 (1H, dd, $J = 10.8, 2.0$ Hz), 7.33 (2H, d, $J = 7.6$ Hz), 7.71 (1H, s), 7.77 (2H, d, $J = 8.0$ Hz); ^{13}C NMR ($CDCl_3$, 100MHz) δ 8.8, 9.4, 10.8, 12.5, 16.2, 18.2, 18.8, 20.0, 21.2, 21.5, 21.6, 25.1, 26.0, 26.7, 28.2, 28.6, 28.9, 29.9, 30.1, 35.0, 36.9, 37.4, 39.2, 39.4, 45.2, 45.4, 46.1, 49.6, 50.4, 50.8, 51.6, 58.1, 63.6, 65.9, 68.4, 69.3, 70.9, 72.8, 74.4, 78.0, 78.5, 81.2, 96.1, 120.1, 102.6, 126.1, 130.2, 147.4, 176.0 ; HRMS (ESI) calc for $[C_{54}H_{90}N_5O_{15} + H]^+$ 1048.6486, found 1048.6427

Descladinose clarithromycin-*N*-benzyltriazolyloctahydroxamic acid (24f)

Reaction of descladinose-4'-ethynylbenzyl clarithromycin **22** (0.10 g, 0.144 mmol) and 8-azidooctahydroxamic acid **17c** (0.049 g, 0.246 mmol) within 2.5 h, according to the protocols of **Method A** described for the synthesis of compound **24a**, followed by prep TLC (silica, 10:1:0.1 CH₂Cl₂/MeOH/conc. NH₄OH) gave 117 mg (90 %) of **24f** as a brown-white solid. ¹H NMR (CDCl₃, 400 MHz) δ 0.81 (3H, t, *J* = 7.2 Hz), 1.10-2.09 (37H, m), 2.18 (3H, s), 2.19-2.68 (7H, m), 2.98-3.73 (4H, m), 3.83 (2H, s), 3.93 (1H, m), 4.36 (3H, m), 5.16 (1H, d, *J* = 8.0 Hz), 7.31 (2H, d, *J* = 8.0 Hz), 7.77 (2H, d, *J* = 8.0 Hz), 7.79 (1H, s); ¹³C NMR (CDCl₃, 100MHz) δ 8.5, 10.6, 12.8, 15.4, 16.4, 17.9, 18.9, 21.4, 21.6, 29.4, 36.1, 36.7, 37.7, 38.9, 44.7, 45.7, 49.8, 58.0, 65.2, 70.0, 70.4, 70.8, 74.4, 76.8, 78.2, 79.2, 83.6, 88.8, 106.9, 121.3, 128.7, 132.5, 139.6, 175.2; HRMS (FAB, thioglycerol) calc for [C₄₆H₇₆N₅O₁₂ + H]⁺ 890.5490, found 890.5562

Clarithromycin-*N*-benzyltriazolylnonahydroxamic acid (24g)

Reaction of 4'-ethynylbenzyl clarithromycin **21** (0.100 g, 0.120 mmol) and 9-azidononahydroxamic acid **17d** (0.043 g, 0.20 mmol) within 2.5 h, according to the protocols of **Method A** described for the synthesis of compound **24a**, followed by prep TLC (silica, 12:1:0.1 CH₂Cl₂/MeOH/conc. NH₄OH) gave 54 mg (42%) of **24g** as a brown-white solid. ¹H NMR (CDCl₃, 400 MHz) δ 0.81 (3H, t, *J* = 7.2 Hz), 1.04-2.02 (47H, m), 2.24 (3H, s), 2.10-2.97 (7H, m), 3.00, 3.09 (6H, s), 3.20-3.82 (6H, m), 3.88 (3H, m), 4.39 (3H, m), 4.88 (1H, d, *J* = 4.0 Hz), 5.05 (1H, d, *J* = 10.0 Hz), 7.35 (2H, d, *J* = 8.0 Hz), 7.77 (3H, d, *J* = 4.0 Hz); ¹³C NMR (CDCl₃, 100MHz) δ 9.0, 9.3, 10.8, 12.5, 16.1, 18.2, 18.8, 21.2, 21.5, 21.6, 21.7, 21.8, 26.3, 26.8, 28.7, 28.9, 29.1, 29.9, 30.3, 35.0, 37.0, 37.4, 39.3, 39.4, 45.2, 45.4, 49.6, 50.5, 50.8, 51.6, 57.8, 63.8, 65.8, 68.8, 69.2, 70.9, 72.7, 74.5, 76.8, 78.1, 78.4, 78.5, 81.1, 96.1, 102.9, 119.7, 125.9, 129.6, 129.9, 138.7, 147.6, 176.1; HRMS (ESI) calc for [C₅₅H₉₁N₅O₁₅ + H]⁺ 1062.6584, found 1062.6586

Clarithromycin-*N*-benzyltriazolyldecahydroxamic acid (24h)

Reaction of 4'-ethynylbenzyl clarithromycin **21** (0.10 g, 0.120 mmol) and 10-azidodecahydroxamic acid **17e** (0.045 g, 0.197 mmol) within 2.5 h, according to the protocols of **Method A** described for the synthesis of compound **24a**, followed by prep TLC (silica, 12:1:0.1 CH₂Cl₂/MeOH/conc. NH₄OH) gave 68 mg (53 %) of **24h** as a brown-white solid. ¹H NMR (CDCl₃, 400 MHz) 0.82 (3H, t, *J* = 7.2 Hz), 1.05-2.12 (49H, m), 2.24 (3H, s), 2.26-2.97 (7H, m), 3.01, 3.10 (6H, s), 3.19-3.80 (6H, m), 3.95 (3H, m), 4.39 (3H, m), 4.89 (1H, d, *J* = 4.0 Hz), 5.04 (1H, d, *J* = 8.0 Hz), 7.35 (2H, d, *J* = 8.0 Hz), 7.76 (1H, s), 7.79 (2H, d, *J* = 8.0 Hz); ¹³C NMR (CDCl₃, 100MHz) δ 9.3, 10.8, 12.5, 16.1, 18.2, 18.8, 20.0, 21.2, 21.5, 21.7, 25.4, 26.2, 28.9, 29.0, 29.3, 29.6, 29.9, 30.2, 35.0, 37.0, 37.4, 39.3, 39.4, 45.2, 45.4, 49.6, 50.5, 50.8, 51.6, 57.8, 63.8, 65.8, 68.8, 69.2, 70.9, 72.7, 74.5, 76.8, 78.1, 78.5, 81.1, 96.1, 102.9, 119.7, 125.9, 129.6, 129.8, 138.9, 147.6, 176.1; HRMS (ESI) calc for [C₅₆H₉₃N₅O₁₅ + H]⁺ 1076.6740, found 1076.6667

HDAC Activity Assay

In Vitro HDAC inhibition was assayed using the HDAC Fluorimetric Assay/Drug Discovery Kit as previously described.^{17, 19} Briefly, 15 µl HeLa Nuclear extract was mixed with 5µl 10X compound and 5µl Assay Buffer. Fluorogenic substrate was added and reaction was allowed to proceed for 15 min at room temperature and then stopped by addition of a developer containing TSA. Fluorescence was monitored after 15 min at excitation and emission wavelengths of 360 and 460 nm, respectively. IC₅₀ values were determined using logit plots.

Cell Culture and Viability

SK-MES-1 and NCI-H69 lung cancer cell lines and DU-145 prostate cancer cell line were obtained from ATCC (Manassas, VA) and were maintained in the recommended complete growth mediums. MCF-7 cell line (a generous gift from Dr. Al Merrill) was maintained in EMEM containing 10% fetal bovine serum. Human mammary epithelial cells (HMEC) (Lonza Biosciences) were maintained in complete MEGM per product instructions. Human normal lung fibroblasts (a generous gift from Dr. Barker) were maintained in EMEM containing 10 percent FBS. All cell lines were maintained in a 37 °C environment containing 5 % CO₂. All compounds to be tested were dissolved to a concentration of 10 mM in DMSO and stored at -80 °C. For cell viability experiments, cells were passaged 24 h prior to dosing. All compounds were diluted to appropriate concentrations in DMSO and fresh medium such that the final concentration of DMSO was 0.1%. To control wells, only fresh medium was added. Viability was assessed after 72 h by Trypan Blue staining.

Western Blotting

SK-MES-1 cells were passaged 24 h prior to dosing with test compounds. Compounds were diluted to appropriate concentrations and were added to fresh medium so that the final concentration of DMSO was 0.1%. Cells were dosed with compounds for 8 h. Cells were washed with ice-cold PBS, recovered by centrifugation, and then lysed on ice in RIPA buffer containing protease inhibitors. Lysates were mixed repeatedly by pipetting and centrifuged at 14,000 rpm at 4 °C. Protein concentration was determined by Bradford assay (Bio-Rad). Protein denaturation and electrophoresis was performed as described.²⁷ Proteins were transferred to a nitrocellulose membrane for 1 h at 4 °C. The membrane was then blocked for 1 h in Odyssey blocking buffer (LI-COR Biosciences). Mouse anti-p21waf1 (Zymed) and rabbit anti-actin (Sigma) primary antibodies, as well as IR-labeled secondary antibodies (LI-COR Biosciences) were added as recommended (LI-COR Biosciences) and two-color detection used to image actin and p21waf1 by incubating with different host animal secondary antibodies. Detection was performed on an Odyssey Infrared Imaging System (LI-COR Biosciences). Blots were imaged simultaneously in both 700 nm and 800 nm channels at 169 µm resolution and analyzed on the imager software.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

We thank Professor Jim Powers for a helpful discussion on the manuscript. We are grateful to professors Al Merrill and Donald Doyle for the use of their cell culture facilities. We thank Professor Thomas Barker for a generous gift of primary lung fibroblasts. We are indebted to Chiaolong Hsiao for his assistance on Molecular Docking. This work was financially supported by Georgia Institute of Technology (AKO and YF), by the Blanchard fellowship to AKO and by Georgia Cancer Coalition Distinguished Cancer Clinicians and Scientists Program and NIH grant GM085261 (YF). PC and WG are recipients of the GAANN predoctoral fellowship from the Georgia Tech Center for Drug Design, Development and Delivery. RH is a Beckman Undergraduate Research Fellow.

References

1. Marks PA. Discovery and development of SAHA as an anticancer agent. *Oncogene* 2007;26:1351–1356. [PubMed: 17322921]
2. Kelly WK, O'Connor OA, Marks PA. Histone deacetylase inhibitors: from target to clinical trials. *Expert Opin Investig Expert Opin Investig Drugs* 2002;11:1695–1713.
3. Miller TA, Witter DJ, Belvedere S. Histone Deacetylase Inhibitors. *J Med Chem* 2003;46:5097–5116. [PubMed: 14613312]

4. Rosato RR, Grant S. Histone deacetylase inhibitors in clinical development. *Expert Opin Invest Drugs* 2004;13:21–38.
5. Monneret C. Histone deacetylase inhibitors. *European Journal of Medicinal Chemistry* 2005;40:1–13. [PubMed: 15642405]
6. Yoo CB, Jones PA. Epigenetic therapy of cancer: past, present and future. *Nature Reviews Drug Discovery* 2006;5:37–50.
7. Finnin MS, Donigian JR, Cohen A, Richon VM, Rifkind RA, Marks PA, Breslow R, Pavletich NP. Structures of a histone deacetylase homologue bound to the TSA and SAHA inhibitors. *Nature* 1999;401:188–193. [PubMed: 10490031]
8. (a) Vannini A, Volpari C, Filocamo G, Casavola EC, Brunetti M, Renzoni D, Chakravarty P, Paolini C, De Francesco R, Gallinari P, Steinkuhler C, Di Marco S. Crystal structure of a eukaryotic zinc-dependent histone deacetylase, human HDAC8, complexed with a hydroxamic acid inhibitor. *Proc Natl Acad Sci USA* 2004;101:15064–15069. [PubMed: 15477595] (b) Somoza JR, Skene RJ, Katz BA, Mol C, Ho JD, Jennings AJ, Luong C, Arvai A, Buggy JJ, Chi E, Tang J, Sang BC, Verner E, Wynands R, Leahy EM, Dougan DR, Snell G, Navre M, Knuth MW, Swanson RV, McRee DE, Tari LW. Structural Snapshots of Human HDAC8 Provide Insights into the Class I Histone Deacetylases. *Structure* 2004;12:1325–1334. [PubMed: 15242608] (c) Schuetz A, Min J, Allali-Hassani A, Schapira M, Shuen M, Loppnau P, Mazitschek R, Kwiatkowski NP, Lewis TA, Maglathin RL, McLean TH, Bochkarev A, Plotnikov AN, Vedadi M, Arrowsmith CH. Human HDAC7 Harbors a Class IIa Histone Deacetylase-specific Zinc Binding Motif and Cryptic Deacetylase Activity. *J Biol Chem* 2008;283:11355–11363. [PubMed: 18285338]
9. <http://www.clinicaltrials.gov/ct/show/NCT00106431>
10. (a) Furumai R, Komatsu Y, Nishino N, Khochbin S, Yoshida M, Horinouchi S. Potent histone deacetylase inhibitors built from trichostatin A and cyclic tetrapeptide antibiotics including trapoxin. *Proc Natl Acad Sci USA* 2001;98:87–92. [PubMed: 11134513] (b) Yurek-George A, Cecil ARL, Mo AHK, Wen S, Rogers H, Habens F, Maeda S, Yoshida M, Packham G, Ganesan A. The First Biologically Active Synthetic Analogues of FK228, the Depsipeptide Histone Deacetylase Inhibitor. *J Med Chem* 2007;50:5720–5726. [PubMed: 17958342] (c) Gomez-Paloma L, Bruno I, Cini E, Khochbin S, Rodriguez M, Taddei M, Terracciano S, Sadoul K. Design and Synthesis of Cyclopeptide Analogues of the Potent Histone Deacetylase Inhibitor FR235222. *ChemMedChem* 2007;2:1511–1519. [PubMed: 17694590]
11. Tsai WC, Standiford TJ. Immunomodulatory Effects of Macrolides in the Lung: Lessons from *In-Vitro* and *In-Vivo* Models. *Current Pharmaceutical Design* 2004;10:3081–3093. [PubMed: 15544499]
12. Shinkai M, Park CS, Rubin BK. Immunomodulatory Effects of Macrolide Antibiotics. *Interstitial, Inflammatory, and Occupational Lung Disease. Clinical Pulmonary Medicine* 2005;12:341–348.
13. Randolph JT, Waid P, Nichols C, Sauer D, Haviv F, Diaz G, Bammert G, Besecke LM, Segreti JA, Mohning KM, Bush EN, Wegner CD, Greer J. Nonpeptide Luteinizing Hormone-Releasing Hormone Antagonists Derived from Erythromycin A: Design, Synthesis, and Biological Activity of Cladinose Replacement Analogues. *J Med Chem* 2004;47:1085–1097. [PubMed: 14971889]
14. (a) Freiberg, LA. US Patent No. 3725385. 1973. (b) Lartey PA, Nellans HN, Faghieh R, Petersen A, Edwards CM, Freiberg L, Quigley S, Marsh K, Klein LL, Plattner JJ. Synthesis of 4"-Deoxy Motilides: Identification of a Potent and Orally Active Prokinetic Drug Candidate. *J Med Chem* 1995;38:1793–1798. [PubMed: 7752203] (c) Stenmark HG, Brazzale A, Ma Z. Biomimetic synthesis of macrolide/ketolide metabolites through a selective N-demethylation reaction. *J Org Chem* 2000;65:3875–3876. [PubMed: 10864780]
15. Awan A, Brennan RJ, Regan AC, Barber J. The conformations of the macrolide antibiotics erythromycin A, azithromycin and clarithromycin in aqueous solution: a 1H NMR study. *J Chem Soc Perkin Trans* 2000;2:1645–1652.
16. Elliott RL, Pireh D, Nilius AM, Johnson PM, Flamm RK, Chu DTW, Plattner JJ, Or YS. Novel 3-Deoxy-3-Descladinose-6-O-Methyl Erythromycin A Analogues. *Synthesis and In Vitro Activity. Bioorg Med Chem Lett* 1997;7:641–646.
17. HDAC Fluorimetric Assay/Drug Discovery Kit - AK-500 Manual. *Fluorescent Assay System* (BIOMOL® International, PA), 2005.

18. Morris, GM.; Goodsell, DS.; Halliday, RS.; Huey, R.; Hart, WE.; Belew, RK.; Olson, AJ. Automated Docking Using a Lamarckian Genetic Algorithm and Empirical Binding Free Energy Function; *J Comput Chem.* 1998. p. 1639-1662. <http://www.scripps.edu/pub/olson-web/doc/autodock/> (b) Wang DF, Wiest O, Helquist P, Lan-Hargest HY, Wiech NL. On the Function of the 14 Å Long Internal Cavity of Histone Deacetylase-Like Protein: Implications for the Design of Histone Deacetylase Inhibitors. *J Med Chem* 2004;47:3409–3417. [PubMed: 15189037] (c) Lu Q, Wang DS, Chen CS, Hu YD, Chen CS. Structure-Based Optimization of Phenylbutyrate-Derived Histone Deacetylase Inhibitors. *J Med Chem* 2005;48:5530–5535. [PubMed: 16107152]
19. Chen PC, Patil V, Guerrant W, Green P, Oyelere AK. Synthesis and Structure-Activity Relationship of Histone Deacetylase (HDAC) Inhibitors with Triazole-Linked Cap Group. *Bioorg Med Chem* 2008;16:4839–4853. [PubMed: 18397827]
20. Numerous examples of Cu (I) catalyzed Huisgen Cycloaddition reaction have appeared in the literature (A comprehensive list is available at www.scripps.edu/chem/sharpless/click.html). Cited here are two pioneering examples: (a) Rostovtsev VV, Green LG, Fokin VV, Sharpless KB. A Stepwise Huisgen Cycloaddition Process: Copper (I)-Catalyzed Regioselective “Ligation” of Azides and Terminal Alkynes. *Angew Chem Int Ed* 2002;41:2596–2599. (b) Tornøe CW, Christensen C, Meldal M. Peptidotriazoles on Solid Phase: [1,2,3]-Triazoles by Regiospecific Copper (I)-Catalyzed 1,3-Dipolar Cycloadditions of Terminal Alkynes to Azides. *J Org Chem* 2002;67:3057–3064. [PubMed: 11975567]
21. (a) Salisbury CM, Cravatt BF. Activity-based probes for proteomic profiling of histone deacetylase complexes. *Proc Natl Acad Sci USA* 2007;104:1171–1176. [PubMed: 17227860] (b) Salisbury CM, Cravatt BF. Optimization of activity-based probes for proteomic profiling of histone deacetylase complexes. *J Am Chem Soc* 2008;130:2184–2194. [PubMed: 18217751]
22. (a) KrennHrubic K, Marshall BL, Hedglin M, Verdin E, Ulrich SM. Design and evaluation of ‘Linkerless’ hydroxamic acids as selective HDAC8 inhibitors. *Bioorg Med Chem Lett* 2007;17:2874–2878. [PubMed: 17346959] (b) Gurard-Levin ZA, Mrksich M. The Activity of HDAC8 Depends on Local and Distal Sequences of Its Peptide Substrates. *Biochemistry* 2008;47:6242–6250. [PubMed: 18470998] (c) Haggarty SJ, Koeller KM, Wong JC, Grozinger CM, Schreiber SL. Domain-selective small-molecule inhibitor of histone deacetylase 6 (HDAC6)-mediated tubulin deacetylation. *Proc Natl Acad Sci USA* 2003;100:4389–4394. [PubMed: 12677000]
23. Freshney, R. *Culture of Animal Cells: A Manual of Basic Technique*. Alan R Liss, Inc.; New York: 1987. p. 117
24. Suzuki T, Nagano Y, Kouketsu A, Matsuura A, Maruyama S, Kurotaki M, Nakagawa H, Miyata N. Novel Inhibitors of Human Histone Deacetylases: Design, Synthesis, Enzyme Inhibition, and Cancer Cell Growth Inhibition of SAHA-Based Non-hydroxamates. *J Med Chem* 2005;48:1019–1032. [PubMed: 15715470]
25. Rundlett SE, Carmen AA, Kobayashi R, Bavykin S, Turner BM, Grunstein M. HDA1 and RPD3 are members of functionally distinct yeast histone deacetylase complexes. *Proc Natl Acad Sci USA* 1996;93:14503–14508. [PubMed: 8962081]
26. Marks P, Rifkind RA, Richon VM, Breslow R, Miller T, Kelly WK. Histone deacetylases and cancer: causes and therapies. *Nat Rev Cancer* 2001;1:194–202. [PubMed: 11902574]
27. Richon VM, Sandhoff TW, Rifkind RA, Marks PA. Histone deacetylase inhibitor selectively induces p21^{WAF1} expression and gene-associated histone acetylation. *Proc Natl Acad Sci USA* 2000;97:10014–10019. [PubMed: 10954755]
28. Danesi R, Lupetti A, Barbara C, Ghelardi E, Chella A, Malizia T, Senesi S, Angeletti CA, Del Tacca M, Campa M. Comparative distribution of azithromycin in lung tissue of patients given oral daily doses of 500 and 1000 mg. *J Antimicrob Chemother* 2003;51:939–945. [PubMed: 12654753]
29. Savariar EN, Aathimanikandan SV, Thayumanavan S. Supramolecular assemblies from amphiphilic homopolymers: Testing the scope. *J Am Chem Soc* 2006;128:16224–16230. [PubMed: 17165775]
30. Heine HW, Becker EB, Lane JF. On Cyclic Intermediates in Substitution Reactions. III. The Alkaline Hydrolysis of ε-Bromocaproic and ζ-Bromoanthranic Acids. *J Am Chem Soc* 1953;75:4514–4515.
31. (a) Collman JP, Devaraj NK, Chidsey CED. ‘Clicking’ Functionality onto Electrode Surfaces. *Langmuir* 2004;20:1051–1053. [PubMed: 15803676] (b) Shon YS, Kelly KF, Halas NJ, Lee TR. Fullerene-Terminated Alkanethiolate SAMs on Gold Generated from Unsymmetrical Disulfides. *Langmuir* 1999;15:5329–5332.

32. Muri D, Bode JW, Carreira EM, Novel A. General Method for the Synthesis of Nitrile Oxides: Dehydration of *O*-Silylated Hydroxamic Acids. *Org Lett* 2000;2:539–541. [PubMed: 10814371]

Abbreviations

HDAC	Histone deacetylase
HDACi	Histone deacetylase inhibitors
HDLP	Histone deacetylase-like protein
SAHA	Suberoylanilide hydroxamic acid
TSA	trichostatin A
ZBG	Zinc-binding group

Figure 1.
Synthesis of prototypical nonpeptide macrocyclic HDACi **8** and **10**.

Figure 2.

SAR studies on nonpeptide macrocyclic HDACi (a) Synthesis of triazole-linked nonpeptide macrocyclic HDACi based on azithromycin 15-membered ring; (b) Synthesis of triazole-linked nonpeptide macrocyclic HDACi based on clarithromycin 14-membered ring.

Figure 3.

Docked structures of SAHA, 14- and 15-membered macrocyclic HDACi at the active site of HDLP. (a) Super-position of the low energy conformation of **16b** (blue) and SAHA (yellow) revealed the pocket binding preferences of inhibitors at the HDLP surface. Ball and stick model of the orientation of the hydroxamate group of SAHA and **16b** (b); **16b** and **16d** (orange) (c) with respect to the active site Zn²⁺ ion (grey ball). (d) Relative orientation of the macrocyclic rings of **16d** and **24d** (yellow) with respect to Phe 338 at the HDLP surface.

Figure 4.

Western blot detection of p21^{WAF1/CIP1} expression levels in SK-MES-1 cells exposed to SAHA and **24d**. Cells are exposed to test agents for 8 hours. Compound numbers and concentrations are displayed in the top panel. Immunoblotting with anti-actin Antibody (in green) is used as a control for equivalent protein loading.

Scheme 1.

(a) Selected examples of acyclic HDAC inhibitors; (b) Representative examples of Cyclic-peptide HDAC inhibitors; (c) Representative examples of Macrolide Antibiotics.

Table 1

In vitro HDAC inhibition (IC₅₀) and isoform selectivity of nonpeptide macrocyclic HDACi. IC₅₀ values were determined using a cell free kit assay.¹⁷ Each data is obtained from three independent experiments.

Compound	HDAC 1/2 (nM)	HDAC 8 (nM)	Isoform Selectivity ^a
7	N.D.	N.T.	– ^b
8	107.1	6,680	62
10	109.8	2,320	21
16a	91.6	4,730	51
16b	88.8	3,740	42
16c	13.9	994	72
16d	10.6	1,020	97
16e	58.9	7,130	121
16f	72.4	6,780	94
16g	145.5	11,050	38
16h	226.7	N.D.	– ^b
24a	37.0	3,990	108
24b	44.3	4,750	107
24c	4.1	1,890	462
24d	1.9	1,390	743
24e	55.6	5,880	106
24f	123.0	4,420	36
24g	169.8	10,550	56
24h	223.4	N.D.	– ^b
SAHA	65.0	1,860	29

N.D. – not determinable N.T. – not tested

^a calculated by dividing the IC₅₀ of HDAC 8 with the IC₅₀ of HDAC 1/2

^b undeterminable

Table 2

Cell growth inhibition data for nonpeptide macrocyclic HDACi. EC₅₀ values were determined from trypan blue exclusion data. Each data is obtained from a duplicate of four simultaneous experiments.

Compound	SKMES 1 (μM)	NCL-H69 (μM)	DU-145 (μM)	lung fibroblast (μM)	HMEC (μM)
8	N.T.	N.T.	>25	N.T.	N.T.
10	N.T.	N.T.	<25	N.T.	N.T.
16a	1.79	1.92	1.45	>10	>10
16b	1.68	1.77	1.24	>10	>10
16c	2.33	3.45	1.88	>10	>10
16d	2.56	3.01	1.97	>10	>10
16e	4.89	4.56	5.89	>10	>10
16f	4.67	3.99	5.68	>10	>10
16g	7.54	8.45	>10	>10	>10
24a	2.15	2.67	2.98	>10	>10
24b	1.95	1.92	3.29	>10	>10
24c	1.33	1.45	1.12	>10	>10
24d	1.28	1.49	1.05	>10	>10
24e	4.89	5.67	6.97	>10	>10
24f	4.45	5.09	5.78	>10	>10
24g	7.12	7.29	8.14	>10	>10
SAHA	2.42	2.06	2.12	>10	>10

N.T. – not tested