Published in final edited form as:

Psychol Med. 2009 March ; 39(3): 355–363. doi:10.1017/S0033291708003930.

HIV Testing among Individuals with a Severe Mental Illness: Review, Suggestions for Research, and Clinical Implications

Theresa E. Senn¹ and Michael P. Carey

Center for Health and Behavior, Syracuse University

Abstract

Background—The prevalence of HIV is elevated among individuals with a severe mental illness (SMI). Because of the benefits of HIV testing, it is important for individuals with SMI to have routine access to testing.

Objectives—The goals of this review are to summarize knowledge about HIV testing prevalence, correlates, and interventions among individuals with a SMI; to identify research needs; and to discuss clinical implications of the studies reviewed.

Methods—Literature searches were conducted using PsycINFO, PubMed, and MEDLINE. Additional articles were obtained from reference lists of relevant articles.

Results—Fewer than one-half of individuals with a SMI have been tested for HIV in the past year. Engaging in sex or drug risk behavior was the only consistent correlate of HIV testing. Interventions for promoting HIV testing among individuals with a SMI have not been well-developed or evaluated.

Conclusions—Research on HIV testing among individuals with a SMI is needed. Mental health settings may be opportune venues for HIV testing, even though providers face ethical challenges when implementing testing programs in these settings.

Introduction

The prevalence of HIV infection is believed to be elevated among individuals with a severe mental illness (SMI; McKinnon et al., 2002); thus, it is important for individuals with a SMI be tested for HIV, which confers both individual and public health benefits. Individuals who test positive will benefit from medical care; the public benefits because antiretroviral therapy reduces infectiousness and knowledge of infection status motivates risk reduction (Weinhardt et al., 1999, Marks et al., 2005). In this article, we summarize knowledge regarding sexual risk behavior and HIV prevalence among individuals with a SMI, review the prevalence and correlates of HIV testing in this population, suggest directions for research on testing, and discuss the clinical implications of the research reviewed.

Sexual Risk Behavior and HIV Prevalence Among Individuals with a SMI

The majority of adults with a SMI are sexually active, and those that are sexually active engage in high rates of sexual risk behavior. Reviews of the literature have found that 54% to 74% of adults with a SMI were sexually active in the last year (Carey et al., 1997, Meade and Sikkema, 2005a). Among sexually active patients, 43% had multiple partners and 46% had not used a condom in the past year; lifetime sex trading was reported by 22%, and 33% reported a history

¹ Corresponding Author: Theresa E. Senn, Center for Health and Behavior, 430 Huntington Hall, Syracuse University, Syracuse, NY 13244; email: tsenn@syr.edu.

of STDs, rates much higher than those found in the general population (Meade and Sikkema, 2005a).

There are many possible causes of the high rates of sexual risk behavior among individuals with a SMI. Sexual risk behavior has been related to: (a) psychiatric symptoms or diagnosis (e.g., Carey et al., 2004, Meade and Sikkema, 2007); (b) factors associated with SMI, such as lack of planning, inaccurate assessment of risk, and poor communication skills (e.g., Gordon et al., 1999); (c) lack of information (e.g., Strauss et al., 2006); (d) lack of motivation to engage in safer sexual behaviors (e.g., Otto-Salaj et al., 1998); and (e) lack of safer sex skills (e.g., Kalichman et al., 2005, Meade and Sikkema, 2005a). In addition, (f) the consequences of mental illness (e.g., poverty, homelessness) may lead to insufficient money to purchase condoms and no privacy for safer sex negotiation (Carey et al., 1997, McKinnon et al., 2002). SMI also leads to periods of hospitalization, which make it difficult to form long-term sexual relationships, and to acquire condoms while hospitalized (McKinnon et al., 2002), and SMI may lead to a lack of social support, which has been associated with sexual risk behavior among individuals with a SMI (e.g., Randolph et al., 2007). Finally, variables that are associated with both SMI and sexual risk behavior, such as (g) childhood sexual abuse (Meade and Sikkema, 2007) and (h) substance use (e.g., Weinhardt et al., 2002, Carey et al., 2004, Meade and Weiss, 2007), may lead to increased sexual risk behavior.

The prevalence of HIV is higher among individuals with a SMI than among individuals in the general population, with rates ranging from 4% to 23% (Cournos and McKinnon, 1997). These rates are much higher than the 0.43% HIV prevalence rate found in the general population (McQuillan et al., 2006). The high rates of HIV in individuals with SMI may be, in part, driven by individuals who are dually diagnosed with a substance use disorder (Himelhoch et al., 2007; Parry et al., 2007). Among the dually diagnosed, HIV prevalence ranged from 6% to 23% (Meade and Weiss, 2007).

Methods

To obtain articles for this review, searches were conducted with PsycINFO, PubMed, and MEDLINE (using the keywords “mental* ill*” and “HIV test*”) and by perusing reference lists of published articles. To be included in the review, studies had to: (a) be published in a peer-reviewed journal; (b) include individuals with a SMI; and (c) report on HIV testing (i.e., prevalence, correlates, interventions). All articles published through 2007 that met inclusion criteria were included in the review ($n = 18$).

Results

HIV Testing Rates

Studies investigating HIV testing rates among individuals with a SMI indicate that the percentage of individuals who have been tested in the past year ranged from 17% to 47% (see Table 1); lifetime prevalence of testing ranged from 11% to 89%. Pirl et al. (2005) reported that 10% of patients in a state psychiatric hospital were tested during a single hospitalization.

In one study that compared HIV testing rates among individuals with and without SMI, adults with major depression, bipolar disorder, or schizoaffective disorder were more likely than individuals without a disorder to have been ever tested for HIV (46% vs. 31%), to have been tested for HIV in the past 12 months (18% vs. 10%), and to expect to be tested for HIV in the next 12 months (14% vs. 7%) (Blumberg and Dickey, 2003). However, in another study, women with schizophrenia or schizoaffective disorders were less likely than women without a SMI to ever be tested for HIV (11% vs. 67%; Miller and Finnerty, 1996). Most adults with a SMI who report ever being tested for HIV were tested for HIV more than once, with 30%

reporting two lifetime tests, 21% reporting three tests, and 35% reporting $\geq$ four HIV tests (Meade and Sikkema, 2005b).

HIV Testing Correlates

Several studies have investigated HIV testing correlates. Among various sociodemographic characteristics, only homelessness has been consistently associated with being tested for HIV (Desai and Rosenheck, 2004, Meade and Sikkema, 2005b). Characteristics such as age, education, and race/ethnicity have not been consistently associated with testing uptake, although these findings are based on only a few studies (Desai et al., 2007, Desai and Rosenheck, 2004, Goldberg, 2004, Meade and Sikkema, 2005b, Walkup et al., 2000).

The relation between testing and psychiatric symptoms also has been investigated. Meade and Sikkema (2005b) found that being tested for HIV was associated with having a non-psychotic disorder as the primary diagnosis and having borderline personality disorder. In an Australian sample, Thompson et al. (1997) found that individuals diagnosed with bipolar disorder were less likely to be tested than individuals diagnosed with schizophrenia. Some studies have found that more psychiatric distress or more severe symptoms and more psychiatric hospitalizations were associated with testing (e.g., Thompson et al., 1997, Desai and Rosenheck, 2004), but Desai et al. (2007) found that fewer psychiatric symptoms was associated with testing, and Walkup et al. (2000) found that being voluntarily admitted to inpatient hospitalization, a proxy for less severe symptoms, was associated with being tested.

The number and types of services used are also associated with testing. Receiving services at an urban (vs. suburban) mental health center (Goldberg et al., 2005), using a greater number of treatment services (Meade and Sikkema, 2005b), and recently using health services (Desai and Rosenheck, 2004) were associated with being tested.

More social support is associated with being tested for HIV (Meade and Sikkema, 2005b); being in an intimate relationship, which could be an indication of greater social support, also was associated with HIV testing (Walkup et al., 2000). In addition, having a stronger therapeutic alliance with the primary clinician (construed here as a type of social support) has also been associated with HIV testing among individuals with a SMI (Desai et al., 2007). Motivational variables related to testing include having more concerns about contracting HIV and knowing someone who has AIDS or who died of AIDS (Knox et al., 1994, Desai and Rosenheck, 2004).

Individuals who engage in risk behaviors are more likely to be tested for HIV. Those with drug problems or a history of injection drug use (Thompson et al., 1997, Walkup et al., 2000, Desai et al., 2007) were more likely to be tested for HIV, although there appears to be no relation between alcohol problems and HIV testing (Walkup et al., 2000). More sexual risk behavior is also associated with HIV testing among SMI, with those who had unprotected sex, engaged in recent sexual risk behavior, exchanged sex, or ever been diagnosed with an STD more likely to be tested for HIV (Knox et al., 1994, Thompson et al., 1997, Weinhardt et al., 1998, Goldberg et al., 2005, Desai et al., 2007). Meade and Sikkema (2005b) found that individuals who had engaged in any sexual or drug use risk behavior were more likely to have been tested (55% vs. 24%).

Reasons for Testing (and Declining Testing)

Research suggests several reasons why individuals get tested. In one study, adults with a SMI who were tested were more likely than adults without SMI to report they simply wanted to know their HIV status; in contrast, adults without SMI were more likely to get tested because of insurance, employment, military, immigration, pregnancy, or hospitalization (Blumberg and

Dickey, 2003). An Australian study found that individuals with a SMI reported being tested for HIV because they requested it, because they were raped, or for a medical reason (e.g., pregnancy, illness, overdose, blood borne infection; Thompson et al., 1997).

To our knowledge, there has not been investigation of the reasons that people with a SMI choose not to be tested. Among the general population, people decline testing for many reasons; for example, some have been tested previously whereas others believe they are not at risk for HIV (Carey et al., in press). Sometimes people fear the results (Simon et al., 1996, CDC, 2003a); others may be reluctant to be tested because of misgivings about provider sensitivity or subsequent stigma (Mimiaga et al., 2007). Research is needed to determine if such reasons are barriers to people living with a SMI.

Returning for Test Results

Among the general population, test return rates are poor; in contrast, 89% to 96% of people with a SMI return for their results (Desai and Rosenheck, 2004, Desai et al., 2007); however, only 50% of adults with a SMI reported receiving post-test counseling (Desai et al., 2007). Returning for results was associated with more education, not having a disability, having been tested previously, not having an STD, having fewer drug problems, and having previously received HIV test results (Desai and Rosenheck, 2004).

Interventions to Promote Testing among Individuals with a SMI

Few interventions have been designed to promote testing among individuals with a SMI. In the most well-described intervention to date, Rosenberg et al. (2004) developed the **Screen, Test, Immunize, Reduce risk, and Refer (STIRR)** intervention. With this program, infectious disease specialists traveled to mental health centers and met with administration and staff to encourage support of the project. Next, a nurse provided HIV and hepatitis testing, pre- and post-test counseling, hepatitis immunization, and referrals for individuals who were HIV positive. At one mental health center, 136 out of 173 (79%) patients were tested for HIV. At a second mental health center, 65 out of 99 (66%) dually diagnosed patients were tested (Rosenberg et al., 2004). Although there was no comparison group, the percentage of patients in the program who were tested for HIV was higher than the 17% to 47% of patients who reported recently being tested for HIV in other studies (Walkup et al., 2000, Blumberg and Dickey, 2003, Desai and Rosenheck, 2004, Meade and Sikkema, 2005b). Meyer (2003) reported that 78% of patients in a state psychiatric hospital were tested for HIV after initiation of a voluntary hepatitis and HIV screening program, but the program was neither well-described nor compared to a control condition.

Conclusions

Studies reported that 17% to 47% of individuals with a SMI were recently tested for HIV, and that 11% to 89% were ever tested for HIV. The variability in estimates reflects variability in many study features including sample size, dates of investigation, sampling (e.g., inpatients vs. outpatients), and policies at the centers from which participants were recruited. Despite these methodological variations, the range of findings indicates that routine HIV testing is not yet a standard practice in most mental health settings.

Studies investigating correlates of testing among individuals with a SMI have reported mixed findings. There was no clear association between HIV testing and either demographic or psychiatric variables; however, many of these findings were based on only one or two studies. The lack of an association between psychiatric symptoms and HIV testing was unanticipated; we expected that individuals with greater distress would be less able to provide informed consent, and would be less likely to be tested for HIV. In a vignette study, clinicians were

reluctant to test for HIV if a patient was impaired, even if the patient had a history of risk behavior (Walkup et al., 2002). The association between testing and psychiatric symptoms or level of functioning merits investigation; it may be important to tailor HIV testing interventions to different levels of functioning.

Engaging in more HIV risk behavior was associated with HIV testing. Thus, among individuals with a SMI, those who are most in need of an HIV test are being tested. Individuals who engage in HIV-risk behaviors may be more concerned about their HIV risk, and therefore seek testing. Clinicians most likely refer patients who report HIV risk behaviors for HIV testing. Although only one study investigated the clinician/patient relationship and HIV testing, a stronger therapeutic alliance was associated with a greater likelihood of testing (Desai et al., 2007), which suggests that therapists could be a valuable resource to encourage their patients to get tested for HIV.

We located only one well-described intervention to encourage HIV testing among individuals with a SMI (viz., the STIRR program; Rosenberg et al., 2004). Testing rates for individuals at mental health centers that participated in the intervention were higher than testing rates in other studies; however, because the study employed a post-test only design, firm conclusions cannot be drawn about the efficacy of the intervention.

There are several limitations to this review. First, most of the findings presented in this article were based on only a few studies. More research is needed before firm conclusions about HIV testing rates and correlates can be drawn. Second, because there are few studies, we could not explore the impact of characteristics of study samples or methodology on findings. Finally, the majority of studies reported in this article were conducted in the US; findings may differ in other countries, where HIV rates, SMI or HIV stigma, and access to HIV testing may differ.

Research Needs

HIV testing among individuals with a SMI deserves further study. First, studies investigating the prevalence of HIV testing among individuals with a SMI should determine the overall prevalence of HIV testing as well as the prevalence among individuals engaging in risk behavior (e.g., Meade & Sikkema, 2005b). Second, it is important to know the timing of HIV diagnosis among individuals with a SMI (Cournos and McKinnon, 1997). Research has found that a large percentage of individuals are diagnosed late in the course of HIV infection (CDC, 2003b). Early diagnosis of HIV allows for earlier entry into care, and facilitates risk reduction. It is important to know if individuals with a SMI are diagnosed later than other individuals. Third, research might explore individual (e.g., low risk awareness) and structural (e.g., logistic, policy) barriers to testing, in order to inform intervention development.

Fourth, studies to encourage HIV testing among individuals with a SMI who engage in HIV-risk behavior are needed. To date, no randomized controlled trials (RCTs) to increase HIV testing rates among individuals with a SMI have been conducted. Interventions focusing on service barriers could evaluate the STIRR intervention in a RCT. Alternatively, if attitudinal barriers were targeted, HIV testing interventions that have been efficacious with individuals without a SMI could be adapted and implemented with individuals with a SMI (e.g., Carey et al., in press).

Finally, the impact of HIV diagnosis on an individual with a SMI merits study. One reason for trying to increase HIV testing uptake is that individuals infected with HIV reduce their sexual risk behavior (Weinhardt et al., 1999, Marks et al., 2005). Given the cognitive and social impairments characteristic of SMI (e.g., social skill deficits), research might determine whether HIV diagnosis in individuals with a SMI results in changes in sexual risk behavior. If not, further intervention may be needed to provide individuals with a SMI who are HIV positive

with the information, motivation, skills, and other support they need to adopt safer sexual practices.

Clinical Implications

To promote the overall health of adults with a SMI, it is important that individuals with a SMI who engage in HIV risk behaviors are tested for HIV. Psychiatric settings may be an opportune venue for HIV testing, for several reasons. First, this is where patients can be found, reducing the need for additional transportation or appointments. Second, mental health clinicians can provide pre- and post-test counseling that is tailored for individuals with a SMI, which other HIV testing sites may be less able to provide (Satriano et al., 2007). Third, therapists can encourage HIV testing, address patients' barriers to testing, and allay concerns about testing, confidentiality, and stigma.

Unfortunately, many mental health settings do not encourage HIV testing. In a survey of psychiatric units in hospitals, 53% reported that they encouraged only a few patients to get tested, and 17% reported encouraging almost no patients to get tested (Walkup et al., 1998). Among outpatient mental health centers in New York state, only 22% provided pre- and post-test counseling, only 16% offered testing on-site, and 13% did not have a procedure to refer patients to other sites for testing (Satriano et al., 2007).

Although providers usually agree that HIV-related services are important (Satriano et al., 2007), there are many barriers to providing HIV services, including lack of training, discomfort with the topic, and competing priorities (Solomon et al., 2007). Clinicians may be uncomfortable delivering risk reduction counseling, or obtaining consent for or administering HIV tests. Patients may be uncomfortable talking about HIV as well. Additionally, because of resource constraints, HIV testing and counseling in mental settings might take time away from the provision of mental health services.

To reduce barriers to HIV testing in mental health settings, therapists should be provided with training around HIV testing and prevention. Fortunately, the CDC supports technical assistance and training programs. In addition, recent advances have led to the development of rapid HIV testing, which is easy to administer, does not require a venipuncture, can be administered by non-medical personnel, and provides results within 20 minutes, eliminating the need to return for test results.

There are several ethical and legal challenges associated with HIV testing of individuals with a SMI. First, there may be limited benefits if individuals learn they are HIV positive, because some doctors delay the prescription of antiretroviral medication for individuals with a SMI (Fairfield et al., 1999; Bogart et al., 2000). This delay in prescribing medication may be due to concerns about interactions among medications, or concerns about HIV-medication adherence (Fairfield et al., 1999).

Second, even if HIV-infected individuals are prescribed medication, they may not benefit, because individuals with a mental illness often have difficulty adhering to antiretroviral medication (e.g., Tucker et al., 2003). Needed are interventions to increase adherence to antiretroviral medications (Simoni et al., 2003; Uldall et al., 2004).

Third, some individuals, because of their mental illness, may not be able to provide informed consent for HIV testing. The laws requiring informed consent for HIV testing for individuals with mental disabilities differ from state to state, and often the need for testing without consent is decided on an individual basis (Haimowitz, 1996).

Fourth, the ethical and legal issues associated with testing are complex. Clinicians must balance confidentiality with the duty to warn (particularly if patients are infected and sexually active in an inpatient setting; see Odunsi, 2007). Such issues require knowledge of state law, and use of clinical judgment (Haimowitz, 1996).

In conclusion, individuals with a SMI are at elevated risk for HIV but many have never been tested. Research on testing should include individuals with a SMI, and interventions to promote testing should be evaluated in RCTs. Mental health settings may be opportune venues to test individuals with a SMI for HIV even though clinicians in these settings will face a number of barriers and ethical challenges. These challenges indicate the need for research, continuing professional education, and the commitment of additional resources to this neglected component of comprehensive health care.

Acknowledgements

This research was supported by NIH grant # R01MH068171 to Michael P. Carey.

References

- Blumberg SJ, Dickey WC. Prevalence of HIV risk behaviors, risk perceptions, and testing among US adults with mental disorders. *Journal of Acquired Immune Deficiency Syndromes* 2003;32:77–79. [PubMed: 12514417]
- Bogart LM, Kelly JA, Catz SL, Sosman JM. Impact of medical and nonmedical factors on physician decision making for HIV/AIDS antiretroviral treatment. *Journal of Acquired Immune Deficiency Syndromes* 2000;23:396–404. [PubMed: 10866232]
- Carey MP, Carey KB, Kalichman SC. Risk for human immunodeficiency virus (HIV) infection among persons with severe mental illness. *Clinical Psychology Review* 1997;17:271–291. [PubMed: 9160177]
- Carey MP, Carey KB, Maisto SA, Schroder KEE, Vanable PA, Gordon CM. HIV risk behavior among psychiatric outpatients: Association with psychiatric disorder, substance use disorder, and gender. *Journal of Nervous and Mental Disease* 2004;192:289–296. [PubMed: 15060403]
- Carey MP, Coury-Doniger P, Senn TE, Vanable PA, Urban MA. Improving HIV rapid testing rates among STD clinic patients: A randomized, controlled trial. *Health Psychology*. in press
- Centers for Disease Control and Prevention. HIV/AIDS special surveillance report: HIV testing survey, 2000. US Department of Health and Human Services; Atlanta: 2003a.
- Centers for Disease Control and Prevention. Late versus early testing of HIV--16 sites, United States, 2000–2003. *Morbidity and Mortality Weekly Report* 2003b;52:581–586.
- Cournos F, McKinnon K. HIV seroprevalence among people with severe mental illness in the United States: A critical review. *Clinical Psychology Review* 1997;17:259–269. [PubMed: 9160176]
- Desai MM, Rosenheck RA. HIV testing and receipt of test results among homeless persons with serious mental illness. *American Journal of Psychiatry* 2004;161:2287–2294. [PubMed: 15569902]
- Desai MM, Rosenheck RA, Desai RA. Prevalence and correlates of human immunodeficiency virus testing and posttest counseling among outpatients with serious mental illness. *Journal of Nervous and Mental Disease* 2007;195:776–780. [PubMed: 17984780]
- Fairfield KM, Libman H, Davis RB, Eisenberg DM, Phillips RS. Delays in protease inhibitor use in clinical practice. *Journal of General Internal Medicine* 1999;14:395–401. [PubMed: 10417596]
- Goldberg RW. Letters to the Editor: Hepatitis and HIV screening, education, and treatment for adults with serious mental illness. *General Hospital Psychiatry* 2004;26:167–168. [PubMed: 15038938]
- Goldberg RW, Himelhoch S, Kreyenbuhl J, Dickerson FB, Hackman A, Fang LJ, Brown CH, Wohlheiter KA, Dixon LB. Predictors of HIV and hepatitis testing and related service utilization among individuals with serious mental illness. *Psychosomatics* 2005;46:573–577. [PubMed: 16288138]
- Gordon CM, Carey MP, Carey KB, Maisto SA, Weinhardt LS. Understanding HIV-related risk among persons with a severe and persistent mental illness: Insights from qualitative inquiry. *Journal of Nervous and Mental Disease* 1999;187:208–216. [PubMed: 10221553]

- Grassi L, Pavanati M, Cardelli R, Ferri S, Peron L. HIV-risk behaviour and knowledge about HIV/AIDS among patients with schizophrenia. *Psychological Medicine* 1999;29:171–179. [PubMed: 10077305]
- Haimowitz, S. An overview of legal issues. In: Cournos, F.; Bakalar, N., editors. *AIDS and People with Severe Mental Illness*. Yale University Press; New Haven: 1996. p. 227-239.
- Himelhoch S, McCarthy JF, Ganoczy D, Medoff D, Dixon LB, Blow FC. Understanding associations between serious mental illness and HIV among patients in the VA health system. *Psychiatric Services* 2007;58:1165–1172. [PubMed: 17766561]
- Kalichman S, Malow R, Devieux J, Stein JA, Piedman F. HIV risk reduction for substance using seriously mentally ill adults: Test of the Information-Motivation-Behavior Skills (IMB) model. *Community Mental Health Journal* 2005;41:277–290. [PubMed: 16131007]
- Katz RC, Watts C, Santman J. AIDS knowledge and high risk behaviors in the chronic mentally ill. *Community Mental Health Journal* 1994;30:395–401. [PubMed: 7956114]
- Kelly JA, Murphy DA, Bahr GR, Brasfield TL, Davis DR, Hauth AC, Morgan MG, Stevenson LY, Eilers MK. AIDS/HIV risk behavior among the chronic mentally ill. *American Journal of Psychiatry* 1992;149:886–889. [PubMed: 1609866]
- Knox MD, Boaz TL, Friedrich MA, Dow MG. HIV risk factors for persons with serious mental illness. *Community Mental Health Journal* 1994;30:551–563. [PubMed: 7835041]
- Levounis P, Galanter M, Dermatis H, Hamowy A, De Leon G. Correlates of HIV transmission risk factors and considerations for interventions in homeless, chemically addicted and mentally ill patients. *Journal of Addictive Diseases* 2002;21:61–72. [PubMed: 12095000]
- Marks G, Crepaz N, Senterfitt W, Janssen RS. Meta-analysis of high-risk sexual behavior in persons aware and unaware they are infected with HIV in the United States. *Journal of Acquired Immune Deficiency Syndromes* 2005;39:446–453. [PubMed: 16010168]
- McKinnon K, Cournos F, Herman R. HIV among people with chronic mental illness. *Psychiatric Quarterly* 2002;73:17–31. [PubMed: 11780595]
- McQuillan GM, Kruszon-Moran D, Kottiri BJ, Kamimoto LA, Lam L, Cowart MF, Hubbard M, Spira TJ. Prevalence of HIV in the US household population: The National Health and Nutrition Examination Surveys, 1988 to 2002. *Journal of Acquired Immune Deficiency Syndromes* 2006;41:651–656. [PubMed: 16652040]
- Meade CS, Sikkema KJ. HIV risk behavior among adults with severe mental illness: A systematic review. *Clinical Psychology Review* 2005a;25:433–457. [PubMed: 15914265]
- Meade CS, Sikkema KJ. Voluntary HIV testing among adults with severe mental illness: Frequency and associated factors. *AIDS and Behavior* 2005b;9:465–473. [PubMed: 16323033]
- Meade CS, Sikkema KJ. Psychiatric and psychosocial correlates of sexual risk behavior among adults with severe mental illness. *Community Mental Health Journal* 2007;43:153–169. [PubMed: 17143728]
- Meade CS, Weiss RD. Substance abuse as a risk factor for HIV sexual risk behavior among persons with severe mental illness: Review of evidence and exploration of mechanisms. *Clinical Psychology: Science and Practice* 2007;14:23–33.
- Meyer JM. Prevalence of hepatitis A, hepatitis B, and HIV among hepatitis C-seropositive state hospital patients: Results from Oregon State Hospital. *Journal of Clinical Psychiatry* 2003;64:540–545. [PubMed: 12755656]
- Miller LJ, Finnerty M. Sexuality, pregnancy, and childrearing among women with schizophrenia-spectrum disorders. *Psychiatric Services* 1996;47:502–506. [PubMed: 8740491]
- Mimiaga MJ, Goldhammer H, Belanoff C, Tetu AM, Mayer KH. Men who have sex with men: Perceptions about sexual risk, HIV and sexually transmitted disease testing, and provider communication. *Sexually Transmitted Diseases* 2007;34:113–119. [PubMed: 16810121]
- Odunsi B. Should caregivers be compelled to disclose patients' HIV infection to the patients' sex partners without consent? *Studies in Family Planning* 2007;38:297–306. [PubMed: 18284044]
- Otto-Salaj LL, Heckman TG, Stevenson LY, Kelly JA. Patterns, predictors, and gender differences in HIV risk among severely mentally ill men and women. *Community Mental Health Journal* 1998;34:175–190. [PubMed: 9620162]
- Parry CD, Blank MB, Pithey AL. Responding to the threat of HIV among persons with mental illness and substance abuse. *Current Opinion in Psychiatry* 2007;20:235–241. [PubMed: 17415075]

- Pirl WF, Greer JA, Weissgarber C, Liverant G, Safren SA. Screening for infectious diseases among patients in a state psychiatric hospital. *Psychiatric Services* 2005;56:1614–1616. [PubMed: 16339630]
- Randolph ME, Pinkerton SD, Somlai AM, Kelly JA, McAuliffe TL, Gibson RH, Hackl K. Severely mentally ill women's HIV risk: The influence of social support, substance use, and contextual risk factors. *Community Mental Health Journal* 2007;43:33–47. [PubMed: 17143730]
- Rosenberg S, Brunette M, Oxman T, Marsh B, Dietrich A, Mueser K, Drake R, Torrey W, Vidaver R. The STIRR model of best practices for blood-borne diseases among clients with serious mental illness. *Psychiatric Services* 2004;55:660–664. [PubMed: 15175463]
- Satriano J, McKinnon K, Adoff S. HIV service provision for people with severe mental illness in outpatient mental health care settings in New York. *Journal of Prevention and Intervention in the Community* 2007;33:95–108. [PubMed: 17298933]
- Simon PA, Weber M, Ford WL, Cheng F, Kerndt PR. Reasons for HIV antibody test refusal in a heterosexual sexually transmitted disease clinic population. *AIDS* 1996;10:1549–1553. [PubMed: 8931791]
- Simoni JM, Frick PA, Pantalone DW, Turner BJ. Antiretroviral adherence interventions: A review of current literature and ongoing studies. *Topics in HIV Medicine* 2003;11:185–198. [PubMed: 14724327]
- Solomon PL, Tennille JA, Lipsitt D, Plumb E, Metzger D, Blank MB. Rapid assessment of existing HIV prevention programming in a community mental health center. *Journal of Prevention and Intervention in the Community* 2007;33:137–151. [PubMed: 17298936]
- Strauss JL, Bosworth HB, Stechuchak KM, Meador KM, Butterfield MI. Knowledge and risks of human immunodeficiency virus transmission among veterans with severe mental illness. *Military Medicine* 2006;171:325–330. [PubMed: 16673748]
- Thompson SC, Checkley GE, Hocking JS, Crofts N, Mijch AM, Judd FK. HIV risk behaviour and HIV testing of psychiatric patients in Melbourne. *Australian and New Zealand Journal of Psychiatry* 1997;31:566–576. [PubMed: 9272267]
- Tucker JS, Kanouse DE, Min A, Koegel P, Sullivan G. HIV risk behaviors and their correlates among HIV-positive adults with serious mental illness. *AIDS and Behavior* 2003;7:29–40. [PubMed: 14534388]
- Uldall KK, Palmer NB, Whetten K, Mellins C. for the HIV/AIDS Treatment Adherence Health Outcomes and Cost Study Group. Adherence in people living with HIV/AIDS, mental illness, and chemical dependency: A review of the literature. *AIDS Care* 2004;16:S71–S96. [PubMed: 15736823]
- Walkup J, McAlpine DD, Olfson M, Boyer C, Hansell S. Recent HIV testing among general hospital inpatients with schizophrenia: Findings from four New York City sites. *Psychiatric Quarterly* 2000;71:177–193. [PubMed: 10832159]
- Walkup J, Satriano J, Berry D, Sadler P, Cournos F. HIV testing policy and serious mental illness. *American Journal of Public Health* 2002;92:1931–1939. [PubMed: 12453811]
- Walkup J, Satriano J, Hansell S, Olfson M. Practices related to HIV risk assessment in general hospital psychiatric units in New York state. *Psychiatric Services* 1998;49:529–530. [PubMed: 9550247]
- Weinhardt LS, Carey MP, Carey KB. HIV-risk behavior and the public health context of HIV/AIDS among women living with a severe and persistent mental illness. *Journal of Nervous and Mental Disease* 1998;186:276–282. [PubMed: 9612444]
- Weinhardt LS, Carey MP, Johnson BT, Bickham NL. Effects of HIV counseling and testing on sexual risk behavior: A meta-analytic review of published research, 1985–1997. *American Journal of Public Health* 1999;89:1397–1405. [PubMed: 10474559]
- Weinhardt LS, Otto-Salaj LL, Brondino MJ, Norberg MM, Kalichman SC. Sex-related alcohol expectancies predict sexual risk behavior among severely and persistently mentally ill adults. *Psychology of Addictive Behaviors* 2002;16:64–67. [PubMed: 11934088]

Table 1
HIV Testing Prevalence Among Individuals with a Severe Mental Illness

Authors	N	Gender	Primary Diagnoses	Substance Use	Setting	Timeframe for Testing	% HIV tested
Blumberg and Dickey (2003)	1834	not reported	major depression, bipolar, or schizoaffective disorder	not reported	national survey of noninstitutionalized adults	past 12 months lifetime	17.6% 46.5%
Desai and Rosenheck (2004)	5890	61% male	major depression, schizophrenia, other psychoses, personality disorder, bipolar disorder, and anxiety disorder	42.8% reported alcohol abuse; 38.4% reported drug abuse	homeless adults recruited through outreach and enrolled in intensive case management	past 3 months	38%
Desai, Rosenheck, and Desai (2007)	487	100% male	schizophrenia, bipolar disorder, major depression with psychotic features, or psychosis not otherwise specified	not reported	outpatient settings (VA medical center and two state mental health centers)	lifetime	58.1%
Goldberg (2004)	170	65% male	schizophrenia	not reported	outpatient	lifetime	69.4%
Goldberg, Himelhoch, Kreyenbuhl, Dickerson, Hackman, Fang, Brown, Wohlheiter, and Dixon (2005)	200	48% male	schizophrenia spectrum disorder or major mood disorder	25% used alcohol, past month; 11% used drugs, past month	outpatient	lifetime	59.5%
Grassi, Pavanati, Cardelli, Ferri, and Peron (1999)	91	65% male	schizophrenia	43% reported using drugs in the past 10 years	inpatient and outpatient psychiatric facilities in Italy	lifetime	36.7%
Katz, Watts, and Santman (1994)	54	74% male	schizophrenia or bipolar disorder	about one third had a drug problem	outpatient	lifetime	61%
Kelly, Murphy, Bahr, Brasfield, Davis, Hauth, Morgan, Stevenson, and Eilers (1992)	60	53% male	schizophrenia, affective disorder, or severe borderline personality	in past month, 54% used alcohol, 12% used marijuana, 5% used crack, 7% used illicit pills, and 2% used intravenous drugs	outpatient	lifetime	33%
Knox, Boaz, Friedrich, and Dow (1994)	120	54% male	schizophrenia, major depressive disorder, schizoaffective disorder, bipolar disorder, mood	one-third had a substance use disorder	inpatient and partial hospitalization	past 12 months lifetime	47% 64%

Authors	N	Gender	Primary Diagnoses	Substance Use	Setting	Timeframe for Testing	% HIV tested
Levounis, Galanter, Dermatis, Hamowy, and De Leon (2002)	104	100% male	disorder not further specified, dysthymia, and paranoid personality disorder	100% co-occurring substance use disorder	therapeutic community for homeless, chemically addicted men with a mental illness	lifetime	89%
Meade and Sikkema (2005)	150	54% male	schizophrenia, schizoaffective disorder, bipolar disorder, major depressive disorder, posttraumatic stress disorder	47% had a co-occurring substance use disorder	community mental health center, intensive outpatient program supportive housing programs, and a crisis/respice center	past 12 months lifetime	41% 81%
Miller and Finnerty (1996)	44	100% women	schizophrenia and schizoaffective disorder	78% reported drug use during pregnancy	inpatient and outpatient psychiatric facilities	lifetime	10.9%
Pirl, Greer, Weissgarber, Liverant, and Safren (2005)	655	70% male (in subset of patients on whom additional information was obtained)	schizophrenia and bipolar disorder (in subset of patients on whom additional information was obtained)	not reported	state psychiatric hospital	while hospitalized	9.5%
Thompson, Checkley, Hocking, Crofts, Mich, and Judd (1997)	145	60% male	schizophrenia, bipolar disorder, and depression	58% reported ever using illegal drugs	inpatient and outpatient psychiatric facilities in Melbourne, Australia	lifetime	34.5%
Walkup, McAlpine, Olifson, Boyer, and Hansell (2000)	300	64% male	schizophrenia or schizoaffective disorder	52% reported alcohol abuse; 54% reported drug abuse	inpatient	past 6 months	17%
Weinhardt, Carey, and Carey (1998)	61	100% female	schizophrenia, schizoaffective disorder, bipolar disorder, major depressive disorder, other psychotic disorders, and pervasive developmental disorder	31% had been in drug or alcohol treatment	outpatient clinics and day treatment programs	lifetime	56%